

CASE REPORT / ПРИКАЗ БОЛЕСНИКА

A combination of acute and delayed contralateral epidural hematoma

Aleksandar Kostić¹, Aleksandar Veselinović², Vesna Nikolov¹, Ivan Stefanović¹, Radisav Mitić¹, Miša Radisavljević¹, Slavko Živković¹

¹Niš Clinical Center, Clinic for Neurosurgery, Niš, Serbia;

²Niš Clinical Center, Clinic for Ophthalmology, Niš, Serbia

SUMMARY

Introduction An acute bilateral extradural hematoma is an uncommon presentation of a traumatic head injury; however, it leads to higher mortality rate than an acute unilateral hematoma. A delayed epidural hematoma (DEDH) is a hematoma not present on the initial computed tomography (CT) scan but is found on a subsequent CT scan. While reviewing the literature, we could not find recently published papers considering supratentorial DEDH after primary operated contralateral epidural hematoma.

Case outline A comatose 14-year-old male patient with Glasgow Coma Scale score of 4 and the right mydriatic pupil on the side of the blunt trauma to the head was admitted to the intensive care unit after he had survived a traffic accident. The initial brain CT scan showed an acute temporoparietal epidural hematoma on the right side of the cranium, with impressive midline shift and bilateral linear skull fracture. Surgery was performed and an intracranial pressure (ICP) monitor was implanted, which showed increased values of ICP. A control brain CT scan performed within 24 hours showed a new contralateral occipitoparietal epidural hematoma. Another operation was performed. A second, control CT brain scan showed favorable findings. The patient was transferred after 25 days to the rehabilitation center, with the disability rating score of 11, which was reduced to 1 after three months.

Conclusion A contralateral DEDH is a life-threatening neurosurgical emergency case which can occur during the first 24 hours after decompressive craniectomy. Control CT scans should be performed one day after the operation in order to verify and treat DEDH timely. A high degree of vigilance and ICP monitoring is recommended in these cases, especially after surgical decompression.

Keywords: brain trauma; delayed bleeding; extradural hematoma; intracranial pressure monitoring



INTRODUCTION

An epidural hematoma (EDH) occurs when blood accumulates between the skull and the dura mater due to a severe cranial trauma. A typical location of these hematomas is the temporal one, as the most common cause of bleeding is a lesion of the middle meningeal artery. The pterion region, which overlies it, is relatively weak and prone to injury. Nevertheless, other locations of EDH are not so uncommon and are reported in 20–30% of cases [1]. It is usually found on the same side of the cranium that was impacted by the blow, but very rarely it can be due to a counter coup brain injury. Epidural blood clot, while expanding, strips the dura from tabula interna of the skull, forming a mass that causes brain shifting, and consequently compression on the cerebral blood vessels and cranial nerves. Usually, EDH grows within the first four to six hours, after which profound deterioration starts in the form of contralateral limb weakness, coma, and ipsilateral pupil fixation and dilatation.

This condition is reported in 1–3% of head injuries [2]; 15–20% of EDHs are fatal, with only 5% in children population [3, 4].

Intracranial EDH is considered to be the most serious complication of head injury, re-

quiring immediate diagnosis and surgical intervention. It usually occurs in young adults, and it is rare before the age of two and after the age of 60. Intracranial EDH may be acute (58%), subacute (31%) or chronic (11%) [5]. Bilateral EDH accounts for 2–10% of all acute EDHs in adults, being exceedingly rare in children [6].

A traumatic delayed epidural hematoma (DEDH) can be defined as a hematoma that is insignificant or not present on the initial computerized tomography (CT) scan made after the trauma; however, a subsequent CT scan shows sizeable epidural bleeding [7]. The reported incidence of DEDH varies from 5.6% to 13.3% [8, 9].

While reviewing the literature, we could not find recently published papers considering supratentorial DEDH after primary operated contralateral EDH. We only found cases after the operation of a contralateral subdural or intracerebral hematoma [10, 11, 12].

CASE REPORT

Herein we present a case of a 14-year-old boy who sustained severe injuries as a pedestrian in a traffic accident. His polytrauma injuries included the left lung contusion, extensive

Received • Примљено:

March 16, 2017

Accepted • Прихваћено:

June 23, 2017

Online first: June 27, 2017

Correspondence to:

Aleksandar KOSTIĆ
Romanijska 11/25
18000 Niš
Serbia
aleko018@yahoo.co.uk

parietal bilateral skull fracture, and a typically located large EDH, temporoparietal on the right side, detected on a CT scan (Figure 1). On admission, the patient's Glasgow Come Scale score was 4, he had the left-sided hemiplegia, and the right pupil was fixed and dilated.

Urgent craniotomy was performed, as well as the evacuation and hemostasis of massive extradural hematoma combined with modest dura and brain laceration. An intracranial pressure (ICP) monitor was implanted in the subdural space, and it showed the values that indicated intracranial hypertension. The initial ICP was higher than 40 mmHg during the first postoperative hour and declined to around 24 (the average ICP value was 24.3 mmHg) in the next 20 hours. The right pupil shrank to the normal size. During the first 12 hours, the patient was sedated so that his state of consciousness was unavailable, except for the short period of time between two sedations. Due to ICHT and reduced possibilities of assessing Glasgow Come Scale score, a control CT brain scan was performed. A new, contralateral occipitoparietal DEDH was found, which was a bit smaller than the initial one (Figure 2). Another operation was performed, and under the contralateral fracture an EDH was evacuated. In the next three days, the values of ICP measured around 20 mmHg. The average daily ICP values were 19.4, 20.8, and 18.2, respectively, and in the last three days of ICP monitoring the values were around 15 mmHg. Meanwhile, blood samples routinely obtained in the emergency room were analyzed for coagulation parameters, including prothrombin time, platelet counts, activated partial thromboplastin time, fibrinogen level, and the international normalized ratio. All values were normal. One day after the second operation, the brain CT scan showed favorable findings (Figure 3). Eye exam was done at the seventh postoperative day, but no signs of papilledema were found. Long-term recovery, tracheotomy, and intensive physical rehabilitation followed the second operation. The patient was discharged from the hospital after three weeks and transferred to the rehabilitation facility and had disability rating score of 11. Three months after the accident, his DRS was 1.

DISCUSSION

The question is whether a hematoma that is not present on the initial CT scan but is found on a subsequent CT scan represents an enlargement of an EDH from invisible to visible on high-quality CT or is a newly-formed subacute extradural hemorrhage promoted by some risk factors.

The risk factors responsible for the delayed appearance of the EDH "tamponade" effect are usually increased endocranial pressure and post-traumatic arterial hypotension, as well as coagulopathy in a limited number of cases [13, 14, 15]. Post-traumatic nasal and ear leakage of the cerebrospinal fluid may result in the occurrence of DEDH. Some of the risk factors are iatrogenic, including the operation of cerebellar expansive lesions, rapid intravenous infusion or transfusion after the development of hemodynamic shock in a traumatized patient, intraven-




Figure 1. Initial epidural hematoma in the right temporoparietal region




Figure 2. Contralateral delayed epidural hematoma in the left occipitoparietal region – a hematoma newly formed after the first operation




Figure 3. A control brain CT scan after the second operation – normal postoperative finding

tricular cerebrospinal fluid drainage, overdrainage during ventriculoperitoneal shunting. A case of a DEDH in the spinal region, after facet joint infiltration in a chronic back pain patient, has also been described [16].

Some authors suggest repeating a brain CT scan within the first 36 hours after sustaining an injury in patients with a small, asymptomatic EDH, as the mean time to EDH enlargement is about eight hours [17]. We performed the control CT brain scan within 24 hours as the neurological status of the patient remained unclear and potentially unfavorable. Nevertheless, ICP monitoring and continuous neurological monitoring could be crucial in the early discovering of DEDH. Post-traumatic restlessness may lead to hyperventilation and blood pressure elevation with lowering of the ICP, both of which may provoke bleeding into EDH. However, in the series of Sakai et al. [18], all of the ICP-monitored patients had either unchanged or increased ICP. Early diagnosis of DEDH can be facilitated by the liberal use of intracranial pressure monitoring as its

sensitivity could be up to 80% [19, 20]. In the case we presented, after the first evacuation of the epidural clot, ICP dropped from 40 mmHg to 24 mmHg and that event could have triggered the subsequent bleeding from a fractured skull contralateral. Still, the ICP values remained higher than normal (average ICP > 20 mmHg) until the time of the second evacuation of the epidural clot, although normalization of ICP values was to be expected. It turned out that DEDH maintained the intracranial hypertension. The ICP values were lower or equal to 20 mmHg for the next three days. Normalization of ICP occurred on the fourth postoperative day. Therefore, ICP monitoring can be of great help, especially in a continuously sedated patient.

Eye examination did not show any signs of the papilloedema since it was performed a week after the accident. Normal results were to be expected according to the fundoscopic examination dynamic finding after severe brain trauma [21]. This dynamic is quite different from fundoscopic examination finding after severe spine injury, in which papilloedema occurs much later [22].

Enlargement of EDH in its acute phase is often explained by continuous hemorrhage and re-hemorrhage from either the arterial or venous vessels. Some authors emphasize that venous hemorrhage does not generate enough pressure to strip dura mater from the bone

or to overcome the clot-induced tamponade effect [23]. A decrease in clot density combined with the formation of membranes with permeable sinusoids usually after the fifth day after the trauma could lead to a new hemorrhage and further enlargement of the hematoma [24]. This slight increase in size between days 5 and 16 is manifested in 50% of cases, and some patients require emergency craniotomy when signs of herniation occur [24]. DEDH in our patient occurred much earlier, in the first 24 hours, probably due to the extensive bilateral fracture.

Rarely reported in mild head traumas [25], DEDH is commonly associated with skull fractures [20]. DEDH development in the infratentorial compartment is tenfold higher in patients with fracture line in the posterior fossa than in patients with supratentorial fractures [26]. Rapid deterioration of an injured patient, especially if DEDH is formed in the posterior fossa, could be lethal despite urgent CT brain scan and prompt operation [25]. This is explained by an irreversible damage to some of the vital centers located in the brain stem.

Atypical occipitoparietal location and modest volume of DEDH presented in this case report are relevant for the patient's prognosis, as it is unlikely to cause brain herniation or pressure on the brain stem. However, if it remains unrecognized timely, DEDH can be life-threatening.

REFERENCES

- Graham DJ, Gennarelli TA. Chapter 5. Pathology of Brain Damage After Head Injury. In: Cooper P, Golfinos G. Head Injury. 4th ed. New York: Morgan Hill; 2000.
- Mishra A, Mohanty S. Contre-coup extradural haematoma: a short report. *Neurol India*. 2001; 49(1):94–5.
- Sanders MJ, McKenna K. Head and Facial Trauma. Chapter 22. In: Mosby's Paramedic Textbook, 2nd revised ed. New York: Mosby; 2001.
- Binder H, Majdan M, Tiefenboeck TM, Fochtmann A, Michel M, Hajdu S, et al. Management and outcome of traumatic epidural hematoma in 41 infants and children from a single center. *Orthop Traumatol Surg Res*. 2016; 102(6):769–74.
- Dähnert W. Radiology Review Manual 7th edition. Epidural Hematoma of Brain. Philadelphia: Lippincott Williams&Wilkins; 2012. p. 289.
- Dharker SR, Bhargava N. Bilateral epidural haematoma. *Acta Neurochir (Wien)*. 1991; 110(1-2):29–32.
- Radulovic D, Janosevic V, Djurovic B, Slavik E. Traumatic delayed epidural hematoma. *Zentralbl Neurochir*. 2006; 67(2):76–80.
- Oertel M, Kelly DF, McArthur D, Boscardin WJ, Glenn TC, Lee JH, et al. Progressive hemorrhage after head trauma: predictors and consequences of the evolving injury. *J Neurosurg*. 2002; 96(1):109–16.
- Ashkenazi E, Constantini S, Pomeranz S, Rivkind AI, Rappaport ZH. Delayed epidural hematoma without neurologic deficit. *J Trauma*. 1990; 30(5):613–5.
- Su TM, Lee TH, Chen WF, Lee TC, Cheng CH. Contralateral acute epidural hematoma after decompressive surgery of acute subdural hematoma: clinical features and outcome. *J Trauma*. 2008; 65(6):1298–302.
- Shen J, Pan JW, Fan ZX, Zhou YQ, Chen Z, Zhan RY. Surgery for contralateral acute epidural hematoma following acute subdural hematoma evacuation: five new cases and a short literature review. *Acta Neurochir (Wien)*. 2013; 155(2):335–41.
- Solomichuk VO, Drizhdov KI. Contralateral delayed epidural hematoma following intracerebral hematoma surgery. *Surg Neurol Int*. 2013; 4:134.
- Domenicucci M, Signorini P, Strzelecki, Delfini R. Delayed post-traumatic epidural hematoma. A review *Neurosurg Rev*. 1995; 18(2):109–22.
- Brown MW, Yilmaz TS, Kasper EM. Iatrogenic spinal hematoma as a complication of lumbar puncture: What is the risk and best management plan? *Surg Neurol Int*. 2016; 7(Suppl 22):S581–9.
- Takahashi Y, Sato T, Hyodo H, Kawamata T, Takahashi E. Symptomatic epidural haematoma after cervical laminoplasty: a report of three cases. *J Orthop Surg (Hong Kong)*. 2016; 24(1):121–4.
- Velickovic M, Ballhause TM. Delayed onset of a spinal epidural hematoma after facet joint injection. *SAGE Open Med Case Rep*. 2016; 4:2050313X16675258.
- Sullivan TP, Jarvika JG, Cohen WA. Follow-up of Conservatively Managed Epidural Hematomas: Implications for Timing of Repeat CT. *AJNR*. 1999; 20:107–13.
- Sakai H, Takagi H, Ohtaka H, Tanabe T, Ohwada T, Yada K. Serial changes in acute extradural hematoma size and associated changes in level of consciousness and intracranial pressure. *J Neurosurg*. 1988; 68(4):566–70.
- Borovich B, Braun J, Guilburd JN, Zaaroor M, Michich M, Levy L, et al. Delayed onset of traumatic extradural hematoma. *J Neurosurg*. 1985; 63(1):30–4.
- Poon WS, Rehman SU, Poon CY, Li AK. Traumatic extradural hematoma of delayed onset is not a rarity. *Neurosurgery*. 1992; 30(5):681–6.
- Joshua SP, Agrawal D, Sharma BS, Mahapatra AK. Papilloedema as a non-invasive marker for raised intra-cranial pressure following decompressive craniectomy for severe head injury. *Clin Neurol Neurosurg*. 2011; 113(8):635–8.
- Catz A, Appel I, Reider-Grosswasser I, Grosswasser Z, Mendelson L, Gepstein R. Late-onset papilloedema following spinal injury. Case report. *Paraplegia*. 1993; 31(2):131–5.
- Bender MB, Christoff N. Nonsurgical treatment of subdural hematomas. *Arch Neurol*. 1974; 31(2):73–9.
- Pang D, Horton JA, Herron JM, Wilberger JE, Vries JK. Nonsurgical management of extradural hematomas in children. *J Neurosurg*. 1983; 59(6):958–71.
- Resigo P, Piquer J, Bottela C, Orozco M, Navarro J, Cabanes J. Delayed extradural hematoma after mild head injury: report of three cases. *Surg Neurol*. 1997; 48(3):226–31.
- Kircelli A, Özel Ö, Can H, Sari R, Cansever T, Elmali İ. Is the presence of linear fracture a predictor of delayed posterior fossa epidural hematoma? *Ulus Travma Acil Cerrahi Derg*. 2016; 22(4):355–60.

Комбинација акутног и одложеног контралатералног епидуралног хематома

Александар Костић¹, Александар Веселиновић², Весна Николов¹, Иван Стефановић¹, Радисав Митић¹, Миша Радисављевић¹, Славко Живковић¹

¹Клинички центар Ниш, Клиника за неурохирургију, Србија;

²Клинички центар Ниш, Клиника за офталмологију, Србија

САЖЕТАК

Увод Акутни билатерални епидурални хематоми (ЕДХ) нису чест налаз после повреде главе, али су узрок знатно веће смртности у поређењу са акутним унилатералним хематомом. Одложени епидурални хематом (ОЕДХ) епидурално је крварење које није присутно на иницијалној компјутеризованој томографији (КТ), али је нађено на накнадној КТ. У литератури нисмо нашли рад са описом супратенторијалног ОЕДХ после операције контралатералног ЕДХ.

Приказ случаја После саобраћајне несреће примљен је 14 година стар дечак у коми (Глазгов кома скор 4) и са проширеном зеницом на страни тупе трауме главе. Иницијална КТ мозга показала је темпоропаријетални ЕДХ на десној страни кранијума, са израженим померењем средњелинијских структура и билатералном линераном фрактуром лобање.

Повређени је оперисан и започет је мониторинг интракранијалног притиска (ИКП), који је био повишен. Контролна КТ мозга унутар прва 24 часа показала је нови контралатерални окципито-паријетални ЕДХ. Предузета је нова операција. Други, контролни налаз КТ мозга је био уредан. После 25 дана болесник је преведен у Рехабилитациони центар, са степеном неспособности оцењеним са 11. После три месеца болесник је био готово без последица повређивања.

Закључак Контралатерални ОЕДХ је стање непосредне животне угрожености, који се може појавити унутар прва 24 часа од краниотомије. Потребно је брижљиво праћење стања болесника, укључујући контролну КТ и праћење ИКП.

Кључне речи: траума мозга; одложено крварење; епидурални хематом; мерење интракранијалног притиска