

REVIEW ARTICLE / ПРЕГЛЕД ЛИТЕРАТУРЕ

Prevalence of spondyloarthritis and its subtypes – are they really comparable?

Mirjana I. Zlatković-Švenda^{1,2}, Roksanda M. Stojanović¹, Sandra B. Šipetić-Grujičić³, Marija M. Radak-Perović¹, Francis Guillemin⁴

¹University of Belgrade, Faculty of Medicine, Institute of Rheumatology, Belgrade, Serbia;

²University of East Sarajevo, Faculty of Medicine, East Sarajevo, Republic of Srpska, Bosnia and Herzegovina;

³University of Belgrade, Faculty of Medicine, Institute of Epidemiology, Belgrade, Serbia;

⁴University of Lorraine, University Hospital, INSERM-CIC-EC and EA 4360 APEMAC, Nancy, France



SUMMARY

Introduction/Objective Increasing spondyloarthritis (SpA) prevalence in the last several decades cannot be attributed to disease manifestations alone.

The objective of this paper is to review the prevalence of SpA and its subtypes: ankylosing spondylitis (AS), psoriatic arthritis (PsA), reactive arthritis (ReA), SpA related to inflammatory bowel disease (IBD) and undifferentiated SpA (UnSpA).

Methods MEDLINE literature search was done via PubMed, Google Scholar, and Embase databases, using terms for spondyloarthritis, and prevalence, with an additional hand searching.

Results As compared with southern European countries, northern European countries (Scotland, Sweden, France) showed lower SpA prevalence rates (0.21–0.45% vs. 1.06% and 1.35% in Italy and Turkey, respectively). The lowest world SpA prevalence was in African and Southeast Asian countries (0–0.19%), and the highest was in Alaska (2.5%). The widest variability in PsA prevalence was in Europe (northern 0.02–0.19%, southern 0.42%). The lowest world PsA prevalence was in Japan (0.001%), followed by China (0.01–0.10%). The European ReA prevalence ranged from 0.04% in Greece to 0.10% in Serbia and Germany, and the European UnSpA prevalence varied from 0.02% in Serbia to 0.67% in Germany; the highest world UnSpA prevalence was in Lebanon (3.4%). Studies aimed at estimating the SpA prevalence differed in sampling strategy and confirmation criteria, different cutoffs for age groups inclusion, presentation of standardized or raw results, etc.

Conclusion Variation in the SpA prevalence cannot be attributed to genetic or geographic distribution only. Differences in methodology of studies add to the diversification, described more in-depth in this review.

Keywords: epidemiology; prevalence; spondyloarthritis; ankylosing spondylitis; psoriatic arthritis

INTRODUCTION

Spondyloarthritis (SpA) comprises ankylosing spondylitis (AS), psoriatic arthritis (PsA), reactive arthritis (ReA), SpA related to inflammatory bowel diseases (IBD), such as Crohn's disease or ulcerative colitis, and undifferentiated SpA (UnSpA). Common SpA clinical features are spinal, sacroiliac, and peripheral joint inflammation, enthesitis and some extra-articular manifestations (uveitis, skin or mucosal changes, cardiovascular or pulmonary changes).

SpA prevalence estimates range widely both in developed and developing countries, from 0.03% in the APLAR region (New Zealand in Oceania to Jordan in the Middle East) to 2.5% in Russian and Alaskan Eskimos [1, 2, 3] (Figure 1). Particularly, African black people populations have shown virtually non-existent SpA prevalence, as in Sub-Saharan Africa, Zimbabwe, and Nigeria [4, 5].

Differences in the prevalence of SpA and its subtypes, apart from geographic, racial, and cultural influences, could also be due to differences in the methodology used in studies (various sampling and case identification

strategy with different classification criteria, diverse age groups included, presentation of raw or standardized results, etc.). For example, in the last decades, PsA has been identified by a number of different criteria [6, 7].

This study aimed to review the estimates of prevalence of SpA and its subtypes: AS, PsA, ReA, IBD SpA, and UnSpA, specifically emphasizing differences in methodology and variations in geography.

METHODS

Review involved a search of MEDLINE literature via PubMed, Google Scholar, and Embase databases with the terms spondyloarthritis, spondyloarthropathy, ankylosing spondylitis, psoriatic arthritis, reactive arthritis, inflammatory bowel disease spondyloarthritis, undifferentiated spondyloarthritis and prevalence, with an additional hand search. A total of 15,325 titles and abstracts were reviewed; the full text of 292 articles was read thoroughly; and data for 67 reports are presented and compared here.

Received • Примљено:

January 10, 2018

Revised • Ревизија:

December 3, 2018

Accepted • Прихваћено:

January 24, 2019

Online first: February 5, 2019

Correspondence to:

Mirjana ZLATKOVIĆ-ŠVENDA
University of Belgrade
Faculty of Medicine
Institute of Rheumatology
Resavska 69
11000 Belgrade, Serbia

mirjanazlatkovicsvenda@gmail.com

Table 1. Spondyloarthritis prevalence

Reference	Country	Year	Age	Population, area, Number of participants	Criteria	Prevalence [%]		
						Men	Women	All
EUROPE								
Steven [14]	Scotland	1992	15+	Mixed, Scottish islands	Doctor dg			0.21*
Saraux [15]	France	2001	18+	Mixed, France 9,395	Doctor dg	0.31	0.29	0.30*
Constantino [16]		2010	35+	Gazel Cohort, 18,757	ASAS			0.43*
Zlatković-Švenda [17]	Serbia	2008	18+	Mixed, Serbia 6,213	Doctor dg	0.34	0.31	0.32*
Haglund [18]	Sweden	2011	15+	Mixed, south Sweden	Register ^p			0.45
Trontzas [19]	Greece	2005	19+	Mixed, 7 areas, 10,647	ESSG			0.49*
Adomavičiute [20]	Lithuania	2008	18+	Urban, Lithuania, 6,542	Doctor dg	1.36	1.34	0.84*
De Angelis [21]	Italy	2007	18+	Mixed, Marche, 3,664	ESSG			1.06
Bruges-Armas [22]	Portugal	2002	50+	Mixed, Terceira, Azores, 936	ESSG	2.7	0.4	1.6
Braun [23]	Germany	1998	18+	Blood donors, 1,871	ESSG			1.9
AFRICA								
Adebajo [4]	Sub-Saharan Africa	1994	A	Negro, rural	Doctor dg			0.00
Mijiyawa [5]	Togo	2000	A	Negro, rural	Doctor dg			0.00
APLAR region [south-east Asia and Pacific]								
Davatachi [1]	Phillipines	1997	15+	Urban, Manila, 3,006	Doctor dg			0.03
Akhter [6]	Pakistan	2011	15+	Rural, 2,090	/			0.10
Veerapen [7]	Malasya	1992	15+	Rural, Banting, 2,594	Doctor dg			0.12
Davatachi [1]	Tailand	1998	15+	Rural, 2,463	/			0.12
Dai [9]	China	2003	15+	Urban, Shanghai, 7,603	Doctor dg			0.11
Zeng [10]		1995	16+	Urban, Shantou 2,040		0.26		
Davatachi [1]	Iran	1995	15+	Urban, 7,000	/			0.34
Onen [12]	Turkey	2008	20+	Urban, Izmir, 2,887	ESSG	0.88	1.22	1.05*
Onen [13]		2015		University Employees, 381	ASAS			1.35
Davatachi [1]	APLAR	2006	15+	APLAR	/			0.19*
Alexeeva [2]	Russia	1994	P	Chukotka, Russia				2.5
NORTH AMERICA								
Strand [24]	USA	2013	18+	Mixed, pt.records, 816	ASAS			0.70
Helmick [25]	USA	2005	18+	NHIS	Doctor dg			0.35–1.31
Boyer [3]	Alaska	1994	20+	Eskimos, 6,749	ESSG/AMOR			2.50

*Prevalence estimates standardized for age and sex in terms of the country population; Doctor dg, diagnosis made by a doctor; ESSG – European Spondyloarthropathy Study Group Classification criteria; ASAS – Assessment of SpondyloArthritis international Society criteria; AMOR – Spondyloarthropathy criteria by Amor et al; A – adult; P – population

SPONDYLOARTHRITIS PREVALENCE

The Asian SpA prevalence ranged from 0.03% in Manila to 1.05–1.3% in Turkey [1, 8–13]. The SpA standardized prevalence in the whole APLAR region was 0.19% [1]. The SpA prevalence in Europe ranged from 0.21% in Scotland to 1.9% in Germany [14–23]. In Serbia, it was 0.32% [17] (Table 1). The estimated SpA prevalence in the United States was 0.35–1.31% [24, 25]. The Australian urban SpA prevalence was 0.21%, but was 0.50% in Australian Aboriginals [1].

Comparison of the world SpA prevalence was limited by methodology discrepancies: various sample sizes, different age limitations (15, 16, 18, 20 years), and different presentation of the achieved results (row or standardized) (Table 1). However, the prevalence was lower in tropical zone countries [Africa or South Asia and Southeastern Asia (Philippines, Pakistan, Malaysia, Thailand, China)] than in northern countries, with the highest rate in Alaska (Figure 1) [2, 4, 7–10]. However, prevalence was lower in northern Europe (Scotland, Sweden, France) than in

southern Europe (Italy and Turkey) (Figure 1), which contrasted with the reported rheumatoid arthritis prevalence [12, 13, 14, 16, 18, 21, 26]. Inequalities in methodologies of SpA studies are described in Table 1.

ANKYLOSING SPONDYLITIS PREVALENCE

A wide range of AS prevalence was reported as well (Table 2, Figure 2). The highest AS prevalence was found in Canadian Indigenous Haida people, 6% [27]. However, no AS cases were found in African countries: Zimbabwe, Nigeria, and Togo [28, 29, 30]. For Southeast Asian and Pacific regions, AS prevalence estimates ranged from 0.01% in Japan to 0.49% in Turkey [12, 31, 32, 33] (Table 2). The European AS prevalence ranged from 0.08% in France and Serbia to 0.86% in Germany [23] [15, 17, 18, 21, 22, 34–40] and was 1.1–1.4% in Norway (Table 2, Figure 2). The AS population prevalence in the United States was estimated at 0.35% and 0.52% (e.g., 0.13% for Caucasians only [24, 25, 41]). The Alaskan Eskimo AS prevalence was 0.4% [3].



Figure 1. Spondyloarthritis prevalence in the world given in percentages

NOTE: The specific population estimate may not represent the full population estimate for a particular country (see Table 1)

AS prevalence estimates were lower in African and South and Southeast Asian countries than in northern Europe and the United States [25, 27–33, 40] (Figure 2). Differences in the studies are further evaluated in Table 2.

PSORIATIC ARTHRITIS PREVALENCE

The widest variability in SpA prevalence was for PsA and was the highest in Europe: 0.42% in Italy [21] (Figure 3). As one of the SpA-related diseases, PsA was not found in certain African countries either (in accordance with the virtual non-existence of SpA), and the non-existence of PsA was confirmed by the studies from Uganda and Sub-Saharan Africa [29, 42]. PsA prevalence was low in Southeast Asia and Pacific regions as well: 0.001% in Japan and 0.10% in China [43, 44]. Europe had the widest PsA prevalence range, lower in the northern than southern part, ranging from 0.02% in Sweden to 0.42% in Italy [15, 17, 21, 23, 34, 36, 37, 45–50] (Figure 3, Table 3). The PsA prevalence was 0.09% in Serbia. The prevalence ranged from 0.07–0.25% in the United States and was 0.12% in Minnesota and 0.10% in Alaskan Eskimos [3, 41, 51, 52]. The prevalence in South America was 0.07% [53].

Differences in PsA prevalence studies are described in more detail in Table 3.

REACTIVE ARTHRITIS PREVALENCE

Results of the ReA prevalence estimation studies cannot be generalized because of periodic and geographic variations in “disease trigger factors,” such as *Salmonella*, *Shigella*, *Yersinia*, and *Campylobacter* bacteria species, as well as *Chlamydia* and *Mycoplasma* subspecies. In fact, only a few ReA prevalence studies were performed, mostly under the SpA disease prevalence estimation. The ReA prevalence was 0.04% in Greece, 0.09% in Italy and 0.10% in Serbia and Germany [17, 19, 21, 23].

INFLAMMATORY BOWEL DISEASE SPONDYLOARTHRTIS PREVALENCE

The prevalence of IBD SpA (Crohn’s disease or ulcerative colitis) has not been studied much, and was less frequently reported than IBDs themselves. According to the few reported studies, the IBD SpA prevalence ranged from 0.01% in Sweden to 0.03% in Serbia, 0.04% in Greece and 0.09% in Italy [17, 18, 19, 21]. It was 0.05–0.25% in the United States [25].

Prevalence related to IBD type was evaluated by one US study: the prevalence of ulcerative colitis SpA was 0.03% and Crohn’s disease 0.01% [41]. Karreman et al. [54] found a SpA prevalence of 13% in IBD patients, with the pooled prevalence for AS of 3%, peripheral arthritis 13% and sacroiliitis 10%.

Table 2. Ankylosing spondylitis prevalence

Reference	Country	Year	Age	Method	Prevalence [%]
EUROPE					
Saraux [15]	France	2001	18+	Questionnaire	0.08*
Zlatković-Švenda [17]	Serbia	2008	18+	Questionnaire	0.08*
Hanova [34]	Czech Republic	2003	16+	Register	0.09
Haglund [18]	Sweden	2011	15+	Register	0.12
Kaipainen [35]	Finland	1997	30+	Sample	0.15
Trontzas [36]	Greece	2005	19+	Questionnaire	0.24*
Anagnostopoulos [37]		2010	A	Questionnaire	0.29
Alamanos [38]		2002	16+	Register	0.29*
Backland [39]	Norway	2005	20+	Register Urban	0.26
Gran [40]		1985			1.1–1.4
De Angelis [21]	Italy	2007	18+	Questionnaire	0.37
Bruges-Armas [22]	Portugal	2002	50+	Questionnaire	0.6
Braun [23]	Germany	1998	18+	Blood donors	0.86
AFRICA					
Stein [28]	Zimbabwe	1990		Population	0.00
Adebajo [29]	Nigeria	1991		Population	0.00
Mijiyawa [30]	Togo	1993		Population	0.00
APLAR region (south-east Asia and Pacific)					
Hukuda [31]	Japan	2001	A	Register	0.01
Dai [10]	Shangai	2003		Urban	0.11*
		1985	16+	Rural	0.22
		1995	16+	Urban	0.2
Yao-Jun [32]	China	2005	16+	Urban	0.2*
Chou [33]	Taiwan	1994	20+	Urban	0.4
				Suburban	0.19
				Rural	0.54
Onen [12]	Turkey	2008	20+	Urban	0.49*
Alexeeva [2]	Chukotka, Russia	1994	P	Population	1.1
NORTH AMERICA					
Boyer [3]	Alaska	1994	20+	Register	0.4
Strand [24]	USA	2013	18+	Pt records	0.35
Helnick [25]		2005	18+	NHIS	0.52*
Lawrence [41]	USA, Caucasians	1998	A	Sample	0.13*
Gofton [27]	Canada, Haida People	1964	15+	Population	6.2

*Prevalence estimates standardized for age and sex in terms of the country population; P – population; A – adult

UNDIFFERENTIATED SPONDYLOARTHRITIS PREVALENCE

The UnSpA impact and its clinical manifestations are not yet fully understood. In general, UnSpA should be considered in patients presented with inflammatory back pain or peripheral joint arthritis, enthesopathy or alternative buttock pain, not accompanied by some SpA-disease specific features such as spondylitis, psoriasis, Crohn's disease, ulcerative colitis, or previously proven urogenital or gastrointestinal infection.

The highest UnSpA prevalence was in Lebanon, 3.4% [55]. In Asia, the representative China had 0.24% prevalence [56]. In Europe, the UnSpA prevalence estimate for Serbia was 0.02%, Greece, 0.03%, France, 0.04%, Sweden, 0.10%, and Germany, 0.67% [15, 17, 18, 19, 23]. The UnSpA prevalence in the United States was 0.37% and in Alaskan Eskimos 1.3% [3, 25, 57].

DISCUSSION

Although historically considered sporadic cases of rheumatoid arthritis and reported at the beginning as “rheumatoid spondylitis” or “seronegative variants of RA,” AS and PsA have over time been shown not only as separate diseases, but also diseases of constantly increasing impact and importance. When the SpA concept was promoted for several seronegative diseases characterized by certain common features, thereby enabling different diseases to be classified into a unified diagnosis a few decades ago, they may have been classified as SpA.

SpA prevalence has increased in the past decades. With the universal health system in France, the AS prevalence was estimated at 0.005% in 1994, 0.08% in 2001 (French population study) and 0.43% in 2015 (French population-based study) [15, 16, 58].

The 21st century European studies showed high SpA prevalence as well: Sweden 0.45%, France 0.43%, Greece 0.49%, Lithuania 0.84%, Turkey 1.05%, Italy 1.06%, and Portugal 1.6%. In the last century, SpA prevalence even



Figure 2. Ankylosing spondylitis prevalence in the world given in percentages;

NOTE: The specific population estimate may not represent the full population estimate for a particular country (see Table 2)

Table 3. Psoriatic arthritis prevalence

Reference	Country	Year	Age	Method	Prevalence [%]
EUROPE					
Hellgren [45]	Sweden	1969	P		0.02
van Romunde [46]	Netherlands	1984	20+	Sample	0.05
Zlatković-Švenda [17]	Serbia	2008	18+	Questionnaire	0.09
Pedersen [47]	Denmark	2002	Twins	Questionnaire	0.15
Madland [48]	Norway	2002	A	Register	0.19*
Hanova [34]	Czech Republic	2003	16+	Register	0.05*
Sarau [15]	France	2001	18+	Questionnaire	0.19*
Braun [23]	Germany	1998	18+	Blood donors	0.29
Alamanos [49]	Greece	2001	A	Register	0.06*
Trontzas [37]		2005	19+	Questionnaire	0.17*
Anagnostopoulos [38]		2010	A	Questionnaire	0.35
Salaffi [50]	Italy	2005	18+	Questionnaire	0.37
De Angelis [21]		2007	18+	Questionnaire	0.42
AFRICA					
Adebajo [29]	Subsaharial Africa	1990			0.00
Lamurezo [42]	Uganda	1980			0.00
APLAR region (south-east Asia and Pacific)					
Hukuda [43]	Japan	2001	A	Register	0.001
Zeng [44]	China	2006	A	Study	0.01–0.1
NORTH AMERICA					
Lawrence [41]	USA white race	1998	A	Sample	0.07*
Gelfand [51]		2005	P	Questionnaire	0.25
Shbeeb [52]	Minesota	1991	A	Register	0.16*
Boyer [3]	Alaska	1994	20+	Register	0.1
SOUTH AMERICA					
Soriano [53]	Argentina	2006	P	Register	0.07

*Prevalence estimates standardized for age and sex in terms of the country population
A – adult; P – population



Figure 3. Psoriatic arthritis prevalence in the world given in percentages;

NOTE: The specific population estimate may not represent the full population estimate for a particular country (see Table 3)

exceeded rheumatoid arthritis (as one of the most prevalent inflammatory rheumatic diseases) in Japan, Germany, Turkey, Lithuania, China, Italy, Australian Aboriginals and the United States, but was similar to the rheumatoid arthritis prevalence in France and Serbia [17, 59]. This fact could be explained by a better understanding of the SpA concept both by the patients and physicians, as well as by introducing new classification criteria, the European Spondyloarthropathy Study Group Classification criteria (ESSG) [60]. The newest Assessment of SpondyloArthritis international Society (ASAS) classification criteria was used by several studies, reporting high SpA prevalence estimates as well: 0.43% in France (0.36% for axial SpA and 0.12% for peripheral SpA), 1.35% in Turkey and 0.70% for axial SpA alone in the United States [13, 16, 24, 61].

The first study on the prevalence of pooled SpA and its subtypes was conducted in 2016 by Stolwijk et al. [62]. The general population pooled prevalence of SpA ranged from 0.20% in Southeast Asia to 1.61% in northern Arctic communities; the AS prevalence ranged from 0.02% in Sub-Saharan Africa to 0.35% in northern Arctic communities; and the PsA prevalence ranged from 0.01% in the Middle East to 0.19% in Europe [62]. However, only a few studies from this meta-analysis were truly representative of the general population, due to a high risk of bias in addition to other methodological variations described more in depth in the article [62].

The limitation of our review is that we could not track the human leukocyte antigen B27 (HLA-B27) prevalence in all studies included, and connect its range with prevalence differences. According to the DESIR (*DEvenir des*

Spondylarthropathies Indifférenciées Récentes) cohort, diagnostic delays are less common in HLA-B27-positive than in B27-negative patients [63]. High HLA-B27 prevalence is generally positively correlated with SpA prevalence, especially axial SpA [64]. Our review found the highest SpA prevalence in Alaskan Eskimos, 2.5% [3], of whom 25–40% are HLA-B27-positive [64]. In the current review, second-place SpA prevalence was found in Russian Chukotka Eskimos, 2.5%, with HLA-B27 prevalence at 34%. The highest AS prevalence (6%) was found in Haida Indians living in the Queen Charlotte Islands of Canada, of whom 50% are HLA-B27-positive [27]. However, although the HLA-B27 prevalence varies from 10–16% in northern European countries, the latest studies have shown an AS prevalence of 0.12% in Sweden, 0.15% in Finland and 0.26% in Norway [18, 35, 39, 64].

CONCLUSION

In conclusion, although the wide SpA prevalence range in different geographic regions could partially be attributed in part to HLA-B27 positivity, other factors may also play the role, such as geographic differences [59]. In addition, prevalence variations could only partially be due to genetic or environmental factors or geographic distribution and are certainly affected by differences in case definition and case confirmation strategy as well as by other variations in applied methodologies [65, 66].

Conflict of interest: None declared.

REFERENCES

- Davatchi F. Rheumatic diseases in the APLAR region. *APLAR J Rheumatol.* 2006; 9:5–10.
- Alexeeva L, Krylov M, Vturin V, Mylov N, Erdesz S, Benevolenskaya L. Prevalence of spondyloarthropathies and HLA-B27 in the native population of Chukotka, Russia. *J Rheumatol.* 1994; 21(12):2298–300.
- Boyer GS, Templin DW, Cornoni-Huntley JC, Everett DF, Lawrence RC, Heyse SF, et al. Prevalence of spondyloarthropathies in Alaskan Eskimos. *J Rheumatol.* 1994; 21(12):2292–7.
- Mijiyawa M, Oniankitan O, Khan MA. Spondyloarthropathies in sub-Saharan Africa. *Curr Opin Rheumatol.* 2000; 12(4):281–6.
- Adebajo A, Davis P. Rheumatic diseases in African blacks. *Semin Arthritis Rheum.* 1994; 24(2):139–53.
- Zlatkovic-Svenda M, Kerimovic-Morina D, Stojanovic RM. Psoriatic arthritis classification criteria: Moll and Wright, ESSG and CASPAR – a comparative study. *Acta Reumatol Port.* 2013; 38(3):172–8.
- Zlatkovic-Svenda MI, Kerimovic-Morina D, Stojanovic RM. Psoriatic arthritis criteria evaluation: CASPAR and Modified CASPAR. *Clin Exp Rheumatol.* 2011; 29(5):899–900.
- Akhter E, Bilal S, Kiani A, Haque U. Prevalence of arthritis in India and Pakistan: a review. *Rheumatol Int.* 2011; 31(7):849–55.
- Veerapen K. Epidemiology of Rheumatic disease in Malaysia. In: AR Nasution, J Darmawan, H Isbagi, eds. *APLAR Rheumatology 1992*:397–9.
- Dai SM, Han XH, Zhao DB, Shi YQ, Liu Y, Meng JM. Prevalence of rheumatic symptoms, rheumatoid arthritis, ankylosing spondylitis, and gout in Shanghai, China: a COPCORD study. *J Rheumatol.* 2003; 30(10):2245–51.
- Zeng QY, Chen R, Darmawan J, Xiao ZY, Chen SB, Wigley R, et al. Rheumatic diseases in China. *Arthritis Res Ther.* 2008; 10(1):R17.
- Onen F, Akar S, Birlik M, Sari I, Khan MA, Gourler O, et al. Prevalence of ankylosing spondylitis and related spondyloarthritides in an urban area of Izmir, Turkey. *J Rheumatol.* 2008; 35(2):305–9.
- Onen F, Solmaz D, Cetin P, Sari I, Balci A, Birlik M, et al. Prevalence of Inflammatory Back Pain and Axial Spondyloarthritis among University Employees in Izmir, Turkey. *J Rheumatol.* 2015; 42(9):1647–51.
- Steven MM. Prevalence of chronic arthritis in four geographical areas of the Scottish Highlands. *Ann Rheum Dis.* 1992; 51(2):186–94.
- Sarax A, Guillemin F, Guggenbuhl P, Roux CH, Fardellone P, Le Bihan E, et al. Prevalence of spondyloarthropathies in France: 2001. *Ann Rheum Dis.* 2005; 64(10):1431–5.
- Constantino F, Talpin A, Said-Nahal R, Goldberg M, Henny J, Chiochia G, et al. Prevalence of spondyloarthritis in reference to HLA-B27 in the French population: results of the GAZEL cohort. *Ann Rheum Dis.* 2015; 74(4):689–93.
- Zlatkovic-Svenda M, Stojanovic R, Sipetic-Grujicic S, Radak-Perovic M, Damjanov N, Guillemin F. Prevalence of spondyloarthritis in Serbia: a EULAR endorsed study. *Ann Rheum Dis.* 2015; 74(10):1940–2.
- Haglund E, Bremander A, Petersson I, Strömbeck B, Bergman S, Jacobsson L, et al. Prevalence of spondyloarthritis and its subtypes in southern Sweden. *Ann Rheum Dis.* 2011; 70(6):943–8.
- Trontzas P, Andrianakos A, Miyakis S, Pantelidou K, Vafiadou E, Garantziotou V, et al. Seronegative spondyloarthropathies in Greece: a population-based study of prevalence, clinical pattern, and management. The ESORDIG study. *Clin Rheumatol.* 2005; 24(6):583–9.
- Adomaviciute D, Pileckyte M, Baranauskaite A, Morvan J, Dadoniene J, Guillemin F. Prevalence survey of rheumatoid arthritis and spondyloarthropathy in Lithuania. *Scand J Rheumatol.* 2008; 37(2):113–9.
- De Angelis R, Salaffi F, Grassi W. Prevalence of spondyloarthropathies in an Italian population sample: a regional community-based study. *Scand J Rheumatol.* 2007; 36(1):14–21.
- Bruges-Armas J, Lima C, Peixoto MJ, Santos P, Mendonca D, da Silva BM, et al. Prevalence of spondyloarthritis in Terceira, Azores: a population based study. *Ann Rheum Dis.* 2002; 61(6):551–3.
- Braun J, Bollow M, Remlinger G, Eggens U, Rudwaleit M, Distler A, et al. Prevalence of spondylarthropathies in HLAB27 positive and negative blood donors. *Arthritis Rheum.* 1998; 41(1):58–67.
- Strand V, Rao SA, Shillington AC, Cifaldi MA, McGuire M, Ruderman EM. Prevalence of axial spondyloarthritis in United States rheumatology practices: Assessment of SpondyloArthritis International Society criteria versus rheumatology expert clinical diagnosis. *Arthritis Care Res (Hoboken).* 2013; 65(8):1299–306.
- Helmick C, Felson D, Lawrence R, Gabriel S, Hirsch R, Kwoh K, et al. Estimates of the Prevalence of Arthritis and Other Rheumatic Conditions in the United States. *Arthritis Rheum.* 2008; 58(1):15–25.
- Zlatkovic-Svenda M, Stojanovic R, Sipetic-Grujicic S, Guillemin F. Prevalence of rheumatoid arthritis in Serbia. *Rheumatol Int.* 2014; 34(5):649–58.
- Gofton JP, Robinson HS, Trueman GE. Ankylosing spondylitis in a Canadian Indian population. *Ann Rheum Dis.* 1966; 25(6):525–7.
- Stein M, Davis P, Emmanuel J, West G. The spondyloarthropathies in Zimbabwe: A clinical and immunogenetic profile. *J Rheumatol.* 1990; 17(10):1337–9.
- Adebajo AO. Spondyloarthropathies in Sub-Saharan Africa. *J Rheumatol.* 1991; 18(7):1115.
- Mijiyawa M. Spondyloarthropathies in patients attending the rheumatology unit of Lom6 Hospital. *J Rheumatol.* 1993; 20(7):1167–9.
- Hukuda S, Minami M, Saito T, Mitsui H, Matsui N, Komatsubara Y, et al. Spondyloarthropathies in Japan: nationwide questionnaire survey performed by the Japan Ankylosing Spondylitis Society. *J Rheumatol.* 2001; 28(3):554–9.
- Yao-Jun Xiang, Sheng-Ming Dai. Prevalence of rheumatic diseases and disability in China. *Rheumatol Int.* 2009; 29(5):481–90.
- Chou C-T, Pei L, Chang D-M, Lee C-F, Schumacher HR, Liang MH. Prevalence of rheumatic diseases in Taiwan: a population study of urban, suburban, rural differences. *J Rheumatol.* 1994; 21(2):302–6.
- Hanova P, Pavelka K, Holcatova I, Pikhart H. Incidence and prevalence of psoriatic arthritis, ankylosing spondylitis, and reactive arthritis in the first descriptive population-based study in the Czech Republic. *Scand J Rheumatol.* 2010; 39(4):310–7.
- Kaipainen-Seppanen O, Aho K, Heliovaara M. Incidence and prevalence of ankylosing spondylitis in Finland. *J Rheumatol.* 1997; 24(3):496–9.
- Trontzas P, Andrianakos A, Miyakis S, Pantelidou K, Vafiadou E, Garantziotou V, et al. Seronegative spondyloarthropathies in Greece: a population-based study of prevalence, clinical pattern and management. The ESORDIG study. *Clin Rheumatol.* 2005; 24(6):583–9.
- Anagnostopoulos I, Zinzaras E, Alexiou I, Papathanasiou A, Davas E, Koutroumpas A, et al. The prevalence of rheumatic diseases in central Greece: a population survey. *BMC Musculoskel Disord.* 2010; 11:98.
- Alamanos Y, Papadopoulos NG, Voulgari PV, Karakatsanis A, Siozos C, Drosos AA. Epidemiology of ankylosing spondylitis in Northwest Greece, 1983–2002. *Rheumatology (Oxford).* 2004; 43(5):615–8.
- Backland G, Nossent HC, Gran JT. Incidence and prevalence of ankylosing spondylitis in Northern Norway. *Arthritis Rheum.* 2005; 53(6):850–5.
- Gran JT, Husby G, Hordvik M. Prevalence of ankylosing spondylitis in males and females in a young middle-aged population of Tromsø, northern Norway. *Ann Rheum Dis.* 1985; 44(6):359–67.
- Lawrence RC, Helmick CG, Arnett FC, Deyo RA, Felson DT, Giannini EH, et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum.* 1998; 41(5):778–99.
- Lamurezo BR. Psoriatic arthritis amongst Ugandan Africans. *East Afr Med J.* 1989; 57:855–60.
- Hukuda S, Minami M, Saito T, Mitsui H, Matsui N, Komatsubara Y, et al. Spondyloarthropathies in Japan: nationwide questionnaire survey performed by the Japan Ankylosing Spondylitis Society. *J Rheumatol.* 2001; 28(3):554–9.
- Zeng QY, Chen R, Darmawan J, Xiao ZY, Chen SB, Wigley R, et al. Rheumatic diseases in China. *Arthritis Res Ther.* 2008; 10:17.
- Hellgren L. Association between rheumatoid arthritis and psoriasis in total populations. *Acta Rheum Scand.* 1969; 15(4):650–7.
- Van Romunde LK, Valkenburg HA, Swart-Bruinsma W, Cats A, Hermans J. Psoriasis and arthritis. I. A population study. *Rheumatol Int.* 1984; 4(2):55–60.
- Pedersen OB, Svendsen AJ, Ejstrup L, Skytthe A, Junker P. The occurrence of psoriatic arthritis in Denmark. *Ann Rheum Dis.* 2008; 67(10):1422–6.

48. Madland TM, Apalset EM, Johannessen AE, Rossebo B, Brun JG. Prevalence, disease manifestations, and treatment of psoriatic arthritis in Western Norway. *J Rheumatol*. 2005; 32(10):1918–22.
49. Alamanos Y, Papadopoulos NG, Voulgari PV, Siozos C, Psychos DN, Tympanidou M, et al. Epidemiology of psoriatic arthritis in northwest Greece, 1982–2001. *J Rheumatol*. 2003; 30(12):2641–4.
50. Salaffi F, De Angelis R, Grassi W. Prevalence of musculoskeletal conditions in an Italian population sample: results of a regional community-based study. I. The MAPPING study. *Clin Exp Rheumatol*. 2005; 23(6):819–28.
51. Gelfand JM, Gladman DD, Mease PJ, Smith N, Margolis DJ, Nijsten T, et al. Epidemiology of psoriatic arthritis in the population of the United States. *J Am Acad Dermatol*. 2005; 53(4):573–7.
52. Shbeeb M, Uramoto KM, Gibson LE, O'Fallon WM, Gabriel SE. The epidemiology of psoriatic arthritis in Olmsted County, Minnesota, USA, 1982–1991. *J Rheumatol*. 2000; 27(5):1247–50.
53. Soriano ER, Rosa J, Vellozo E, Schpilberg M, Imamura PM, Diaz J, et al. Incidence and prevalence of psoriatic arthritis in Buenos Aires, Argentina: a 6-year health management organization-based study. *Rheumatology (Oxford)*. 2011; 50(4):729–34.
54. Karreman MC, Luime JJ, Hazes JM, Weel AEAM. The Prevalence and Incidence of Axial and Peripheral Spondyloarthritis in Inflammatory Bowel Disease: A Systematic Review and Meta-analysis. *J Crohns Colitis*. 2017; 11(5):631–42.
55. Baddoura R, Awada H, Okais J, Habis T, Attoui S, Abisaab M. Validation of the European Spondylarthropathy Study Group and Amor criteria for spondyloarthropathies in Lebanon. *Rev Rheum Engl Ed*. 1997; 64(7-9):459–64.
56. Wu ZB, Zhu P, Wang HK, Zheng ZH, Jia Y, Ding J, et al. Prevalence of seronegative spondyloarthritis in the army force of China. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2004; 25(9):753–5.
57. Boyer GS, Lanier AP, Templin DW. Prevalence rates of spondyloarthropathies, rheumatoid arthritis, and other rheumatic disorders in an Alaskan Inupiat Eskimo population. *J Rheumatol*. 1988; 15(4):678–83.
58. Fender P, Paita M, Ganay D, Benech JM. Prevalence of thirty long term disorders for French health insurance members in 1994. *Rev Epidemiol Sante Publique*. 1997; 45(6):454–64. French.
59. Zlatkovic-Svenda M, Rouse M, Radak-Perovic M, Stojanovic R, Vujsinovic-Stupar N, Lazovic-Popovic B, et al. Adaptation and validation of the Rheumatoid Arthritis Quality of Life (RAQoL) questionnaire for use in Serbia. *Rheumatol Int*. 2017; 37(4):641–6.
60. Dougados M, van der Linden S, Juhlin R, Huitfeldt B, Amor B, Calin A, et al. The European Spondylarthropathy Study Group preliminary criteria for the classification of spondylarthropathy. *Arthritis Rheum*. 1991; 34(10):1218–27.
61. Rudwaleit M, van der Heijde D, Landewé R, Listing J, Akkoc N, Brandt J, et al. The development of Assessment of SpondyloArthritis international Society classification criteria for axial spondyloarthritis (part II): validation and final selection. *Ann Rheum Dis*. 2009; 68(6):777–83.
62. Stolwijk C, van Onna M, Boonen A, van Tubergen A. Global Prevalence of Spondyloarthritis: A Systematic Review and Meta-Regression Analysis. *Arthritis Care Res (Hoboken)*. 2016; 68(9):1320–31.
63. Chung HY, Machado P, van der Heijde D, D'Agostino MA, Dougados M. HLA-B27 positive patients differ from HLA-B27 negative patients in clinical presentation and imaging: results from the DESIR cohort of patients with recent onset axial spondyloarthritis. *Ann Rheum Dis*. 2011; 70(11):1930–6.
64. Khan MA. Epidemiology of HLA-B27 and Arthritis. *Clin Rheumatol*. 1996; 15 Suppl 1:10–2.
65. Adebajo A, Boehncke W, Gladman D, Mease P. Psoriatic arthritis and psoriasis: pathology and clinical aspects. Classification criteria for psoriasis. Switzerland: Springer International Publishing; 2016 (XIV):35.
66. Stojanovic R, Vlajinac H, Palic-Obradovic D, Janosevic S, Adanja B. Prevalence of rheumatoid arthritis in Belgrade, Yugoslavia. *Br J Rheumatol*. 1998; 37(7):729–32.

Преваленција спондилоартритиса и његових подтипова – да ли је заиста упоредива?

Мирјана И. Златковић-Швенда^{1,2}, Роксанда М. Стојановић¹, Сандра Б. Шипетић-Грујичић³, Марија М. Радак-Перовић¹, Франсис Гилман⁴

¹Универзитет у Београду, Медицински факултет, Институт за реуматологију, Београд, Србија;

²Универзитет у Источном Сарајеву, Медицински факултет, Република Српска, Босна и Херцеговина;

³Универзитет у Београду, Медицински факултет, Институт за епидемиологију, Београд, Србија;

⁴Универзитет у Лорени, Универзитетска болница, *INSERM-CIC-EC* и *EA 4360 APEMAC*, Нанси, Француска

САЖЕТАК

Увод/Циљ Пораст преваленције спондилоартритиса у последњих неколико деценија не може се приписати само манифестацијама болести.

Циљ овог рада је преглед преваленције спондилоартритиса (SpA) и његових подтипова: анкилозирајућег спондилитиса (АС), псоријазног артритиса (ПсА), реактивног артритиса (РеА), SpA повезаног са инфламаторним болестима црева и недиферентованог SpA (HeSpA), са посебним освртом на методолошке разлике и географске варијације.

Метод Претражена је литература са базама података *PubMed*, *Google Scholar* и *Embase* уз додатно ручно претраживање у односу на термин спондилоартритис и његове подтипове у комбинацији са преваленцијом.

Резултати Земље северне Европе (Шкотска, Шведска, Француска) имају ниже стопе преваленције SpA (0,21–0,45%) у поређењу са јужним (Италија – 1,06% и Турска – 1,35%). Најнижа преваленција SpA у свету забележена је у афричким

и југоисточним азијским земљама (0–0,19%), а највиша на Аљасци (2,5%). Највећа варијабилност у преваленцији ПсА запажена је у Европи (северни део 0,02–0,19%, јужни 0,42%). Најнижа преваленција ПсА у свету забележена је у Јапану – 0,001%, а затим у Кини 0,01–0,1%. У Европи се преваленција РеА креће од 0,04% у Грчкој до 0,1% у Србији и Немачкој, док је преваленција УнSpA од 0,02% у Србији до 0,67% у Немачкој. Највиша преваленција УнSpA у свету нађена је у Либији – 3,4%. Студије које одређују преваленцију SpA разликују се у бројним карактеристикама: различит метод формирања узорака и употребљених критеријума, укључивање различитих старосних група, различит приказ резултата итд.

Закључак Варијације у преваленцији SpA не могу се приписати само генетским или географским разликама. Различита преваленција делимично је условљена и неуједначеним методама истраживања, детаљније описаним у овом чланку.

Кључне речи: епидемиологија; преваленција; спондилоартритис; анкилозирајући спондилитис; псоријазни артритис