



**СРПСКИ АРХИВ**  
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО  
**SERBIAN ARCHIVES**  
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: [office@srpskiarhiv.rs](mailto:office@srpskiarhiv.rs), Web address: [www.srpskiarhiv.rs](http://www.srpskiarhiv.rs)

**Paper Accepted\***

**ISSN Online 2406-0895**

**Case report / Приказ болесника**

Anita Krsman<sup>1,2,\*</sup>, Branislava Baturan<sup>1,2</sup>, Dmtar Vlahović<sup>3</sup>, Zorica Grujić<sup>1,2</sup>,  
Đorđe Petrović<sup>1,2</sup>, Biljana Radovanović<sup>3</sup>

**Mature ovarian teratoma-associated encephalitis**

Зрели тератом јајника повезан са енцефалитисом

<sup>1</sup>Clinical Center of Vojvodina, Department of Obstetrics and Gynecology, Novi Sad, Serbia;

<sup>2</sup>University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia;

<sup>3</sup>Clinical Center of Vojvodina, Department of Neurology, Novi Sad, Serbia

**Received: December 3, 2020**

**Revised: December 21, 2021**

**Accepted: December 22, 2021**

**Online First: January 13, 2022**

**DOI: <https://doi.org/10.2298/SARH201203004K>**

\***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal.

The date the article was made available online first will be carried over.

**\*Correspondence to:**

Anita KRSMAN

Department of Obstetrics and Gynecology, Clinical Centre of Vojvodina

Branimira Ćosića 37, Novi Sad, Serbia

E mail: [anitaanaaleksic@gmail.com](mailto:anitaanaaleksic@gmail.com)

## Mature ovarian teratoma-associated encephalitis

### Зрели тератом јајника повезан са енцефалитисом

#### SUMMARY

**Introduction** Autoimmune encephalitis associated with ovarian teratoma is a serious and potentially fatal pathology. While this clinical entity is known to neurologists, the available literature rarely mentions the role of a gynecologist in diagnostic imaging and treatment. Although several months have passed from the appearance of the symptoms to surgical treatment, this case shows that even then a complete recovery is possible.

**Case presentation** The patient was a 28-year-old female, brought to the hospital because a sudden onset of unusual behavior - an acute psychosis with suicidal thoughts and auditory hallucinations. Soon after the admission she became delirious, uncooperative and agitated. Blood check, neurological assessment and cranial computed tomography yielded normal results. Therefore, a psychiatric disorder was suspected. Electroencephalogram revealed a diffuse encephalitic insufficiency. As cerebrospinal fluid was negative for infections, the autoimmune etiology of the disease was suspected. Abdominal computer tomography showed a complex right ovarian mass measuring 50 x 40 x 30 mm, confirmed by vaginal ultrasound. Laparoscopy with right adnexectomy was performed. The pathohistological finding showed a mature teratoma. In the meantime, the result of the cerebrospinal fluid test came positive for anti NMDAR antibodies. Six months after surgery, the patient was in a good mental and neurological status without symptoms.

**Conclusion** Gynecologists should be aware of the presence of ovarian tumors in encephalitis cases. A timely diagnosis of the underlying gynecological cause of a neurological condition, allows for prompt treatment and can remarkably improve clinical conditions and, thus, be lifesaving.

**Keywords:** autoimmune encephalitis; teratoma; tumor; young women

#### САЖЕТАК

**Увод** Аутоимуни енцефалитис повезан са тератомом јајника је озбиљна и потенцијално фатална болест. Иако је ово познат клинички ентитет неуролозима, желели бисмо да скренемо пажњу гинеколозима, јер имају посебну улогу у дијагностици и оперативном лечењу пацијенткиња оболелих од аутоимуног енцефалитиса. Приказан је случај пацијенткиње чији је исход лечења успешан, иако је од појаве првих симптома па до постављања дијагнозе и операције прошло више месеци.

**Приказ случаја** Пацијенткиња стара 28 година, хоспитализована је због изненадог развоја психозе са звучним халуцинацијама и суицидалним мислима. Убрзо након пријема пацијенткиња постаје изразито узнемирена и агресивна. Лабораторијске анализе, неуролошки статус и компјутеризована томографија ендокранијума били су уредни. Електроенцефалограм је показао дифузни енцефалитис. Прегледом ликвора искључен је инфективни узрочник. Постављена је сумња на аутоимуно етиологију обољења. Компјутеризованом томографијом абдомена и вагиналним ултразвучним прегледом уочена је комплексна туморска промена десног јајника димензија 50 x 40 x 30 мм. Урађена је десна аднексектомија лапароскопски. Патохистолошка анализа је показала да се ради о зром тератому. Накнадно је пристигао позитиван узорак ликвора на анти НМДАР антитета. Шест месеци након операције пацијенткиња је уредног психичког и неуролошког статуса без тегаба.

**Закључак** Гинеколози треба да буду свесни постојања везе између енцефалитиса и тумора јајника младих жена. Брзо постављање дијагнозе, утврђивање гинеколошког узрока неуролошког обољења и адекватан терапијски поступак воде излечењу овог озбиљног, а понекад и смртоносног обољења.

**Кључне речи:** аутоимуни енцефалитис, тератом, тумор, младе жене

## INTRODUCTION

Anti-N-Methyl-D-Aspartate (NMDA) receptor encephalitis has been recognized as the second most common cause of autoimmune encephalitis, after acute demyelinating encephalomyelitis [1]. Anti-NMDAR encephalitis in young women is usually associated

with the presence of ovarian masses. [1, 2]. Due to the prevalence of psychiatric symptoms, 77% of patients are initially hospitalized by a psychiatrist [3, 4, 5]. The above symptoms are followed by a memory deficit, autonomic instability, seizures, muscle rigidity, movement disorder and hypoventilation [6]. It is not uncommon for such patients to require ventilation support and intensive care to stay alive [7]. In this patient, withdrawal symptoms and recovery occurred immediately after surgical treatment, although literature data show that recovery can take up to 24 months [1, 2, 7]. However, some patients still have problems with cognition (understanding, memory, attention and expression) and motor function (walking, taking care of yourself, swallowing) [1]. Anti-NMDAR encephalitis is well known to neurologist. Still, this is less well known to gynecologists, who may have a decisive role in etiological management. Raising awareness about this disease among gynecologists can be of great importance for earlier diagnosis and thus faster healing [8].

We present a rare case of Anti-NMDAR encephalitis induced by ovarian teratoma. Although five months passed from the appearance of the first symptoms to removal of the tumor, our patient completely recovered without consequences. The purpose of the study was to present our first experience, as gynecologists, with such a rare pathology and to point out the challenge in diagnosis and treatment patient with NMDAR encephalitis.

## **CASE REPORT**

A 28-year-old woman was hospitalized at the Clinic for Psychiatry, Clinical Center of Vojvodina, complaining of symptoms of an acute episode of psychotic behavior. According to heteroanamnesic data, the patient had a sudden onset of unusual behavior, an acute psychosis with suicidal thoughts, auditory hallucinations and decreased food intake. Soon after admission, she became delirious, uncooperative and agitated. She had no past medical

history and there was no history of substance abuse. Fear, loss of appetite, decreased oral intake and insomnia were the dominant symptoms. The neurological assessment and basic blood evaluation yielded negative results, cranial computed tomography was unremarkable. The patient exhibited severe psychomotor agitation which led to self-inflicted harm. In the further course of hospitalization, she destroyed her lower lip and caused a fracture of the alveolar extension of the lower jaw. An extensive array of microbiological and serological tests were done and all were negative. The autoimmune antibodies and tumor marker examination results were also negative. The brain magnetic resonance imaging (MRI) and angiography (MRA) were unremarkable. Electroencephalogram (EEG) revealed a diffuse encephalitic insufficiency. The sample of cerebrospinal fluid was negative for infections and was placed for further testing of anti-NMDAR antibodies. The patient was started on intravenous (IV) steroids, intravenous immunoglobulin (IVIG) and plasma exchange. There was a slight improvement after the applied therapy.

Finally, since autoimmune encephalitis (anti-NMDAR) is usually caused by the presence of a dermoid cyst in young women [1], a gynecologist was consulted. The computer tomography scan (CT) showed the presence of a complex right ovarian mass measuring 50x40x20 mm, confirmed by vaginal ultrasound (Figures 1 and 2). Laparoscopy with right adnexectomy was performed. The ovarian mass contained cystic components as well as skin, hair, brain tissue and cartilage. The final pathology result was mature teratoma. In the meantime, the result of the cerebrospinal fluid test was positive for anti-NMDAR antibodies. After four months, the patient had problems with short-term memory. Six months after the surgery, the patient was in a good mental and neurological status, without any symptoms.

Written consent to write and publish this case report was obtained from the patient. The review was approved by the Ethics Committee of the Clinical Center of Vojvodina.

## DISCUSSION.

Teratoma-associated encephalitis (Anti-NMDAR encephalitis) was first noted in 1997 [1, 2, 9, 10]. Dalmau et al. [11] were the first to describe the progressive and potentially fatal course of this phenomenon. NMDA receptors are involved in cognitive processes: behavior, memory and learning [3]. Delay in diagnosis adversely affects the outcome which ranges from complete recovery, recovery followed by psycho-neurological sequelae, to death. An accurate and rapid diagnosis, followed by an early removal of the tumor in combination with immunotherapy, leads to a favorable outcome. This disease affects mostly young women and is associated with ovarian teratomas, which can contain nervous tissue expressing NMDA receptors [1, 2, 12]. However, cases have been reported in patients aged from 8 months to 85 years [1]. Although there is a possibility of extra-ovarian teratomas (2%) and other tumors (4%, for instance lung, breast, thymus), ovarian teratomas are most often associated with this type of encephalitis [13]. Anti-NMDA receptor encephalitis often (70%) begins with prodromal flu-like symptoms (fever, headache and fatigue) [2]. After this uncharacteristic clinical picture, psychological symptoms develop with a predominance of the following: agitation, hallucinations, changes in behavior and anxiety [14]. There is a significant overlap of neurological and psychiatric symptoms associated with autoimmune encephalitis. Therefore, cases like this can manifest as a whole spectrum of neurological or psychiatric diseases (schizophrenia, paranoia, hallucinations, agitation, depression, anxiety or substance abuse) [15]. The diagnosis of autoimmune encephalitis is challenging because the differential diagnosis involves the exclusion of the entire spectrum of the diseases (HSV encephalitis, CMV encephalitis, Hashimoto's encephalopathy, systemic lupus erythematosus encephalopathy, antiphospholipid antibody syndrome, Sjögren's syndrome, and primary central nervous angiitis). However, inadequate response to psychiatric therapy and neurological status should lead clinicians to consider encephalitis. MRI has been

reported to be negative in up to 50-70% of cases, and in our case as well [15]. EEG abnormalities are present in most cases (90%) [1]. The final diagnosis is based on the presence of anti-NMDAR autoantibodies in cerebrospinal fluid [2]. After anti-NMDAR encephalitis has been confirmed, imaging studies are performed, such as vaginal ultrasound examination, MRI, CT and positron emission tomography (PET CT) [16, 17]. It is important to note that if autoimmune etiology is suspected it is necessary to intensively search for tumor changes in the ovaries [18]. Transvaginal ultrasound should be the first step in the diagnosis of ovarian masses [2]. Therefore, gynecologists must be aware of the connection between ovarian tumors and autoimmune encephalitis and look in detail for even minimal ovarian changes. Sometimes this syndrome has been associated to patients with no detectable underlying neoplasms. It is theorized that the syndrome may be triggered by microscopic germ cell tumors undetectable by imaging diagnostic methods [4]. Moreover, it has been reported that ovarian teratomas were discovered years after the onset of symptoms of anti-NMDAR encephalitis [4]. Surgical intervention is imperative, since a study by Titulaer MJ et al. showed that five out of six patients with a confirmed NMDAR encephalitis and an ovarian teratoma who did not undergo surgery died [1]. The removal of the tumor will lead to an improvement in the patient's neurological performance within a few days or weeks [15]. The data from previous research indicate that a surgical procedure with immunotherapy have the best results in the treatment of this potentially fatal condition [2, 11]. Any delay in treatment can result in worsening of the patient's condition and even death [1]. It has been found that the outcome is more favorable if the operation is conducted after less than 4 months from the onset of symptoms [11]. In the case of this patient, the surgical treatment was performed 5.5 months after the appearance of the first symptoms. Women in the reproductive period are most often affected with this disease so the biggest challenge for gynecologists is how to completely remove the tumor while preserving fertility

in the patient. Because a laparoscopic examination for determining ovarian teratoma is less-invasive than laparotomy, trial laparoscopy is acceptable if an ovarian tumor cannot be detected by various imaging tests [8]. Additionally, there have been cases where, despite the fact that the imaging procedures did not determine the existence of ovarian tumors, an ovariectomy was performed in order to stop the production of antibodies [3, 19, 20]. Postoperative histological examination in these cases confirmed the existence of occult teratoma consisting of skin, hair, cartilage and nervous tissue. Types of surgical treatment range from selective removal of lesions by ultrasound-guided laparoscopy [21], cystectomy, uni/bilateral ovariectomy and uni/bilateral adnexectomy. Although some studies emphasize that cystectomy and adnexectomy have similar outcomes [22, 23], we decided to perform a unilateral adnexectomy. As a possible limitation of the type of treatment in this case, we can mention the extent of the surgical procedure, which is not in accordance with the previous statements about preserving the fertility of young women. Pathohistological examination confirmed the existence of a mature teratoma with components of nervous tissue. This is consistent with literature reports showing that dermoid cysts, containing nervous tissue, are the most common pathohistological finding in patients with anti-NMDAR encephalitis [12].

As a limitation of the study, we state the period of postoperative monitoring. We followed the patients for 6 months postoperatively and it is not known whether the patient will develop new symptoms in the subsequent period. However, this case report may help the identification of this rare disease and therefore is an important educational article.

In conclusion, it should be highlighted that it is extremely important to discover the cause of the disease as soon as possible, in order to give the patient a greater chance of complete recovery. Since gynecologists play an important role in the treatment of these patients, they should be aware of the presence of tumors in patients with encephalitis. Determining the gynecological etiology of this condition, with the earliest possible surgical

treatment, can lead to a favorable outcome and thus save the lives of young women.

#### **ACKNOWLEDGMENT**

The abstract of this paper was awarded as the best case report presented at 9th Symposium of Pediatric and Adolescent Gynecology, Belgrade (Serbia), June 26, 2020.

**Conflict of interest:** None declared.

Paper accepted




## REFERENCES

1. Titulaer MJ, McCracken L, Gabilondo I, Armangue T, Glaser C, Iizuka T et al. Treatment and prognostic factors for long-term outcome in patients with anti-NMDA receptor encephalitis: an observational cohort study. *Lancet Neurol* 2013; 12:157-65. DOI: 10.1016/S1474 4422(12)70310-1. PMID: 23290630.
2. Dalmau J, Armangué T, Planagumà J, Radosevic M, Mannara F, Leypoldt F, Geis C, Lancaster E, Titulaer MJ, Rosenfeld MR, Graus F. An update on anti-NMDA receptor encephalitis for neurologists and psychiatrists: mechanisms and models. *Lancet Neurol* 2019; 18: 1045-1057. DOI: 10.1016/S1474-4422(19)30244-3. PMID: 31326280.
3. Abdul-Rahman et al. Anti-N-methyl-D-aspartate receptor encephalitis with an imaging-invisible ovarian teratoma: a case report. *J Med Case Reports* 2016; 10:296. DOI 10.1186/s13256-016-1067-4.
4. Mann AP, Grebenciucova E, Lukas RV. Anti-N-methyl-d-aspartate-receptor encephalitis: diagnosis, optimal management and challenges. *Ther Clin Risk Manag* 2014; 10:517-25. DOI 10.2147/TCRM.S61967. PMID: 25061311.
5. Wang HY, Li T, Li XL, Zhang XX, Yan ZR, Xu Y. Anti-N-methyl-D-aspartate receptor encephalitis mimics neuroleptic malignant syndrome: case report and literature review. *Neuropsychiatr Dis Treat* 2019; 15:773-778. DOI:10.2147/NDT.S195706. PMID: 31040676.
6. Li W, Jia D, Tong L, Lun Z, Li H. Anti-N-methyl-daspartate receptor encephalitis induced by bilateral ovarian teratomas with distinct histopathologic types: a case report and brief literature review. *Medicine* 2019; 98:48. PMID: 31770255. DOI: 10.1097/MD.000000000018148.
7. Schubert J, Bramer D, Huttner HB et al. Management and prognostic markers in patients with autoimmune encephalitis requiring ICU treatment. *Neurol Neuroimmunol Neuroinflamm* 2019; 6:514. DOI: 10.1212/NXI.0000000000000514. PMID: 30568992.
8. Yan B, Wang Y, Zhang Y, Lou W. Teratoma-associated anti-N-methyl-D-aspartate receptor encephalitis: a case report and literature review. *Medicine (Baltimore)* 2019; 98(21):e15765. DOI: 10.1097/MD.000000000015765. PMID: 31124965.
9. Dai Y, Zhang J, Ren H, Zhou X, Chen J, Cui L, Lang J, et al. Surgical outcomes in patients with anti-N-methyl D-aspartate receptor encephalitis with ovarian teratoma. *Am J Obstet Gynecol* 2019; 221(5): 481-485. DOI: 10.1016/j.ajog.2019.05.026. PMID: 31128109.
10. Azizyan A, Albrektson JR, Maya MM, Pressman BD, Moser F. Anti NMDA encephalitis: an uncommon autoimmune mediated form of encephalitis. *J Radiol Case Rep* 2014; 8 (8):1-6. DOI: 10.3941/jrcr.v8i8.1566. PMID: 25426239.
11. Dalmau J, Tuzun E, Wu HY et al. Paraneoplastic anti-N-methyl-d-aspartate receptor encephalitis associated with ovarian teratoma. *Ann Neurol* 2007; 61 (1):25-36. DOI: 10.1002/ana.21050. PMID: 17262855.
12. Chefdeville A, Treilleux I, Mayeur ME, Couillaud C, Picard G, Bost C, et al. Immunopathological characterization of ovarian teratomas associated with Anti-N-Methyl-D aspartate receptor encephalitis. *Acta Neuropathol Commun* 2019; 38. DOI: 10.1186/s40478- 019-0693-7. PMID: 30857565.
13. Liu CY, Zhu J, Zheng XY, Ma C, Wang X. Anti-N-Methyl-D-aspartate receptor encephalitis: a severe, potentially reversible autoimmune encephalitis. *Mediators Inflamm* 2017. DOI: 10.1155/2017/6361479. PMID: 28698711.
14. Subeh G K, Lajber M, Patel T et al. Anti-N-Methyl-D-Aspartate receptor encephalitis: A detailed review of the different psychiatric presentations and red flags to look for in suspected cases. *Cureus* 2021; 13(5): 15188. DOI:10.7759/cureus.15188. PMID: 34178509.
15. Lwin S, San Yi M, Mardiana K, Woon SY, Nwe TM. Ovarian teratoma-associated anti-NMDAR encephalitis in a 12-year-old girl. *Med J Malaysia* 2020; 75(6):731-733. PMID: 33219185.
16. Mitra AD, Afify A. Ovarian teratoma associated Anti-N-methyl-D-aspartate receptor encephalitis: a difficult diagnosis with a favorable prognosis. *Autops Case Rep* 2018; 18:8(2):e2018019. DOI:10.4322/acr.2018.019. PMID: 29780755.
17. Habib E, Patel J, Kerndt C, Summers D, Bragg T. A rare case of anti-NMDA receptor encephalitis associated with an ovarian teratoma. *American Journal of Medical Case Reports* 2021; 9(7):354-357. DOI: 10.12691/ajmcr-9-7-3.
18. Thiagarajan M, Sebastian A, Thomas DS, Thomas A, Peedicayil A, Mathew V. Ovarian teratoma and N-Methyl-D-Aspartate Receptor autoimmune encephalitis: Insights into imaging diagnosis of teratoma and timing of surgery. *J Clin Gynecol Obstet* 2021; 10(1):22-27. DOI: <https://doi.org/10.14740/jcgo715>.
19. Mahendra INB, Saspriyana KY, Ekawati NP. Mature ovarian teratoma associated with anti-N-Methyl-D-aspartate receptor encephalitis: A case report. *The Medical Journal of Malaysia* 2021; 76(1):110-113. PMID: 33510121.
20. Boeck A, Logemann F, Krauss T, Hussein K, Bültmann E, Trebst C, Stangel M. Ovariectomy despite negative imaging in anti-NMDA receptor encephalitis: effective even late. *Case Rep Neurol Med* 2013. DOI: 10.1155/2013/843192. PMID: 23533859.

21. Jones BP, Rees R, Saso S, Stalder C, Smith JR et al. Ultrasound-guided laparoscopic ovarian preserving surgery to treat anti-NMDA receptor encephalitis. *BJOG* 2017; 124: 337-41. DOI: 10.1111/1471-0528.14214. PMID: 27425649.
22. Asai S, Ishimoto H, Yabuno A, Asada H, Seki M et al. Laparoscopic cystectomy of ovarian teratoma in anti-NMDAR encephalitis: 2 case reports. *J Minim Invasive Gynecol* 2011; 18:135-37. DOI: 10.1016/j.jmig.2010.10.002. PMID: 21195970.
23. Schiavi, Michele Carlo, et al. Mature ovarian teratoma: neurological implications in a young woman. *Case Reports in Obstetrics and Gynecology* 2021; 2021. <https://doi.org/10.1155/2021/3085559>

Paper accepted



**Figure 1.** Computed tomography scan of the patient's pelvis

Paper accepted



**Figure 2.** Transvaginal ultrasound of the ovarian teratoma in our patient

Paper accepted