



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Violeta Knežević^{1,2,*}, Tijana Azaševac^{1,2}, Dragana Milijašević^{1,3}, Uroš Milošević^{1,4},
Lada Petrović^{1,2}

**Predictors of renal function non-recovery in critically ill patients with acute
kidney injury treated with continuous renal replacement therapy**

Предиктори неопоравка функције бубрега код критично оболелих пацијената са
акутним оштећењем бубрега лечених континуираном дијализом

¹University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia;

²University Clinical Center of Vojvodina, Clinic for Nephrology and Clinical Immunology, Novi Sad, Serbia;

³Institute of Public Health of Vojvodina, Novi Sad, Serbia;

⁴University Clinical Center of Vojvodina, Clinic for Abdominal and Endocrine Surgery, Novi Sad, Serbia

Received: September 9, 2022

Revised: February 15, 2024

Accepted: February 29, 2024

Online First: March 6, 2024

DOI: <https://doi.org/10.2298/SARH220909020K>

***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. Srp Arh Celok Lek. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

***Correspondence to:**

Violeta KNEŽEVIĆ

University of Novi Sad, University Clinical center of Vojvodina, Clinic for Nephrology and Clinical Immunology, Hajduk Veljkova 1–7, 21000 Novi Sad, Serbia

E-mail: vknezevic021@gmail.com

Predictors of renal function non-recovery in critically ill patients with acute kidney injury treated with continuous renal replacement therapy

Предиктори неопоравка функције бубрега код критично оболелих пацијената са акутним оштећењем бубрега лечених континуираном дијализом

SUMMARY

Introduction/Objective Acute kidney injury (AKI) is a highly prevalent complication among the critically ill individuals who are admitted to the intensive care unit (ICU). This study deals with identifying the frequency and predictors of the lack of renal function recovery in non-renal functions among critically ill patients requiring dialysis for AKI (AKI-D).

Methods The study included 440 ICU patients from the University Clinical Center of Vojvodina in the period from 2014 to 2018. The patients required Continuous Renal Replacement Therapy (CRRT). In this study, we analyzed various factors including demographic features, clinical characteristics, laboratory parameters, co-morbidities, as well as the need for vasopressor therapy and mechanical ventilation on the day when AKI was confirmed. Additionally, we examined the different modalities of CRRT, which were used.

Results A retrospective analysis of the results included discovered that out of 440 patients with AKI-D, 242 (55%), average age 63.14, did not recover renal function. Significant predictors of renal function non-recovery in critically ill patients with AKI-D were: the patients age over 65 ($p = 0.044$), starting time of CRRT ($p = 0.043$), mechanical ventilation ($p = 0.044$) and previous kidney disease ($p = 0.005$). Significant predictors of renal function non-recovery in critically ill septic patients with AKI-D were: the patients age over 65 ($p = 0.002$), diabetes mellitus ($p = 0.023$), previous kidney disease ($p = 0.045$), CRP values $< 100 \text{ mg/l}$ ($p = 0.033$) and procalcitonin ($p = 0.010$), while in non-septic patients, the significant predictors of renal function non-recovery includes previous kidney disease ($p = 0.035$).

Conclusion Out of all examined predictors, both in septic and non-septic patients, previous kidney damage presents the greatest risk for renal function non-recovery in critically ill patients with AKI-D.

Keywords: acute kidney injury; recovery of renal function; critically ill; sepsis

САЖЕТАК

Увод/Циљ Акутно оштећење бубрега (АОБ) је честа компликација код критично оболелих пацијената хоспитализованих у одељењима интензивног лечења (ОИЛ). Циљ ове студије је утврђивање преваленције и предиктора неопоравка функције бубрега код критично оболелих пацијената са АОб-зависним од дијализе (АОб-Д).

Методе Студијом је било обухваћено 440 пацијената лечених на ОИЛ, Ургентног центра, Универзитетског Клиничког Центра Војводине у периоду 2014. до 2018. Године који су захтевали континуирану замену функције бубрега (ЗФБ). Анализирани су демографски, клинички и лабораторијски параметри, коморбидитети, потреба за вазопресорном терапијом и механичком вентилацијом у дану потврђеног АОб и модалитети ЗФБ.

Резултати Ретроспективном анализом резултата установљено је да од укупно 440 пацијената са АОб-Д, 242 (55%), просечне старости 63.14 година, није опоравило функцију бубрега. Значајни предиктори неопоравка функције бубрега код критично оболелих са АОб-Д били су: старост изнад 65 година ($p = 0.044$), време почетка ЗФБ ($p = 0.043$), механичка вентилација ($p = 0.044$) и претходне болести бубрега ($p = 0.005$). Значајни предиктори неопоравка функције бубрега код септичних пацијената са АОб-Д били су: старост изнад 65 година ($p = 0.002$), *diabetes mellitus* ($p = 0.023$), претходне болести бубрега ($p = 0.045$), *CRP* вредности $< 100 \text{ mg/l}$ ($p = 0.033$) и прокалцитонин ($p = 0.010$), док је код критично оболелих пацијената без сепсе значајан предиктор неопоравка функције бубрега претходне болести бубрега ($p = 0.035$).

Закључак Од свих испитиваних предиктора на укупном узорку, код септичних и несептичних пацијената, претходне бубрежне болести представљају највећи ризик за неопоравак функције бубрега код критично оболелих пацијената са АОб-Д.

Кључне речи: акутно оштећење бубрега; опоравак бубрежне функције; критично оболели; сепса

INTRODUCTION

The development of acute kidney injury (AKI) is associated with various adverse outcomes such as prolonged intensive care unit (ICU) stay, development of chronic kidney disease (CKD), increased the mortality rate along with the increased treatment costs [1, 2].

The reported prevalence of AKI in ICUs varies from study to study and runs the gamut from 5% to 67% [3]. According to Opgenorth et al about 5–10% of these patients require renal replacement therapy (RRT) [4].

In patients with AKI requiring RRT, the recovery of renal function occurs in only 20–60% of patients and patients survival without the need for RRT is considered successfully [4, 5]. Only 1–6% recovery of renal function occurs after 90 days of starting RRT [6]. AKI stage and time required for renal function recovery significantly influence morbidity/mortality rates after an acute episode of patients requiring dialysis for AKI (AKI-D) [7].

The results so far have indicated numerous predictors of renal function recovery and include those related to patients (age, CKD, proteinuria, co-morbidity, etiology, stage and duration of AKI, exposure to nephro-toxic drugs and/or diagnostic contrast agents, multi organ dysfunction), as well as potential predictors related to dialysis procedures itself (modalities, membrane, dose intensity, lasting and frequency of procedures) [5, 7, 8]. The aim of this study is to determine the prevalence and predictors of renal function non-recovery in critically ill patients with AKI-D.

METHODS

This retrospective study was conducted between 2014 and 2018 at the University Clinical Center of Vojvodina. The study included ICU and ER 440 adult surgical and non-surgical patients suffering from AKI and AKI on CKD who required continuous renal replacement therapy (CRRT).

The study analyzed various parameters, including demographic features, co-morbidities, laboratory and clinical analyses such as urea, creatinine, C-reactive protein (CRP), procalcitonin (PCT), quick sequential organ failure assessment score (qSOFA) score, the need for vasopressor therapy (VT), and mechanical ventilation (MV) on the day when AKI was confirmed.

Different modalities of continuous CRRT were utilized, including continuous veno-venous hemodiafiltration (CVVHDF), continuous veno-venous hemofiltration (CVVHF), continuous veno-venous hemodialysis (CVVHD), and CVVHD combined with CVVHDF. Discontinuation of CRRT was guided by the establishment of diuresis of 1000-1500ml/day without diuretics. The recovery of renal function after AKI-D was defined independently from CRRT within 90 days of starting RRT. The criteria for early start of CRRT were stages 2 or 3 AKI according to KDIGO guidelines and/or hypervolemia with poor response to conservative treatment within 24h of AKI diagnosis, while manifested clinical complications related to AKI were criteria for i.e. late start of CRRT after 24 hours.

The choice of CRRT modalities was determined by the clinicians, guided by international guidelines. The CRRT prescription included specific details such as treatment modality, circulation, dilution mode, replacement and dialysis fluid flow, patient weight, and heparin anticoagulation.

The study also highlights the reasons for delayed initiation of RRT, such as organizational issues, unavailability of equipment, difficulty in catheter placement, or the need for surgical interventions or radiological tests before starting RRT. Some patients received intermittent dialysis initially based on factors like hemodynamic stability or equipment availability.

Certain exclusion criteria were applied to patients with urgent need for RRT, including specific laboratory abnormalities (urea > 50 mmol/l, K > 6.5 mmol/l, pH < 7.15) and acute pulmonary edema, and those treated with conservative therapy.

CRRT was performed on the Multifilter and the Prismaflex; standard high-flux filters and membranes/adsorbers were used in septic patient. The CRRT prescription included: treatment modality, blood flow, dilution mode, replacement and dialysis fluid flow, and the patient's weight and heparin anticoagulation, according to clinical practice guideline [9].

The study was carried out according to the principles of the Helsinki Declaration and it was approved by the local Ethics Committee, decision number 00-116.

Statistical analysis

IBM® SPSS® Statistics software version 20.0 was used for statistical data processing. The obtained data were analyzed using the chi-square test and t test of independent samples (when examining individual underlying possible predictors) and binomial logistic regression analysis (when examining individual continuous potential predictors, as well as when analyzing the association of all possible predictors along with no renal function recovery). The statistical significance of the influence of potential risk factors for the absence of renal function recovery was examined in the course of the study. The Box-Tidwell test was previously applied in order to confirm that the correlation between each continuous predictor and the corresponding logarithm of the odds ratio is linear.

RESULTS

Out of 440 patients with AKI-D, 242 (55%), average age 63.14, did not recover renal function. In patients who did not recover kidney function, earlier kidney diseases ($p = 0.002$) and vasopressor therapy ($p = 0.010$) were significantly more prevalent, and those patients also

had a significantly higher average ultrafiltration (ml) ($p = 0.046$) and a significantly lower percentage of patients started earlier CRRT ≤ 24 hours ($p = 0.050$). There were more male patients in the group who did not recover their renal function in comparison to the patients who did (69.7% vs. 64.5%) (Table 1).

Significant predictors such as age ($p = 0.044$), CRRT start time ($p = 0.043$), lung MV ($p = 0.044$) and previous kidney disease ($p = 0.005$) were singled out in the entire sample of critically ill patients. CRRT start within 24 hours of AKI diagnosis was associated with a 1.353 times higher risk (95% CI 1.009–1.814) for non-recovery of kidney function, age was associated with a 1.437 times higher risk (95% CI 1.010–2.043), mechanical ventilation was associated with a 1.63 -fold higher risk (95% CI 1.013–2.633) and previous renal disease with a 5.49-fold higher risk for renal function non-recovery (95% CI 1.661–18.148) (Table 2).

In the septic patient group, age ($p = 0.002$), CRP ($p = 0.033$), PCT ($p = 0.010$), diabetes mellitus ($p = 0.023$) and previous renal disease ($p = 0.045$) were significant predictors of renal function non-recovery. The patients over 65 years of age were associated with a 1.63-fold higher risk of renal function non-recovery (95% CI 1.063–2.504). The patients over 65 years of age were at three times higher risk (95% CI 1.486–5.760), the ones with CRP higher than 100 mg/l were associated with 2.49 times higher risk (95% CI 1.077–5.748), diabetes mellitus with 2.71 times higher risk (95% CI 1.144–6.414), PCT with 2.84 times higher risk (95% CI 1.279–6.308) and previous kidney disease with 7.08 times higher risk of renal function non-recovery (95% CI 1.046–47.884) (Table 3). In the non-septic group of patients, previous kidney disease was associated with a 6.25 times greater risk of renal function non-recovery ($p = 0.035$, 95% CI 1.14–34.273) (Table 4).

DISCUSSION

Although CRRT has been used for many years in critically ill hemodynamically unstable patients with AKI, the prognosis and recovery of renal function are still uncertain.

A retrospective analysis of the results found that out of a total of 440 patients with AKI, 198 (45%) did not recover their kidney function, which corresponds with previous results [4]. Having analyzed the entire sample, we found that age, the starting time of CRRT, mechanical ventilation and previous kidney disease play a significant predictive role in the outcome of renal function. The subgroup analysis has also shown that these results were consistent after adjustment for multiple variables. Namely, patients over 65 have a 1.63 times greater risk of renal function non-recovery. In a retrospective study by Jiang et al. [10] that included cardiac surgery patients with AKI, multivariate analysis showed that age was an independent prognostic factor for renal function non-recovery during the first week of CRRT treatment. The authors of previous studies investigated the outcomes in patients with AKI treated with CRRT and their results, as well as in our study, indicate that older age has a significant prediction for renal function non-recovery [11]. It is highly likely that the reason for such results is the higher frequency of co-morbidities in the elderly population, but also the structural and functional changes in the kidney that occur with aging [12]. In critically ill patients AKI is manifested by varying degrees of uremia, volemia, acid-base status disorders, physiological and non-renal disorders, and often has a variable course. The decision to start RRT in these patients may depend on numerous factors and it represents a very complex process [13]. In relation to that, the start time of RRT has been difficult to study and has shown considerable variations of clinicians and institutions [14]. Due to its changeable clinical course of AKI, constant monitoring of patients is necessary to ensure that RRT starts timely, i.e. before the appearance of uremic and metabolic complications, but when there are signs of general improvement in the clinical condition and spontaneous recovery of renal function. The results of our study

indicate that "early" initiation of RRT in critically ill patients with AKI significantly affects the rate of renal function non-recovery. Contrary to our results, Lin et al. [15] have concluded that the "early" start of RRT does not significantly affect the recovery of kidney function. The aforementioned results correspond to the meta-analysis published by Ponce et al. [16], where it was found that "early" initiation of RRT increases the chance of recovery of renal function by 30% compared to the "late" one. Our previous study, which was aimed at identifying predictors of kidney function recovery, with the very same criteria used both for "early" and "late" CRRT start time just as in this study, did not find any predictors of recovery of renal function in the "early" group, while in the "late" CRRT group, non-diabetic patients were found to have a 3.5 times higher chance of recovery of function compared to patients with diabetes [17]. Completely different results were published in a research by Castro et al. [18] and indicated that "early" initiation of RRT has no significant effect on the recovery of renal function. However, the studies included in this analysis were highly prone to be biased due to the study design itself, mixed cases, definition of timing of RRT initiation and variation in outcome determination [18]. Mechanical ventilation, while essential in providing respiratory support for critically ill patients, can potentially exacerbate pre-existing lung injury or lead to the development of new lung injury, commonly known as ventilator-associated lung injury (VALI) [19]. According to the available literature, we have observed that the majority of studies investigated the impact of mechanical ventilation on the development of AKI in critically ill patients, but we have not found any research on whether mechanical ventilation has an impact on the recovery of renal function. In our study, we found that MV was associated with a 1.63-fold higher risk for failure to recover renal function. In a randomized clinical study, Yang et al. [20] found that protective mechanical ventilation compared to conventional MV had no significant effect on AKI progression, although AKI progression occurred in both groups of patients. Ostermann et al. [21] performed a murine model of ventilator-associated

lung injury and concluded that a short period of MV with high tidal volumes (17 mL/kg) causes increased lung permeability and liver and kidney inflammation. As pulmonary mechanical ventilation affects both the onset and progression of AKI, it probably has an effect on the recovery of renal function as well, which needs to be determined in future prospective randomized trials.

In the subgroup of septic patients, diabetes mellitus was linked to a 2.71-fold higher risk and previous kidney disease with a 7.08-fold higher risk of renal function non-recovery. Various studies indicate that diabetes mellitus potentially increases morbidity and mortality in patients with AKI [22]. In a multicenter study by Chung et al. [23], diabetes mellitus did not increase the risk of developing AKI, but significantly decreased the probability of renal function recovery. Regarding previous kidney conditions and renal function recovery, our results correlate with most studies, one of which was conducted by Vijayan et al. [24], which included patients with previous kidney diseases were significantly less likely to recover renal function after AKI. However, we must take into account that the aforementioned authors followed the long-term effects of co-morbidities on the recovery of renal function.

In a cohort analysis of the 131 critically ill patients with AKI who underwent the CRRT, along with the older age, two other determined predictors that we did not examine (coronary artery disease and admission to the ICU) were associated with a lower rate of renal function recovery [11].

While our study provides valuable insights, it is important to acknowledge the limitations associated with it. Firstly, this study was conducted retrospectively in a single center, which may limit the generalizability of the findings to other settings or patient populations.

Additionally, we lacked specific data on the etiology of AKI-D, diuresis, and the use of diuretics. These factors could potentially influence the recovery of renal function and should be considered in future studies for a more comprehensive analysis.

Furthermore, we did not have detailed information on the parameters of the CRRT procedure. Factors such as the specific modality of CRRT, dialysis dose, choice of dialysis membrane, and duration of dialysis could potentially impact the chances of renal function recovery. Although there is no definitive evidence linking these factors to the recovery of renal function, their potential influence cannot be entirely ruled out.

Despite these limitations, our study contributes valuable information on the predictors of renal function non-recovery in critically ill patients with dialysis-dependent AKI. It underscores the importance of individual consideration of potential predictors and highlights the need for further research to address the limitations and expand our knowledge in this area [25, 26, 27].

CONCLUSION

After conducting both group and subgroup analyses, it was found that among all the predictors investigated, previous kidney conditions pose the highest risk for the failure to recover kidney function in critically ill patients with dialysis-dependent AKI. It is important to look at individual and overall predictors so that clinicians can assess the potential for recovery of kidney function and implement appropriate interventions to improve outcomes.

ACKNOWLEDGEMENTS

We would like to thank dr Dragana Milijašević for assistance with statistical analysis.

Conflict of interest: None declared.

REFERENCES

1. Bani Hani A, Abu Abeeleh M, Al-Najjar S, Alzibdeh A, Mansour S, Bsisu I et al. Incidence, risk factors and outcomes of acute kidney injury in surgical intensive care unit octogenarians at the Jordan University Hospital. *BMC Geriatr.* 2023;23(1):266. [DOI: 10.1186/s12877-023-03975-2] [PMID: 37142956]
2. Singh J, Singh S. Review on kidney diseases: types, treatment and potential of stem cell therapy. *Ren Replace Ther.* 2023;9(1):21. [DOI: 10.1186/s41100-023-00475-2] [PMID: 37131920]
3. Komaru Y, Oguchi M, Sadahiro T, Nakada TA, Hattori N, Moriguchi T et al. Urinary neutrophil gelatinase-associated lipocalin and plasma IL-6 in discontinuation of continuous venovenous hemodiafiltration for severe acute kidney injury: a multicenter prospective observational study. *Ann Intensive Care.* 2023;13(1):42. [DOI: 10.1186/s13613-023-01137-6] [PMID: 37184598]
4. Opgenorth D, Bagshaw SM, Lau V, Graham MM, Fraser N, Klarenbach S et al. A study protocol for improving the delivery of acute kidney replacement therapy (KRT) to critically ill patients in Alberta - DIALYZING WISELY. *BMC Nephrol.* 2022;23(1):369. [DOI: 10.1186/s12882-022-02990-6] [PMID: 36384465]
5. Sancho-Martínez SM, Casanova AG, Düwel AG, Rivero-García K, García-Garrido T, Morales AI et al. Identification of Pre-Renal and Intrinsic Acute Kidney Injury by Anamnestic and Biochemical Criteria: Distinct Association with Urinary Injury Biomarkers. *Int J Mol Sci.* 2023;24(3):1826. [DOI: 10.3390/ijms24031826] [PMID: 36768149]
6. Lee BJ, Hsu CY, Parikh R, McCulloch CE, Tan TC, Liu KD et al. Predicting Renal Recovery After Dialysis-Requiring Acute Kidney Injury. *Kidney Int Rep.* 2019;4(4):571–81. [DOI: 10.1016/j.ekir.2019.01.015] [PMID: 30993232]
7. Lam ICH, Wong CKH, Zhang R, Chui CSL, Lai FTT, Li X et al. Long-term post-acute sequelae of COVID-19 infection: a retrospective, multi-database cohort study in Hong Kong and the UK. *EClinicalMedicine.* 2023;60:102000. [DOI: 10.1016/j.eclinm.2023.102000] [PMID: 37197226]
8. Yang JN, Li Z, Wang ML, Li XY, Li SL, Li N. Preoperative dipstick albuminuria is associated with acute kidney injury in high-risk patients following non-cardiac surgery: a single-center prospective cohort study. *J Anesth.* 2022;36(6):747–56. [DOI: 10.1007/s00540-022-03113-z] [PMID: 36178550]
9. Ruan X, Li M, Pei L, Lan L, Chen W, Zhang Y et al. Association of intraoperative hypotension and postoperative acute kidney injury after adrenalectomy for pheochromocytoma: a retrospective cohort analysis. *Perioper Med (Lond).* 2023;12(1):17. [DOI: 10.1186/s13741-023-00306-2] [PMID: 37194032]
10. Jiang Y, Chen J, Yu Y, Yang F, Hamza M, Zou P et al. Risk factors for the in-hospital mortality of CRRT-therapy patients with cardiac surgery-associated AKI: a single-center clinical study in China. *Clin Exp Nephrol.* 2022;26(12):1233–9. [DOI: 10.1007/s10157-022-02274-1] [PMID: 36083528]
11. Kahindo CK, Mukuku O, Mokoli VM, Sumaili EK, Wembonyama SO, Tsongo ZK. Predictors of Mortality in Adults with Acute Kidney Injury Requiring Dialysis: A Cohort Analysis. *Int J Nephrol.* 2022;2022:7418955. [DOI: 10.1155/2022/7418955] [PMID: 36132538]
12. Yang G, Tan L, Yao H, Xiong Z, Wu J, Huang X. Long-Term Effects of Severe Burns on the Kidneys: Research Advances and Potential Therapeutic Approaches. *J Inflamm Res.* 2023;16:1905–21. [DOI: 10.2147/JIR.S404983] [PMID: 37152866]
13. Schmid N, Ghinescu M, Schanz M, Christ M, Schrickler S, Ketteler M et al. Algorithm-based detection of acute kidney injury according to full KDIGO criteria including urine output following cardiac surgery: a descriptive analysis. *BioData Min.* 2023;16(1):12. [DOI: 10.1186/s13040-023-00323-3] [PMID: 36927544]
14. Li Y, Li H, Zhang D. Timing of continuous renal replacement therapy in patients with septic AKI: A systematic review and meta-analysis. *Medicine (Baltimore).* 2019;98(33):e16800. [DOI: 10.1097/MD.00000000000016800] [PMID: 31415389]
15. Lin WT, Lai CC, Chang SP, Wang JJ. Effects of early dialysis on the outcomes of critically ill patients with acute kidney injury: a systematic review and meta-analysis of randomized controlled trials. *Sci Rep.* 2019;9(1):18283. [DOI: 10.1038/s41598-019-54777-9] [PMID: 31797991]
16. Ponce D, Zamoner W, Addad V, Batistoto MM, Balbi A. Acute Renal Replacement Therapy in Intensive Care Units versus Outside Intensive Care Units: Are They Different? *Int J Nephrol Renovasc Dis.* 2020;13:203–9. [DOI: 10.2147/IJNRD.S251127] [PMID: 32943905]
17. Knežević V, Azaševac T, Simin Šibalić M, Sladojević V, Urošević I, Čelić D. Early initiation of renal replacement therapy improves survival in patients with acute kidney injury. *Vojnosanit Pregl* 2021; 78(10):1028–35. [DOI: 10.2298/VSP191117019K]
18. Castro I, Relvas M, Gameiro J, Lopes JA, Monteiro-Soares M, Coentrão L. The impact of early versus late initiation of renal replacement therapy in critically ill patients with acute kidney injury on mortality and clinical outcomes: a meta-analysis. *Clin Kidney J.* 2022;15(10):1932–45. [DOI: 10.1093/ckj/sfac139] [PMID: 36158157]
19. Joannidis M, Forni LG, Klein SJ, Honore PM, Kashani K, Ostermann M et al. Lung-kidney interactions in critically ill patients: consensus report of the Acute Disease Quality Initiative (ADQI) 21 Workgroup. *Intensive Care Med.* 2020;46(4):654–72. [DOI: 10.1007/s00134-019-05869-7] [PMID: 31820034]

20. Yang C, Guo J, Ni K, Wen K, Qin Y, Gu R et al. Mechanical Ventilation-Related High Stretch Mainly Induces Endoplasmic Reticulum Stress and Thus Mediates Inflammation Response in Cultured Human Primary Airway Smooth Muscle Cells. *Int J Mol Sci.* 2023;24(4):3811. [DOI: 10.3390/ijms24043811] [PMID: 36835223]
21. Ostermann M, Lumlertgul N. Acute kidney injury in ECMO patients. *Crit Care.* 2021;25(1):313. [DOI: 10.1186/s13054-021-03676-5] [PMID: 34461966]
22. Infante B, Conserva F, Pontrelli P, Leo S, Stasi A, Fiorentino M et al. Recent advances in molecular mechanisms of acute kidney injury in patients with diabetes mellitus. *Front Endocrinol (Lausanne).* 2023;13:903970. [DOI: 10.3389/fendo.2022.903970] [PMID: 36686462]
23. Chung MC, Hung PH, Hsiao PJ, Wu LY, Chang CH, Hsiao KY et al. Sodium-Glucose Transport Protein 2 Inhibitor Use for Type 2 Diabetes and the Incidence of Acute Kidney Injury in Taiwan. *JAMA Netw Open.* 2023;6(2):e230453. [DOI: 10.1001/jamanetworkopen.2023.0453] [PMID: 36811856]
24. Vijayan A, Abdel-Rahman EM, Liu KD, Goldstein SL, Agarwal A, Okusa MD et al. AKI!NOW Steering Committee. Recovery after Critical Illness and Acute Kidney Injury. *Clin J Am Soc Nephrol.* 2021;16(10):1601–9. [DOI: 10.2215/CJN.19601220] [PMID: 34462285]
25. Carey LI, Kaimba S, Nyirenda S, Chetcuti K, Joeke E, Henrion MYR, et al. Prospective cohort study to identify prevalence, risk factors and outcomes of infection associated kidney disease in a regional hospital in Malawi. *BMJ Open.* 2022;12(11):e065649. [DOI: 10.1136/bmjopen-2022-065649] [PMID: 36442901]
26. Wang M, Wang X, Zhu B, Li W, Jiang Q, Zuo Y, et al. The effects of timing onset and progression of AKI on the clinical outcomes in AKI patients with sepsis: a prospective multicenter cohort study. *Ren Fail.* 2023;45(1):1-10. [DOI: 10.1080/0886022X.2022.2138433] [PMID: 37096423]
27. Soum E, Timsit JF, Ruckly S, Gruson D, Canet E, Klouche K, et al. Predictive factors for severe long-term chronic kidney disease after acute kidney injury requiring renal replacement therapy in critically ill patients: an ancillary study of the ELVIS randomized controlled trial. *Crit Care.* 2022;26(1):367. [DOI: 10.1186/s13054-022-04233-4] [PMID: 36447221]

Table 1. Demographic and clinical parameters in critically ill patients and its association with renal recovery function

Variables	Renal function non-recovery n/%	Renal function recovery n/%	p
	242 (55)	198 (45)	
Sex			p = 0.264
Male	138/69.7	156/64.5	
Female	60/30.3%	86/35.5	
Mean age-years	63.14	60.93	
Comorbidities			
Hypertension	86 (43.4)	103 (42.6)	p = 0.923
Chronic pulmonary disease	25 (12.5)	25 (10.3)	p = 0.455
Diabetes mellitus	43 (21.7)	51 (21.1)	p = 0.907
Cerebrovascular disease	15 (7.6)	23 (9.5)	p = 0.500
Digestive disease	26 (13.1)	29 (12.0)	p = 0.773
Previous kidney disease	16 (8.1)	4 (1.7)	p = 0.002*
MV support	61 (69.3)	65 (64.4)	p = 0.537
Vasopressor support	138 (69.7)	139 (57.4)	p = 0.010*
Start of CRRT ≤ 24 hours	121 (61.1)	170 (70.2)	p = 0.050*
qSOFA score			
0	47(23.7)	76 (31.4)	p = 0.296
1	53 (26.8)	64 (26.4)	
2	26 (13.1)	25 (10.3)	
3	72 (36.4)	77 (31.8)	
Types of CRRT modalities			
CVVHDF	81 (40.9)	88 (36.4)	p = 0.282
CVVHD	72 (36.4)	91 (37.6)	
CVVH	8 (4)	14 (5.8)	
CVVHD + CVVHDF	37 (18.7)	49 (20.2)	
Urea (mmol) + (Mean, SD)	26.32 ± 15.306	26.33 ± 13.962	p = 0.995
Creatinine (μmol) (Mean, SD)	401.43 ± 203.56	373.70 ± 191.51	p = 0.145
CRP (mg/L) (Mean, SD)	153.99 ± 121.44	154.62 ± 126.79	p = 0.966
PCT (ng/l) (Mean, SD)	42.80 ± 68.88	42.59 ± 94.32	p = 0.987
Ultrafiltration (ml) (Mean, SD)	3367.88 ± 5066.81	2592.8 ± 2946.46	p = 0.046*

CRRT – continuous renal replacement therapy; CVVHDF – continuous veno-venous hemodiafiltration; CVVH – continuous veno-venous hemofiltration; CVVHD – continuous veno-venous hemodialysis; CRP – C-reactive protein; PCT – procalcitonin; MV – mechanical ventilation; qSOFA score – quick sequential organ failure assessment score

Table 2. Predictors of renal function non-recovery in critically ill patients with AKI-D with demographic, laboratory and clinical parameters

Parameters	Exp(B)	95% C.I. for EXP(B)		Sig.
		Lower	Upper	
Gender	1.472	0.940	2.304	0.091
Age	1.437	1.010	2.043	0.044*
Ultrafiltration	1.081	0.767	1.524	0.657
Modalities of CRRT	0.950	0.787	1.147	0.595
Urea at admission	1.079	0.860	1.353	0.512
Creatinine at admission	1.004	0.979	1.031	0.739
CRP	1.424	0.880	2.304	0.150
PCT	0.620	0.371	1.038	0.069
Length of hospitalization	1.070	0.763	1.502	0.694
Oliguria/ Anuria	0.649	0.414	1.018	0.060
Start of CRRT \leq 24 hours	1.353	1.009	1.814	0.043*
Vasopressors support	0.958	0.507	1.808	0.894
MV support	1.633	1.013	2.633	0.044*
Sepsis	0.247	0.732	0.431	1.242
CVD	0.886	0.560	1.402	0.606
Pulmonary diseases	1.306	0.850	2.007	0.223
Digestive disease	1.191	0.629	2.254	0.592
Diabetes mellitus	1.299	0.766	2.201	0.331
CD	0.753	0.355	1.599	0.460
Previous kidney disease	5.490	1.661	18.148	0.005*
Other comorbidities	0.901	0.563	1.443	0.664
qSOFA score	1.111	0.699	1.763	0.657
Constant	0.173			0.167

CRRT – continuous renal replacement therapy; CRP – C-reactive protein; PCT – procalcitonin;

MV – mechanical ventilation; CD – cerebrovascular diseases; CVD – cardiovascular disease;

qSOFA score – quick sequential organ failure assessment score

Table 3. Association between non-recovery of renal function in septic critically ill patients and AKI-D with demographic, laboratory and clinical parameters

Parameters	Exp(B)	95% C.I. for EXP(B)		Sig.
		Lower	Upper	
Gender	1.728	0.833	3.582	0.142
Age	2.551	1.422	4.579	0.002*
Ultrafiltration	1.128	0.616	2.066	0.696
Modalities of CRRT	0.997	0.701	1.417	0.986
Urea at admission	1.137	0.749	1.726	0.547
Creatinine at admission	1.012	0.965	1.063	0.616
CRP	2.488	1.077	5.748	0.033*
PCT	2.841	1.279	6.308	0.010*
Length of hospitalization	1.234	0.760	2.002	0.395
Oliguria/ Anuria	0.818	0.380	1.759	0.607
Start of CRRT ≤ 24 hours	1.522	0.926	2.501	0.098
Vasopressors support	0.999	0.296	3.374	0.999
MV support	1.572	0.577	4.284	0.377
CVD	1.391	0.626	3.088	0.418
Pulmonary diseases	1.904	0.776	4.670	0.159
Digestive disease	2.604	0.874	7.757	0.086
Diabetes mellitus	2.709	1.144	6.414	0.023*
CD	1.562	0.371	6.583	0.543
Previous kidney disease	7.077	1.046	47.894	0.045*
Other comorbidities	1.522	0.699	3.314	0.290
qSOFA score	0.763	0.354	1.645	0.489
Constant				

CRRT – continuous renal replacement therapy; CRP – C-reactive protein; PCT – procalcitonin;

MV – mechanical ventilation; CD – cerebrovascular diseases; CVD – cardiovascular diseases;

qSOFA score – quick sequential organ failure assessment score

Table 4. Association between non-recovery of renal function in non-septic critically ill patients with AKI-D with demographic, laboratory and clinical parameters

Parameters	Exp(B)	95% C.I. for EXP(B)		Sig.
		Lower	Upper	
Gender	1.541	0.816	2.911	0.183
Age	0.981	0.604	1.594	0.939
Ultrafiltration	0.852	0.531	1.367	0.507
Modalities of CRRT	0.938	0.735	1.197	0.608
Urea at admission	1.040	0.776	1.394	0.794
Creatinine at admission	1.002	0.969	1.037	0.887
CRP	1.093	0.578	2.070	0.784
Length of hospitalization	0.776	0.420	1.435	0.419
Oliguria/ Anuria	1.985	1.053	3.743	0.034*
Start of CRRT ≤ 24 hours	1.321	0.872	2.002	0.189
Vasopressors support	0.814	0.402	1.647	0.567
MV support	1.423	0.749	2.702	0.282
CVD	0.640	0.338	1.211	0.170
Pulmonary diseases	0.756	0.310	1.843	0.538
Digestive disease	0.840	0.344	2.052	0.702
Diabetes mellitus	0.822	0.389	1.737	0.607
CD	0.621	0.238	1.619	0.330
Previous kidney disease	6.251	1.140	34.273	0.035*
Other comorbidities	0.632	0.322	1.242	0.183
qSOFA score	1.440	0.746	2.780	0.277
Constant	0.524			0.703

CRRT – continuous renal replacement therapy; CRP – C-reactive protein; MV – mechanical ventilation; CD – cerebrovascular diseases; CVD – cardiovascular diseases; qSOFA score – quick sequential organ failure assessment score