



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

History of Medicine / Историја медицине

Srđan S. Putnik^{1,*}, Miroslav D. Ilić^{2,3}, Mia Manojlović³

Development of bariatric/metabolic surgery in Vojvodina

Развој баријатријске/метаболичке хирургије у Војводини

¹Vršac General Hospital, Department of General Surgery, Vršac, Serbia;

²Institute for Pulmonary Diseases of Vojvodina, Clinic for Thoracic Surgery, Sremska Kamenica, Serbia;

³University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia

Received: February 12, 2022

Revised: March 25, 2022

Accepted: April 13, 2022

Online First: April 20, 2022

DOI: <https://doi.org/10.2298/SARH220212040P>

***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

***Correspondence to:**

Srdjan S. PUTNIK

Vršac General Hospital, Department of General Surgery, Abraševićeva 13, 26300 Vršac, Serbia

E mail: putniksrdjan@outlook.com

Development of bariatric/metabolic surgery in Vojvodina

Развој баријатријске/метаболичке хирургије у Војводини

SUMMARY

The paper presents the chronological development of bariatric/metabolic surgery in Vojvodina, the region with the largest incidence of obesity in Serbia, and in which 13.2% of the population suffers from diabetes with a mortality of 15.7 / 100.000, which is one of the highest rates in Europe.

Bariatric surgery began in the 1950s in the United States, with the consecutive development of various restrictive, malabsorptive, and combined procedures, which are intensified by the adoption of laparoscopic techniques.

After initial, European knowledge and preparation of obese patients for surgery at the end of the 1970s, the first laparoscopic bariatric/metabolic interventions started in Vojvodina at the Clinical Center in Novi Sad in 2006. Next year, the multidisciplinary team for bariatric surgery is being prepared in Slovenia and Austria, and the first interventions are performed in 2008 at the Clinic for Thoracic Surgery of the Institute in Sremska Kamenica, in the first place - laparoscopic sleeve gastrectomy. Since then, bariatric/metabolic surgery at this Institute has been performed continuously. By establishing international cooperation, three courses were held with demonstration operations in bariatric/metabolic surgery and the participation of eminent surgeons and other experts from the country and abroad. Several bariatric surgeries were performed on children (Children's Surgery Clinic in Novi Sad), as well as in Sremska Mitrovica.

Their own experience in this field has been published in the international literature and through several regional meetings dedicated to metabolic surgery.

Keywords: bariatric surgery; laparoscopic sleeve gastrectomy; mini gastric bypass

САЖЕТАК

У раду је дат хронолошки развој баријатријске/метаболичке хирургије у АП Војводини, регији која има највећу инциденцу гојазних особа и у којој 13,2% популације болује од шећерне болести са морталитетом од 15,7/100.000 становника, што је једна од највиших стопа у Европи. Баријатријска хирургија почиње педесетих година прошлог века у САД, са консекутивним развојем различитих рестриктивних, малапсорптивних и комбинованих захвата који се интензивирају усвајањем лапароскопских техника.

Након почетних, европских сазнања и припреме гојазних пацијената за операцију крајем седамдесетих година, прве лапароскопске баријатријске/метаболичке интервенције започињу у АП Војводини у Клиничком центру у Новом Саду 2006. године постављањем „шведске прилагодљиве траке“. Следеће године мултидисциплинарни тим за баријатријску хирургију припрема се у Словенији и Аустрији, те се прве интервенције лапароскопске рукавне ресекције желуца изводе 2008. године на Клиници за грудну хирургију, Института у Сремској Каменици. Од тада се баријатријска/метаболичка хирургија на овом месту изводи у континуитету. Успостављајући међународну сарадњу, одржана су три курса са показним операцијама из баријатријске/метаболичке хирургије и учешћем еминентних хирурга и других експерата из земље и иностранства. Неколико баријатријских операција изведено је у Сремској Митровици као и код деце на Клиници за дечију хирургију у Новом Саду.

Сопствена искуства из ове области публикована су у међународној литератури и представљена на велики број регионалних састанака посвећених метаболичкој хирургији.

Кључне речи: баријатријска хирургија; лапароскопска рукавна ресекција желуца; мини желудачно премошћавање

INTRODUCTION

Bariatric surgery or weight loss surgery is considered different operative procedures with the aim of treating obesity. The word bariatric has origin from the Greek root of the two words which mean weight and treatment [1]. Obesity is a serious public health problem [2]. People who have elevated BMI have a significantly increased risk of developing cardiovascular, endocrine and infectious diseases, diabetes mellitus, cancer and asthma, as well as sleep apnea,

musculoskeletal disease and sudden death [3, 4, 5]. In addition, obesity affects the social, economic and psychological function of a person in society. Although conservative treatment is first in the treatment of obesity and includes a diet, exercise, lifestyle change, and anti-obesity drugs, it is not successful in a long way [6]. Today, the surgical treatment of obesity and associated "metabolic syndrome" is recommended by the World Health Organization, but also by endocrine and other international associations [7, 8]. Due to the effects on metabolism, primarily on the metabolism of glucose and fat, bariatric surgery is also called metabolic surgery. Vojvodina is according to the epidemiological and social medical indicators of the regions in Serbia and Europe, especially and most affected by obesity and associated metabolic syndrome, primarily diabetes and hypertension. According to the Institute of Public Health Institute in Serbia report of 2016, 13.2% of the adult population suffers from diabetes mellitus type 2, with a mortality rate of 15.7 / 100.000, which represents the highest rate in Europe. Vojvodina and its "districts" (Bačka, Central-Banat and South-Banat district) have the highest incidence of diabetes, over 160:100,000 [9, 10]. In the past decades sporadically and relatively constant in the last ten years, bariatric/metabolic surgery in Vojvodina has been performed.

The aim of the paper is to present the development of bariatric/metabolic surgery in Vojvodina.

THE HISTORY OF BARIATRIC/METABOLIC SURGERY IN THE WORLD

The development of open bariatric surgery began in the fifties of the last century, using the intestinal bypass [11]. In this procedure, the upper part of the intestine (jejunum) merged with the lower part of the small intestine (ileum), which left a large part of the small intestine to be excluded from digestion and caused weight loss by malabsorption [12]. Thereafter in the sixties years of twentieth century, a jejuno-colic bypass was made: the proximal small intestine is transected and anastomosed to the colon [13]. In 1967, on the basis of experiments in the laboratory, the "father" of bariatric surgery Edward Mason developed the original gastric bypass [14]. In 1977, following the work of Alden (who made gastro-jejunostomy - a surgical procedure in which an anastomosis is created between the stomach and the proximal loop of the jejunum), Griffen introduced Roux-en-Y anastomosis into bariatric surgery and this procedure became a gold standard in bariatric/metabolic surgery [11, 15].

As a combined restrictive and malabsorptive procedure, the Roux-en-Y gastric bypass was followed by dumping syndrome, iron deficiency and vitamin B12 deficiency. In the seventies and eighties in the last century, purely restrictive surgery – gastroplasty - was brought to bariatric surgery [16]. The procedures on the stomach and its reduction were followed by the set of rings or staples [17]. These operations showed fewer complications but were accompanied by minor weight loss.

The modern era in bariatric surgery begins with the operation of the biliopancreatic diversion performed by the Italian surgeon - Nicola Scopinaro. The first report on 18 patients was published in 1979 [18]. He performed a horizontal hemi-gastrectomy with pouch drainage by a Roux limb of at least 250 cm length, joining a long biliopancreatic limb to form 50 cm long the common ileal segment, or channel. This procedure was converted by Picard Marceau et al. [19] in Canada to a vertical sleeve gastrectomy with cross-stapling of the duodenum and an approximately 100-cm common channel.

In 1998, Hess and Hess in the USA made a duodenal switch, by performing a sleeve gastrectomy, with pyloric preservation, duodenal division, a proximal duodeno-ileostomy, and a common channel of approximately 100 cm [20]. In this version of the biliopancreatic diversion called "biliopancreatic diversion with duodenal separation" (BPD-DS), which began to be performed laparoscopic, the volume of the stomach was reduced to about 200 millilitres, drawing the future sleeve gastrectomy.

Kuzmak introduced a silicone tape in 1986 that could be filled through the subcutaneous chamber and "adjusted" with clamping around the cardia area, preventing excessive filling of the stomach [21]. Forsell (1993) first set up a laparoscopic "adjustable gastric band" [22].

Laparoscopic sleeve gastrectomy (LSG) or vertical gastric resection was first performed by Michel Gagner in 1999 within the BPD-DS. In 2003, the same surgeon did the LSG as the first stage of two-stage Roux-en-Y gastric bypass (RYGB) surgery in patients at high risk and BMI higher than 50 kg / m². Shortly after this description, the articles followed in which the LSG was performed as an independent and sufficient procedure for the treatment of obesity and metabolic disorders due to good results [23, 24]. Today, this operation is the most commonly performed procedure in metabolic surgery in the whole world [25].

DEVELOPMENT OF BARIATRIC/METABOLIC SURGERY IN VOJVODINA

The first theoretical knowledge on bariatric surgery and the first preparation of patients were done in Novi Sad by the associate professor of surgery Borislav Savić, who worked in the Surgical Clinic from 1979 to 1981. He came from Germany as an affiliated surgeon and author of the book "Septic Surgery", published in Stuttgart [26]. The cooperation of surgeons and internist-endocrinologists was done on the Faculty of Medicine in Novi Sad. The sudden death of Professor Savić stopped the realization of the first bariatric interventions, but the idea continued to live in the followed generations of surgeons and internists. The first laparoscopic interventions on biliary tract were carried out in 1996 in Sremska Kamenica at the Clinic for Cardiovascular Surgery by Professor Branislav Daničić and Dr Miroslav Ilić, who were abdominal surgeons (equipment for laparoscopic procedures was only available to this institution). After the introduction of laparoscopic methods in the Clinical Center of Vojvodina at the Clinic for Abdominal and Endocrine Surgery and the complete experience in this surgery (laparoscopic adrenalectomy, laparoscopic reparation of hiatal hernia, laparoscopic appendectomy), the first operation of setting up a "Swedish adjustable gastric band" (SAGB - Johnson & Johnson®) Professor Radovan Cvijanović was performed with his team in 2006. He continued this procedure to work in a private surgical hospital "Parks" in Novi Sad. In 2007, after obtaining the necessary equipment for bariatric procedures at the Institute for Lung Diseases at the Clinic for Thoracic Surgery, the multidisciplinary team (surgeons, psychologist, nutritionist, anesthesiologist, nurses) was in Slovenj Gradec (Slovenia) for conducting bariatric operations under the management of Dr Brane Breznikar.

Immediately afterward, courtesy of Johnson & Johnson medical company, Dr Miroslav Ilić was at the Clinic for Bariatric Surgery under the mentorship of Associate Professor Karl Miller (Head of the Surgical Department, Hallein Clinic, Austria), where he was educated primarily for the SAGB procedure and Roux-en-Y gastric bypass.

The first bariatric/metabolic laparoscopic procedures were performed at the Clinic for Thoracic Surgery of the Institute of Pulmonary Diseases of Vojvodina on 31 October 2008, with the guidance of Dr Miller, two SAGB procedures and one laparoscopic sleeve gastrectomy on mega - obese patient (BMI 70 kg/m²) [27]. At the same time, Professor Miller held a public lecture about bariatric medicine and surgery in front of representatives of provincial authorities

in order to point out the financial advantages of early treatment of obesity and metabolic syndrome.

The first accredited two-day multidisciplinary Alma Mons course (Continuing Medical Education – CME) was held in October 2013 with lecturers from Serbia, Bosnia and Herzegovina, Croatia, Romania and Turkey, with live performance and monitoring with the possibility of asking questions to surgeons during the procedure. The lecturers were: internist endocrinologists Professor Edita Stokic (Novi Sad, Serbia), Dr Snežana Polovina (Belgrade, Serbia), internist pulmonologist Assistant Professor Ivan Kopitović, anesthesiologists Dr Milana Komarčević (Sremska Kamenica, Serbia) and surgeons Professor Miroslav Bekavac-Beslin (Zagreb, Croatia), Assistant Professor Fuad Pašić (Tuzla, Bosnia and Herzegovina), Professor Catalin Copăescu (Bucharest, Romania), Professor Alper Celik (Istanbul, Turkey), Assistant Professor Miloš Koledin and Professor Miroslav Ilić (Sremska Kamenica, Serbia). The following operations were performed: laparoscopic "resleeve" gastrectomy and laparoscopic ileal interposition. Both of these operations were first performed in Serbia.

The second Alma Mons course was held in November 2014 with the participation of surgeons from the Netherlands, Croatia and Serbia. The lecturers were Dr Arlond van de Laar, Head of Bariatric Surgery at the Slotervaart Hospital in Amsterdam (Netherlands), the European Center of Excellence for Bariatric Surgery, with associates Dr Yair Acherman and Dr M. Hoen, whose focus was on Roux-en-Y gastric bypass, then Professor Davor Štimac (Clinical Hospital Center Rijeka, Croatia) - about intragastric balloons. Cooperation and meeting with Professor Bešlin-Bekavac (University Hospital Center "Sestre Milosrdnice", Zagreb, Croatia) and Assistant Professor F. Pašić (University Clinical Center Tuzla, Tuzla, Bosnia and Herzegovina). The third Alma Mons course was dedicated to only one surgeon and surgery: Dr Robert Rutledge (Center for Excellence in Laparoscopic Obesity Surgery, Henderson, USA) and his metabolic laparoscopic procedure - "mini-gastric bypass" (MGB/OAGB). This course was held on May 30, 2016, with the theoretical part and the performance of two MGB/OAGB operations carried out by Dr Robert Rutledge and Professor Miroslav Ilić (Figure 1). These interventions were performed for the first time in the region of Southeast Europe.

From 2008 onwards, bariatric/metabolic surgeries are continuously performed at the Clinic for Thoracic Surgery in Sremska Kamenica under the leadership of Professor Miroslav Ilić [28].

According to the data from the operational protocols and electronic patient records, in the period from 2008 to 2020, about 500 bariatric procedures were performed.

The most commonly performed procedure is laparoscopic sleeve gastrectomy with 380 operations. Complications included three postoperative bleeding, four anastomotic leaks, two chronic fistulas and one neofundus that were resolved surgically and one fistula that was treated conservatively.

Other bariatric procedures have been performed: 18 Swedish adjustable gastric bands, 9 laparoscopic mini-gastric bypass, 5 Roux-en-Y gastric bypass, 3 laparoscopic "resleeve" gastrectomy, 1 laparoscopic ileal interposition, 2 single anastomosis sleeve ileal bypass and more open LSG and mini-gastric bypass.

Professor Radoica Jokić, a pediatric and abdominal surgeon, has performed several bariatric interventions on children (laparoscopic sleeve gastrectomy), with his colleague Professor Dragan Kravarušić (Schneider Children's Medical Center, Israel), with excellent results at the Children's Surgery Clinic in Novi Sad. Attempts to carry out the SAGB bariatric procedure were also recorded in the General Hospital of Sremska Mitrovica by Dr Goran Ivić in 2010 but without further continuity.

The results of their own experiences and guest appearances abroad were presented in several publications, as well as doctoral theses, in which the general surgeon from Vršac was very diligent, Dr Srđan Putnik (General Hospital Vršac) [27, 28, 29].

Apart from Vojvodina, bariatric surgery is also performed at the University Clinical Center of Serbia. Thanks to Academician Professor Dragan Micić, the Multidisciplinary Center for Obesity Treatment was opened in 2010. Of the bariatric procedures, RYGB and LSG are the most common. According to the data from the literature, about 100 Roux-en-Y gastric bypass were performed [30].

Other centers in Serbia are also showing great interest in this type of increasingly popular surgery.

ACKNOWLEDGMENT

This paper is part of a doctoral thesis by Srđan Putnik, for which the permission of the Ethics Committee of the Institute for Pulmonary Diseases of Vojvodina, Sremska Kamenica, was given (No. 72-XIII/24). The authors thank Prof. R. Jokić and Prof. R. Cvijanović for the obtained data.

Conflict of interest: None declared.

Paper accepted

REFERENCES

1. Baltasar A, Bou R, Bengochea M, Pérez N. Beginnings of bariatric and metabolic surgery in Spain. *JONNPR*. 2020; 5(4):448-67. [DOI: 10.19230/jonnpr.3324]
2. Jaacks LM, Vandevijvere S, Pan A, McGowan CJ, Wallace C, Imamura F, et al. The obesity transition: stages of the global epidemic. *Lancet Diabetes Endocrinol*. 2019; 7(3):231-240. [DOI: 10.1016/S2213-8587(19)30026-9] [PMID: 30704950]
3. La Sala L, Pontiroli AE. Prevention of Diabetes and Cardiovascular Disease in Obesity. *Int J Mol Sci*. 2020; 21(21):8178. [DOI: 10.3390/ijms21218178] [PMID: 33142938]
4. Peters U, Dixon AE, Forno E. Obesity and asthma. *J Allergy Clin Immunol*. 2018; 141(4):1169-1179. [DOI: 10.1016/j.jaci.2018.02.004] [PMID: 29627041]
5. Krupa-Kotara K, Dakowska D. Impact of obesity on risk of cancer. *Cent Eur J Public Health*. 2021; 29(1):38-44. [DOI: 10.21101/cejph.a5913] [PMID: 33831285]
6. Gadde KM, Martin CK, Berthoud HR, Heysmsfield SB. Obesity: Pathophysiology and Management. *J Am Coll Cardiol*. 2018; 71(1):69-84. [DOI: 10.1016/j.jacc.2017.11.011] [PMID: 29301630]
7. Affinati AH, Esfandiari NH, Oral EA, Kraftson AT. Bariatric Surgery in the Treatment of Type 2 Diabetes. *Curr Diab Rep*. 2019; 19(12):156. [DOI: 10.1007/s11892-019-1269-4] [PMID: 31802258]
8. Pratt JSA, Browne A, Browne NT, Bruzoni M, Cohen M, Desai A, et al. ASMBS pediatric metabolic and bariatric surgery guidelines, 2018. *Surg Obes Relat Dis*. 2018; 14(7):882-901. [DOI: 10.1016/j.soard.2018.03.019] [PMID: 30077361]
9. Incidence and mortality of diabetes in Serbia. Institute of Public Health of Serbia "Dr Milan Jovanovic Batut" 2016
10. Atlas of health in Europe. World Health Organization. Regional Office for Europe: Copenhagen, Denmark, 2003
11. Buchwald H, Buchwald JN. Evolution of operative procedures for the management of morbid obesity 1950-2000. *Obes Surg*. 2002; 12(5):705-17. [DOI:10.1381/096089202321019747] [PMID:12448398]
12. Griffen WO Jr, Bivins BA, Bell RM. The decline and fall of the jejunoileal bypass. *Surg Gynecol Obstet*. 1983; 157(4):301-8. [PMID:6623319]
13. Payne JH, DeWind LT, Commons RR. Metabolic observations in patients with jejunocolic shunts. *Am J Surg*. 1963; 106:273-89. [DOI:10.1016/0002-9610(63)90017-5] [PMID:14042557]
14. Mason EE, Ito C. Gastric bypass in obesity. *Surg Clin North Am*. 1967; 47(6):1345-51. [PMID:6073761]
15. Griffen WO Jr, Young VL, Stevenson CC. A prospective comparison of gastric and jejunoileal bypass procedures for morbid obesity. *Ann Surg*. 1977; 186(4):500-9. [PMID:907395]
16. Laws HL. Standardized gastroplasty orifice. *Am J Surg*. 1981; 141(3):393-4. [DOI: 10.1016/0002-9610(81)90206-3] [PMID:7011080]
17. Fobi M. Why the Operation I Prefer is Silastic Ring Vertical Gastric Bypass. *Obes Surg*. 1991; 1(4):423-426. [DOI:10.1381/096089291765560854] [PMID:10775946]
18. Scopinaro N, Gianetta E, Civalleri D, Bonalumi U, Bachi V. Bilio-pancreatic bypass for obesity: II. Initial experience in man. *Br J Surg*. 1979; 66(9):618-20. [DOI: 10.1002/bjs.1800660906] [PMID:497645]
19. Marceau P, Biron S, Bourque RA, Potvin M, Hould FS, Simard S. Biliopancreatic Diversion with a New Type of Gastrectomy. *Obes Surg*. 1993; 3(1):29-35. [DOI:10.1381/096089293765559728] [PMID:10757900]
20. Hess DS, Hess DW. Biliopancreatic diversion with a duodenal switch. *Obes Surg*. 1998; 8(3):267-82. [DOI:10.1381/096089298765554476] [PMID:9678194]
21. Kuzmak, L.I. Silicone gastric banding: a simple and effective operation for morbid obesity. *Contemp Surg*. 1986;28:13-18.
22. Forsell P, Hallberg D, Hellers G. Gastric Banding for Morbid Obesity: Initial Experience with a New Adjustable Band. *Obes Surg*. 1993; 3(4):369-374. [DOI:10.1381/096089293765559052] [PMID:10757948]
23. Castro Vázquez J, Saravia Barahona F, Loureiro González C, Leturio Fernández S, García Fernández M, Moro Delgado A, et al. Sleeve gastrectomy as a surgical technique in bariatric surgery: Results of safety and effectiveness. *Cir Esp (Engl Ed)*. 2022; 100(2):88-94. [DOI: 10.1016/j.cireng.2022.01.002] [PMID: 35131227]
24. Kraljević M, Cordasco V, Schneider R, Peters T, Slawik M, Wölnerhanssen B, et al. Long-term Effects of Laparoscopic Sleeve Gastrectomy: What Are the Results Beyond 10 Years? *Obes Surg*. 2021; 31(8):3427-3433. [DOI: 10.1007/s11695-021-05437-3] [PMID: 33890225]
25. Welbourn R, Hollyman M, Kinsman R, Dixon J, Liem R, Ottosson J, et al. Bariatric Surgery Worldwide: Baseline Demographic Description and One-Year Outcomes from the Fourth IFSO Global Registry Report 2018. *Obes Surg*. 2019; 29(3):782-795. [DOI: 10.1007/s11695-018-3593-1] [PMID: 30421326]
26. Eckert P, Aigner P, Savic B. *Septische Chirurgie*. Stuttgart; New York: Schattauer, 1980.
27. Ilić M, Putnik SS, Raspopović K. Long-term results of laparoscopic gastric sleeve resection due to morbid obesity and metabolic syndrome. *SrpArhCelokLek*. 2018;146(3-4):211-214. [DOI: 10.2298/SARH170425129I]
28. Ilic M, Putnik SS. Surgical Technique: Laparoscopic Gastric Sleeve Resection in Super-Obese Patients. *J Invest Surg*. 2018;31(3):165-167. [DOI: 10.1080/08941939.2017.1289284] [PMID: 28387579]

29. Ilić M, Putnik SS. Laparoscopic sleeve gastrectomy in a superobese patient with restenosis of the trachea. *SrpArhCelokLek*. 2019; 147(3-4):215-217. [DOI: 10.2298/SARH180411052I]
30. Nedeljkovic-Arsenovic O, Banovic M, Radenkovic D, Rancic N, Polovina S, Micic D, et al. Five-Year Outcomes in Bariatric Surgery Patients. *Medicina (Kaunas)*. 2020; 56(12):669. [DOI: 10.3390/medicina56120669] [PMID: 33287109]

Paper accepted



Figure 1. Dr Robert Rutlage and Professor Miroslav Ilić

Paper