



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Case report / Приказ болесника

Slaviša Zagorac^{1,2,*}, Radovan Mijalčić^{1,3}, Miloš Vasić², Uroš Dabetić², Uroš Novaković²

**Occipitocervical fusion as treatment of instability in
Chiari malformation**

Окципитоцервикална фузија у лечењу нестабилности код
Кјари малформације

¹University of Belgrade, Faculty of medicine, Belgrade, Serbia;

²University Clinical Center of Serbia, Clinic for Orthopedic Surgery and Traumatology, Belgrade, Serbia;

³University Clinical Center of Serbia, Clinic for Neurosurgery, Belgrade, Serbia

Received: February 3, 2022

Revised: April 3, 2022

Accepted: April 6, 2022

Online First: April 21, 2022

DOI: <https://doi.org/10.2298/SARH220203041Z>

***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

***Correspondence to:**

Slaviša ZAGORAC

University Clinical Center of Serbia, Clinic for Orthopedic Surgery and Traumatology

Pasterova 2, 11000 Belgrade, Serbia

slavisa.zagorac@gmail.com

Occipitocervical fusion as treatment of instability in Chiari malformation

Окципитоцервикална фузија у лечењу нестабилности код Кјари малформације

SUMMARY

Introduction Occipitocervical (OC) fusion is a method for fixation of the OC junction when there is instability of that segment. Arnold Chiari malformation is a congenital disorder where cerebellar tonsils descend through the foramen magnum, which can lead to cervicomedullar compression and formation of syrinx. While treating this condition, for the purpose of decompression, the foramen magnum is expanded which can potentially harm the stability of the OC junction.

Case outline We are presenting the case of a 16 year old female who was surgically treated (suboccipital craniectomy and decompression) because of Arnold Chiari malformation type I. One-year post-op multislice computed tomography verified a slight enlargement of basion-axial and basion-dens intervals, with signs of spinal cord compression. Surgery was performed – occipitocervical fusion, canal decompression on C1 and C2 levels with a plate on occipital bone and screws placed in the third, fourth, and fifth cervical vertebra.

Conclusion Occipitocervical fusion is a complex surgical procedure (vital neurovascular structures), but it is a reliable method for treatment of instability of the OC junction.

Keywords: occipitocervical instability; fusion; Arnold Chiari

САЖЕТАК

Увод Окципитоцервикална (ОЦ) фузија је метода фиксације окципитоцервикалног прелаза када постоји нестабилност тог сегмента. Арнолд–Кјаријева малформација је конгенитални поремећај где долази до миграције церебеларних тонзила кроз форамен магнум, што може довести до цервикомедуларне компресије и стварања сиринкса. Приликом лечења овог поремећаја, због декомпресије се шири форамен магнум што потенцијално може довести до нестабилности ОЦ прелаза.

Приказ болесника Приказујемо случај пацијенткиње старости 16 година са Арнолд–Кјаријевом малформацијом типа И која је хируршки лечена (субокципитална краниектомија и декомпресија). Годину дана након операције на мултисјајсној компјутеризованом томографији је верификовано благо увећање бацион-аксис и бацион-денс интервала, са знацима компресије кичмене мождине. Учињен је хируршки захват – окципитоцервикална фузија, декомпресија спиналног канала на нивоима Ц1 и Ц2, плоча је позиционирана на окципиталну кост док су шрафови пласирани у Ц3, Ц4 и Ц5 пршљенове.

Закључак Окципитоцервикална фузија је комплексна хируршка процедура (анатомска позиција виталних неуроваскуларних структура), али представља поуздану методу лечења нестабилности ОЦ прелаза.

Кључне речи: окципитоцервикална нестабилност; фузија; Арнолд–Кјари

INTRODUCTION

Occipitocervical fusion is a surgical method for fixation of the OC junction when there is instability of the occipitocervical segment caused by a variety of conditions (trauma, rheumatoid arthritis, congenital anomalies, neoplasm, iatrogenic instability due to decompression) [1].

Arnold Chiari malformation is a congenital disorder where cerebellar tonsils descend through the foramen magnum, which can lead to corticomedullar compression and creation of

syrinx. It can also be associated with numerous anomalies (syringomyelia, spina bifida, hydromyelia, kyphosis, scoliosis and tethered cord syndrome), as well as hereditary syndromes [2].

To this date the treatment protocol is debatable, with some authors preferring conservative treatment to surgery, [3] but available literature is in favor of surgical treatment [4, 5, 6].

While treating this condition, for the purpose of decompression, the foramen magnum is expanded which can potentially harm the stability of the OC junction [7].

CASE REPORT

We are presenting the case of a 16-year-old female who was surgically treated (suboccipital craniectomy and decompression) because of Arnold Chiari malformation type I. She was initially hospitalized because of repetitive episodes of weakness in both arms and legs, that lasted for a few minutes and disappeared spontaneously. They first appeared 6 months prior to hospital admission. After careful examination she was diagnosed with Chiari malformation type I that was magnetic resonance (MRI) verified – a prolapse (herniation) of the cerebellar tonsils through the foramen magnum by 20 mm, with compression of the spinal cord. The ventricular system was in an orderly position and shape, without signs of hydrocephalus. In addition to this obvious malformation, the existence of syringomyelia (starting from C5 and caudally) and malformation of the base of the skull in terms of platybasia and an abnormal angle between the medulla oblongata and mesencephalon were observed (Figure 1).

Follow up nuclear magnetic resonance was performed (Figure 2) following the first hospitalization and the findings were unchanged and the neurosurgical procedure (suboccipital craniectomy and decompression) was performed. Approximately one year postop the patient

presents with following symptoms: headache, left arm paresthesia eventually followed by left side hemiparesis. New MSCT and MRI scans were obtained and showed progression of syringomyelia expanding cranially to syringobulbia, decompression in posterior cranial fossa was still intact. Two months later the symptoms persisted and new MRI scans were obtained – in comparison with former scans progression of hydrosyringomyelia was observed, as well as expanded central medullar canal with oedema. Conservative treatment was tried with antiedema therapy, but there was no clinical improvement. The patient was hospitalized again and another MRI evaluation was performed, and neurosurgeon and orthopedic spinal surgeon indicated occipitocervical stabilization.

Surgery was performed – occipitocervical fusion, canal decompression on C1 and C2 levels with a plate on occipital bone and screws placed in 3rd, 4th and 5th cervical vertebra. Neuromonitoring was used throughout the whole surgery (Figure 3 and 4).

Postoperatively the patient has no significant symptoms, with notable reduction of pain and paresthesia of the left arm. Movements of cervical spine are limited, in accordance with the stated surgery. The patient is feeling well and she is back to her daily activities.

The study was done in accordance with the institutional Committee on Ethics.

DISCUSSION

Occipitocervical fusion is a surgery indicated for treatment of craniocervical junction instability caused by a variety of different pathologies (congenital, traumatic, degenerative, inflammatory, infective, or neoplastic) [1, 8].

This surgery represents a huge challenge for the operator and his team, considering that C0-C1-C2 is the most mobile portion of the spine, and it is the portion of the spine that must resist force in 8 axes of rotation [9].

Each patient that presents with craniocervical junction instability, whatever the cause, must be carefully and minutely assessed, appropriate diagnostic methods used. Only then – while taking into account clinical presentation (functional stability, neurological status, accompanying symptoms), as well as MRI/CT findings – after preoperative planning the decision about the right treatment should be made [10].

The advances made in the field of neuroradiology have made it possible to understand biomechanics and structure of this region, especially considering substantial number of anatomical variations of vascular and neural structures [11]. Today, these radiologic findings are crucial in preoperative planning, and intraoperative visualization.

In 1900s occipitocervical instability was considered inoperable and a terminal condition, and in the last 90 years there was a large number of surgical techniques developed to perform occipitocervical fusion [12].

Despite of the chosen method the main goal is the same – restore and maintain alignment, decompress neural elements and provide good conditions for the bone fusion to occur.

The spine surgeon performing OC fusion must be well aware of spinal biomechanics and anatomy, and he must be familiar with the procedures to achieve decompression, alignment immobilization and fusion. Also, the surgeon should be aware of the perioperative risks and complications rates that are significant [13], and the readmission rates following OC fusion – that although lower in elective OC fusion surgeries aren't negligible [14].

Taking into account the aforementioned, it is clear that indicating this surgery is a big decision and the surgeon must consider all of the methods available as well as evaluate each patient individually before deciding this is the best solution.

While official surgical guidelines are yet to be established advancements are being made and teams consisting of experts in diagnosing and treating this condition were put together to discuss this matter. [15] Most authors agree that a decision should be made based on the clinical

condition of the patient [5–8]. In their research, Asghar Ali Turabi et al. [16] concluded that occipitocervical fusion along with decompression surgery, had better outcome than decompression surgery alone. In order to be able to make the best patient-oriented decision, larger studies focusing on treatment outcomes are necessary as the available studies have shown that outcomes vary relative to a number of factors like chosen surgical procedure, symptoms duration and syringomyelia [17].

In our experience, occipitocervical fusion is a complex surgical procedure (involving vital neurovascular structures), demanding a multidisciplinary approach (spinal surgeon, neurosurgeon, anesthesiologist) but it is a reliable method for treatment of instability of the OC junction. Every case should be carefully considered, and if decided that occipitocervical fusion is the appropriate method of treatment, it requires detailed planning.

Conflict of interest: None declared.

REFERENCES

1. Kukreja S, Ambekar S, Sin A, Nanda A. Occipitocervical Fusion Surgery: Review of Operative Techniques and Results. *J Neurol Surg B Skull Base*. 2015 Sep; 76(5): 331–339
2. Holly L, Batzdorf U. Chiari malformation and syringomyelia. *J Neurosurg Spine*. 2019 Nov 1;31(5):619-628.
3. Landridge B, Philips E, Choi D. Chiari Malformation Type 1: A Systematic Review of Natural History and Conservative Management. *World Neurosurg*. 2017 Aug;104:213-219.
4. Vinchon M. Surgery for Chiari 1 malformation: the Lille experience. *Childs Nerv Syst*. 2019 Oct;35(10):1875-1880.
5. Fric R, Ringstad G, Eide P. [Chiari malformation type 1 – diagnosis and treatment]. *Tidsskr Nor Laegeforen*. 2019 Jun 11;139(10).
6. Chatterjee S, Shivhare P, Verma S. Chiari malformation and atlantoaxial instability: problems of co-existence. *Childs Nerv Syst*. 2019 Oct;35(10):1755-1761.
7. Rocque B, Oakes W. Surgical Treatment of Chiari I Malformation. *Neurosurg Clin N Am*. 2015 Oct;26(4):527-31.
8. Joaquim A, Osorio J, Riew K. Occipitocervical Fixation: General Considerations and Surgical Technique. *Global Spine J*. 2020 Aug;10(5):647-656.
9. Ling FP, Chevillotte T, Leglise A, Thompson W, Bouthors C, Le Huec JC. Which parameters are relevant in sagittal balance analysis of the cervical spine? A literature review. *Eur Spine J*. 2018 Feb;27(Suppl 1):8-15.
10. Macki M, Hamilton T, Pawloski J, Chang V. Occipital fixation techniques and complications. *J Spine Surg*. 2020 Mar;6(1):145-155.
11. Izzo R, Popolizio T, Balzano RF, Simeone A, Gasparotti R, Scarabino T, Muto M. Imaging of cranio-cervical junction traumas. *Eur J Radiol*. 2020 Jun;127:108960.
12. Winegar CD, Lawrence JP, Friel BC, Fernandez C, Hong J, Maltenfort M, Anderson PA, Vaccaro AR. A systematic review of occipital cervical fusion: techniques and outcomes. *J Neurosurg Spine*. 2010 Jul;13(1):5-16.
13. Elia C, Takayanagi A, Arvind V, Goodmanson R, von Glinski A, Pierre C, Sung J, Qutteineh B, Jung E, Chapman J, Oskouian R. Risk Factors Associated with 90-Day Readmissions Following Occipitocervical Fusion: A Nationwide Readmissions Database Study. *World Neurosurg*. 2021 Mar;147:e247-e254. doi: 10.1016/j.wneu.2020.12.031. Epub 2020 Dec 13. PMID: 33321249.
14. Zileli M, Akintürk N. Complications of occipitocervical fixation: retrospective review of 128 patients with 5-year mean follow-up. *Eur Spine J*. 2022 Feb;31(2):311-326. doi: 10.1007/s00586-021-07037-2. Epub 2021 Nov 2. PMID: 34725722.
15. Ciaramitaro P et al; International Experts Jury of the Chiari Syringomyelia Consensus Conference, Milan, November 11-13, 2019. Diagnosis and treatment of Chiari malformation and syringomyelia in adults: international consensus document. *Neurol Sci*. 2022 Feb;43(2):1327-1342. doi: 10.1007/s10072-021-05347-3. Epub 2021 Jun 15. Erratum in: *Neurol Sci*. 2021 Nov 17;; PMID: 34129128.
16. Asghar Ali Turabi MA, Pruthi N, Bhat D, Bhagavatula ID. Complex Chiari Malformation: Proposal of a New Set of Diagnostic Criteria and Outcomes After fusion Surgery. *World Neurosurg*. 2021 Oct;154:e677-e682. doi: 10.1016/j.wneu.2021.07.115. Epub 2021 Jul 31. PMID: 34343684.
17. Basaran R, Bozdogan C, Senol M, Gundogan D, Isik N. Long-term outcomes of surgical management in subtypes of Chiari malformation. *Neurol Res*. 2021 Sep;43(9):760-766. doi: 10.1080/01616412.2021.1934314. Epub 2021 May 30. PMID: 34057045



Figure 1. First magnetic resonance imaging showing herniation of cerebellar tonsils through the foramen magnum (20 mm) with spinal cord compression



Figure 2. First follow up magnetic resonance imaging, before the first surgery




Figure 3. Intraoperative finding – occipital plate and screws



Figure 4. Postoperative X-ray

Paper accepted