



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Case Report / Приказ болесника

Dragan Erić¹, Boris Tadić^{2,3}, Nikola Grubor^{2,3,*}, Borislav Tosković^{3,4},
Vladimir Milosavljević⁴

Giant spleen as a surgical challenge – case report and literature review

Масивна спленомегалија као хируршки изазов –
приказ болесника и преглед литературе

¹National Health Insurance Fund of Serbia, Belgrade, Serbia;

²University Clinical Centre of Serbia, Clinic for Digestive Surgery, Department for HBP Surgery, Belgrade, Serbia;

³University of Belgrade, Faculty of Medicine, Department for Surgery with Anesthesiology, Belgrade, Serbia;

⁴Bežanijska Kosa University Hospital Medical Center, Department for HBP Surgery, Belgrade, Serbia

Received: March 12, 2022

Accepted: May 18, 2022

Online First: May 25, 2022

DOI: <https://doi.org/10.2298/SARH220312051E>

*Accepted papers are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

*Correspondence to:

Nikola GRUBOR

Department for HBP Surgery, Clinic for Digestive Surgery, University Clinical Centre of Serbia, 6 Koste Todorovica Street, Belgrade 11000, Serbia

E mail: n.grubor@yahoo.com

Giant spleen as a surgical challenge – case report and literature review

Масивна спленомегалија као хируршки изазов – приказ болесника и преглед литературе

SUMMARY

Introduction According to the guidelines of the European Association of Endoscopic Surgery (EAES), any case where the maximum craniocaudal splenic diameter exceeds 20 cm is considered as massive splenomegaly. In addition to metabolic, hematological, and hemodynamic problems enlarged spleen may cause mechanical difficulties due to the pressure to surrounding organs and vascular structures. The aim of this paper is to present the surgical challenges and technique applied in massive splenomegaly, in a patient who had neglected the importance of regular medical checkups.

Case outline We present a 62-year-old male patient who was admitted to hospital for treatment of previously clinically and radiologically verified splenomegaly but who neglected the importance of regular checkups and medical treatment. Splenectomy was performed with splenic specimen 38 cm in its maximal diameter.

Conclusion Taking into consideration all the possible benefits and possible complications of surgical treatment, including the quality of life of splenectomized patients, comprehensive preoperative assessment should be made, and surgical treatment selectively applied.

Keywords: splenomegaly; massive splenomegaly; splenectomy

САЖЕТАК

Увод Према критеријумима Европског удружења за ендоскопску хирургију масивна спленомегалија подразумева случајеве у којима је максимални кранио-каудални дијаметар слезине превазилази 20 cm. Осим метаболичких, хематолошких и хемодинамских поремећаја, значајно увећана слезина може испољавати непожељан компресивни ефекат на суседне органе и васкуларне структуре. Циљ овог приказа случаја је да се прикажу изазови и техника хируршког лечења масивне спленомегалије.

Приказ болесника Болесник стар 62 године са раније дијагностикованом спленомегалијом који је занемарио значај редовних контролних прегледа, примљен је на клинику због тежоба узрокованих масивном спленомегалијом. Након пажљиво спроведене преоперативне припреме одстрањена је слезина кранио-каудалног дијаметра 38 cm.

Закључак Узимајући у обзир све могуће бенефите али и компликације хируршког лечења болесника са масивном спленомегалијом, неопходна је свеобухватна преоперативна хируршка процена и припрема, а хируршко лечење се планира у пажљиво изабраним случајевима.

Кључне речи: спленомегалија; масивна спленомегалија; спленектомија

INTRODUCTION

Massive splenomegaly is an enlargement of the spleen that can occur in a wide array of diseases and conditions. According to the guidelines of the European Association of Endoscopic Surgery (EAES), any case where the maximum craniocaudal splenic diameter exceeds 20 cm is considered as massive splenomegaly [1].

In addition to metabolic, hematological, and hemodynamic problems, an enlarged spleen may also cause mechanical difficulties. Due to the pressure to surrounding organs and vascular structures, patients often complain of a sense of heaviness in the abdomen, frequent perspiration, nausea, indigestion, etc. [2].

The diagnostics in splenomegaly, depending on the cause, entails a series of procedures, starting with a physical exam, and also including abdominal ultrasonography, computerized tomography (CT), magnetic resonance imaging (MRI), laboratory analyses, and finally splenectomy. A definitive diagnosis is established after histopathological examination [2, 3, 4].

Splenomegaly treatment may be nonsurgical, which includes the treatment of the basic cause of spleen enlargement, as well as surgical. Surgical treatment involves the removal of the spleen, necessitated by the difficulties that its size causes, and also by rupture or imminent risk of rupture. Also, when preoperative diagnostics does not offer reliable data, splenectomy is performed for diagnostic purposes [5].

The aim of our paper is to present the surgical technique applied in massive splenomegaly, in a patient who had neglected to observe the importance of regular checkups and medical treatment.

CASE REPORT

We present a 62-year-old male patient who was admitted to hospital for treatment of previously clinically and radiologically verified splenomegaly. Abdominal CT detected an enlarged spleen of enormous size - 39.3 cm (Figure 1). At admission, we learned from the patient that he had been diagnosed with an enlarged spleen a number of years before, that he had had bone marrow biopsy several times and that the results of the biopsies had been normal. We also found out that he had not been keeping up with regular check-ups and examinations during the previous several years. Medical records revealed that he had undergone urinary bladder surgery several years before, due to calculosis, as well as surgery of bilateral inguinal hernia, two years before.

Two months before admission to our hospital the patient visited his doctor complaining of abdominal pain, the sensation of heaviness, discomfort and increased perspiration. After complete radiological diagnostics had been performed, the patient was examined and tested by a hematologist. Immunophenotyping of peripheral blood was performed and it was determined that the patient had the B monoclonic population which was Pan B+, CD5-, CD10-, FMC7+

hetero, CD23+ hetero, CD38+ hetero, with IgM/kappa+, with a HILL score of 1. In other laboratory results there was a marked monoclonic IgM kappa paraprotein in low concentration.

Upon the completion of hematological diagnostics and preparation, further surgical treatment was decided on during a clinical case conference. After preoperative preparations were performed, the patient was placed under general anesthesia and the abdomen was opened with midline laparotomy. The abdomen revealed an enlarged spleen interspersed with white patches (Figure 2), which was pressing the surrounding organs and vascular structures to the opposite side of the abdomen. The next step was the mobilization of the spleen with the aid of the LigaSure device (SurgRx, Redwood City, CA), for severing splenic ligaments. We first identified the lienal artery and vein, which were then, through careful preparation, ligated and severed. Then, we dealt with the short gastric blood vessels, also with the use of LigaSure. After this, splenectomy was completed without incident and removed from the abdomen in its entirety. After hemostasis was established, an abdominal drain was placed in the left subphrenic space, and the abdomen was closed by anatomical layers.

The splenic tissue specimen taken after the removal of the spleen from the abdomen (Figure 3) was sent for definitive histopathological examination, which confirmed hairy cell leukemia.

The patient was discharged from hospital on the 6th postoperative day with prescribed antibiotic prophylaxis and a recommendation for post splenectomy immunization. Also, the patient was referred for further treatment and follow-up with a hematologist.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

DISCUSSION

Clinically massive splenomegaly represents a finding corresponding with a palpable mass in the abdomen below the left costal margin at a distance > 8 cm towards the superior iliac spine. The recommendation of the EAES is to use metric units for defining splenomegaly in preoperative imaging diagnostics. The golden standard in diagnosing and determining

massive splenomegaly is calculating the splenic volume, however this procedure requires more time than standard radiological methods [5, 6].

An enlarged spleen can occur within numerous infectious diseases, also, in hereditary diseases, immunological diseases, as well as in hematological diseases. The most frequent causes of splenomegaly are shown in Table 1 [7].

Pronounced symptoms are mostly present in massive splenic enlargement, which is why patients commonly complain of symptoms manifesting after food intake, of the inability to adequately empty their bladders, of obstructive complaints, as well as of the inability to flex their trunk [8, 9, 10].

In the patient we are presenting, the symptoms were the main reason for his visiting the doctor again. It is our belief, based on insight into the medical records, that bilateral inguinal hernia was directly caused by splenic enlargement and increased pressure within the abdomen.

Splenomegaly can generally be treated with medication, with reductive therapy (radiation therapy), although, in massive splenomegaly, potential complications related to this condition should always be taken into consideration, which is why, in this sense, surgical treatment should not be delayed. As a treatment modality in potential rupture, preoperative embolization of the splenic artery should be considered, when there are technical conditions for this procedure, since, in this way, the possibility of bleeding is reduced and partial reduction of the volume of the spleen is achieved, which further facilitates surgical treatment [11, 12, 13].

Since the diagnosis had been established preoperatively and since other treatment options had been exhausted, splenectomy was performed in our patient as a form of palliative treatment, i.e., debulking.

Surgical treatment is a second, sometimes a third therapeutic option, in treating splenomegaly. Taking into consideration all the possible benefits and possible complications of surgical treatment, including the quality of life of splenectomized patients, preoperative assessment should be made, and surgical treatment should be selectively applied, bearing in mind that massive splenomegaly is a great surgical challenge.

Conflict of interest: None declared.

REFERENCES

1. Habermalz B, Sauerland S, Decker G, Delaitre B, Gigot JF, Leandros E, et al. Laparoscopic splenectomy: the clinical practice guidelines of the European Association for Endoscopic Surgery (EAES). *Surg Endosc.* 2008 Apr;22(4):821-48. doi: 10.1007/s00464-007-9735-5. PMID: 18293036
2. Fieldman L, Munshi A, Al-Mahroos M, Fried G. The Spleen. *Maingot's abdominal operations.* 13th ed. New York: McGraw-Hill; 2019. p. 1239-49.
3. Aldulaimi S, Mendez AM. Splenomegaly: Diagnosis and Management in Adults. *Am Fam Physician.* 2021;104(3):271-276. PMID: 34523897.
4. Lee MW, Yeon SH, Ryu H, Song IC, Lee HJ, Yun HJ, et al. Volumetric Splenomegaly in Patients With Polycythemia Vera. *J Korean Med Sci.* 2022 Mar 21;37(11):e87. doi: 10.3346/jkms.2022.37.e87. PMID: 35315598;
5. Lemaire J, Rosière A, Bertrand C, Bihin B, Donckier JE, Michel LA. Surgery for massive splenomegaly. *BJS Open.* 2017 Apr 6;1(1):11-17. doi: 10.1002/bjs5.1. PMID: 29951600
6. Sjoberg BP, Menias CO, Lubner MG, Mellnick VM, Pickhardt PJ. Splenomegaly: A Combined Clinical and Radiologic Approach to the Differential Diagnosis. *Gastroenterol Clin North Am.* 2018 Sep;47(3):643-666. doi: 10.1016/j.gtc.2018.04.009. PMID: 30115442.
7. Chapman J, Bansal P, Goyal A, Azevedo AM. Splenomegaly. 2021 Aug 11. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2022. PMID: 28613657.
8. Tago M, Fujiwara M, Nakashima T, Makio S, Hirakawa Y, Tokushima M, et al. Massive splenomegaly requiring differential diagnosis of hematologic malignancies. *Clin Case Rep.* 2022;10(3):e05512. doi: 10.1002/ccr3.5512. PMID: 35310319;
9. Pavlovic-Popovic, Zora, Bojan Zaric, Zdravko Kosjerina, Dragana Petrovic. Splenomegaly in Sarcoidosis: Frequency, Treatment, Prognosis and Long-Term Follow-Up. *Srpski Arhiv Za Celokupno Lekarstvo.* 2015; 143 5–6): 279–83. <https://doi.org/10.2298/SARH1506279P>.
10. Ammad Ud Din M, Shapiro J, Sham R. Massive splenomegaly in hairy cell leukemia causing urinary retention. *Clin Case Rep.* 2020 Dec 4;9(2):1027-1028. doi: 10.1002/ccr3.3624. PMID: 33598291.
11. Hudson VE, Rogers PS, Bennett JMH. Management of massive splenomegaly. *Clin Case Rep.* 2021 Aug 21;9(8):e04603. doi: 10.1002/ccr3.4603. PMID: 34457283.
12. Markovic O, Divac A, Lukic F, Mrda D, Vidovic A, Zdravkovic M, et al. Successful Treatment of Spontaneous Rupture of the Spleen by Embolization of Splenic Artery in a Patient with Acute Promyelocytic Leukaemia and COVID-19 Infection. *Open Journal of Emergency Medicine.* 2021; 9, 60-65. doi: 10.4236/ojem.2021.93008.
13. Zaorsky NG, Williams GR, Barta SK, Esnaola NF, Kropf PL, Hayes SB, et al. Splenic irradiation for splenomegaly: A systematic review. *Cancer Treat Rev.* 2017 Feb;53:47-52. doi: 10.1016/j.ctrv.2016.11.016. PMID: 28063304




Figure 1. Abdominal computed tomography detected an enlarged spleen of enormous size, 39.5 in diameter




Figure 2. Intraoperative photo shows the enlarged spleen that occupied most of the abdominal cavity



Figure 3. Splenic specimen

Table 1. Causes of splenomegaly

Liver disease	cirrhosis, hepatitis
Hematologic malignancies	lymphomas, leukemias, myeloproliferative disorders
Venous thrombosis	portal or hepatic vein thrombosis
Splenic congestion	venous thrombosis, portal hypertension
Cytopenias	Immune thrombocytopenic purpura, autoimmune hemolytic anemia, immune-mediated neutropenia, Felty syndrome
Splenic sequestration	pediatric sickle cell disease, hemolytic anemias, thalassemias
Acute or chronic infection	bacterial endocarditis, infectious mononucleosis, HIV, malaria, tuberculosis, histiocytosis, abscess
Connective tissue diseases	systemic lupus erythematosus, rheumatoid arthritis, adult-onset Still's disease, and some familial autoinflammatory syndromes
Infiltrative disorders	sarcoidosis, amyloidosis, glycogen storage disease
Focal lesions	hemangiomas, abscess, cysts, metastasis