



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Ivan Palibrk^{1,2}, Marija Đukanović^{1,2}, Maja Maksimović-Mandić^{2,*}, Bojana Miljković²,
Dubravka Đorović², Jelena Veličković^{1,2}

**Analysis of comorbidity and anesthesia technique in patients undergoing
bariatric surgery at the University Clinical Center of Serbia**

Анализа коморбидитета и технике анестезије код пацијената на
баријатријској хирургији у Универзитетском клиничком центру Србије

¹University of Belgrade, Faculty of Medicine, Belgrade, Serbia;

²University Clinical Center of Serbia Center for Anesthesiology and Reanimatology, Belgrade, Serbia

Received: April 16, 2024

Revised: June 10, 2024

Accepted: July 1, 2024

Online First: July 3, 2024

DOI: <https://doi.org/10.2298/SARH240416055P>

***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

***Correspondence to:**

Maja MAKSIMOVIĆ-MANDIĆ

Druge Srpske Armije 17/12, 11000 Belgrade, Serbia

E-mail maja_maksimovic14@hotmail.com

Analysis of comorbidity and anesthesia technique in patients undergoing bariatric surgery at the University Clinical Center of Serbia

Анализа коморбидитета и технике анестезије код пацијената на баријатријској хирургији у Универзитетском клиничком центру Србије

SUMMARY

Introduction/Objective Altered physiology and metabolism of obese patients represents a big challenge for the anesthesiologist.

The objectives of the study are to investigate numbers of comorbidities, choice of anesthesia techniques, intraoperative and postoperative complications between bariatric and non-bariatric patients.

Methods This retrospective study included 469 patients. The study group of patients included obese patients with Body Mass Index ≥ 30 . The control group included patients in whom elective laparoscopic cholecystectomy was performed, on the same day as bariatric surgery in the control group.

Results The study group included 235 patients who underwent bariatric surgery, while control group included 234 patients. More patients in study group had comorbidities compared with the control group (84.4% vs. 63.2%, $p < 0.001$). In the study group, total intravenous anesthesia and target control anesthesia were statistically significant more delivered than in the control group (74% vs. 0.9%, $p < 0.001$; 7.2% vs. 1.7, $p < 0.001$, respectively). Difficult intubation was statistically significant more in the control group (5.6% vs. 0.9%, $p = 0.004$).

There was a statistically significant difference in the incidence of intraoperative desaturation and hypotension during induction of anesthesia between the study and the control group (9.8% vs. 2.1%, $p < 0.001$; 14.5% vs. 2.1, $p < 0.001$, respectively). There was statistically significant difference between the study and control group in minor complication according Clavian-Dindo classification, (20.8% vs. 5.1%, $p < 0.001$).

Conclusion Obesity is associated with higher number of comorbidities and intraoperative complications. There was no statistically difference in major postoperative complications between bariatric and non-bariatric patients.

Keywords: obesity; bariatric surgery; comorbidities; body mass index

САЖЕТАК

Увод/Циљ Измењена физиологија и метаболизам гојазних пацијената представља изазов за анестезиолога. Циљеви овога рада су испитивање учесталости коморбидитета, типа анестезије, учесталост интраоперативних и постоперативних компликација код пацијената који су подвргнути баријатријској хирургији и пацијената којима је учињена елективна холецистектомија.

Метод Ова ретроспективна студија је обухватила 469 пацијената. У испитиваној групи су укључени гојазни пацијенти са индексом телесне масе ≥ 30 . Контролну групу су чинили пацијенти за елективну лапароскопску холецистектомију оперисани истог дана када и баријатријски.

Резултати Испитивана група је укључила 235 пацијената, док је контролна група обухватила 234 пацијента. Учесталост коморбидитета била је статистички значајно већа у испитиваној у односу на контролну групу (84,6% и 63,2%, $p < 0,001$). Постојала је статистички значајна разлика у анестезиолошкој техници – тотална интравенска анестезија и анестезија циљано контролисаном инфузијом су више примењиване у испитиваној групи (74% наспрам 0,9%, $p < 0,001$; 7,2% наспрам 1,7, $p < 0,001$). Број отежаних интубација је био статистички значајно већи у контролној групи (5,6% наспрам 0,9%, $p = 0,004$). Постојала је статистички значајна разлика у инциденци десатурације и хипотензије током увода у анестезију – ове компликација забележене су чешће у испитиваној у односу на контролну (9,8% наспрам 2,1%, $p < 0,001$; 14,5% наспрам 2,1 $p < 0,001$). Статистички значајна разлика је постојала у инциденци малих компликација између испитиване и контролне групе према Клавијен–Диндо класификацији (20,8% наспрам 5,1%, $p < 0,001$).

Закључак Гојазност је повезана са бројним коморбидитетима и вишом инциденцом интраоперативних компликација. Упркос томе, не постоји статистички значајна разлика у великим постоперативним компликацијама између ове две групе оперисаних болесника.

Кључне речи: гојазност; баријатријска хирургија; коморбидитети; индекс телесне масе

INTRODUCTION

According to the definition of the World Health Organization (WHO), obesity represents “abnormal or excessive fat accumulation that presents a health risk” [1]. Obesity is defined by a body mass index (BMI). BMI between 25 and 29.9 kg/m^2 defines overweight, while BMI over 30 kg/m^2 considers obesity [1, 2]. The obesity or overweight affects about 60% of the adult population. Also, in one of three children obesity is detected. Overall, obesity has been identified as the fourth-leading cause of noncommunicable diseases (NCDs) [1]. In 2019, 20.8% of the population over the age of fifteen was obese in Serbia [3]. Comorbidities such as cardiovascular disease, type 2 diabetes, rheumatoid arthritis, major depressive illness, polycystic ovarian syndrome, asthma, and obstructive sleep apnea are more likely in obese patients. [2, 4, 5, 6].

The metabolic, anatomical, and physiological aspects of obese patients make induction and maintenance of anesthesia challenging [7, 8]. Obese patients often have upper airway obstruction, decreased lung capacities and compliance, higher respiratory exertion, and impaired gas exchange. Respiratory pathophysiology is altered [7, 8]. Difficult ventilation and intubation are expected during anesthesia induction for bariatric surgery [8, 9]. Determine the dose of the anesthetic drugs in obese patients may be particularly challenging. Lipophilic drugs such as propofol, barbiturates, and benzodiazepines characterize high volume of distribution (V_d). To achieve adequate serum concentrations, larger loading doses are needed, therefore doses are calculated based on total body weight (TBW). For maintenance of anesthesia, dosing of these medications should be calculated based on the ideal body weight (IBW) or lean body weight (LBW). Loading succinylcholine dose is calculated based on TBW. Nondepolarizing muscle relaxants dose is calculated based on IBW as in non-obese patients. Fentanyl, sufentanil, alfentanil should be estimated based on LBW, whereas remifentanil on IBW [8, 10].

The obese patients are at greater risk for developing postoperative complications. Wound infection, intra-abdominal infection, bowel injury, myocardial and other major complications are more common in obese patients [11]. An adequate preoperative assessment is mandatory as optimal intraoperative anesthetic management and postoperative care to prevent postoperative complications after bariatric surgery [7, 8, 12].

We hypothesized that obese patients have higher prevalence of comorbidities, and are more prone to postoperative complications.

The objectives of this study are to examine the prevalence of comorbidities in patients undergoing bariatric surgery compared to patients undergoing elective cholecystectomy. Also, we analyzed the choices of the anesthesia techniques in bariatric and elective surgery, the incidence of intraoperative and postoperative complications between bariatric and non-bariatric patients.

METHODS

This retrospective study included 469 patients, scheduled for bariatric surgery or elective cholecystectomy at the Hospital for digestive surgery, University Clinical Center of Serbia, in the period from June 2011 to November 2022. This study was approved by the ethics committee of University Clinical Center of Serbia, protocol number (661/2).

Obese patients with a BMI ≥ 30 undergoing bariatric surgery were included in the study group (SG). The control group (CG) included patients admitted to the hospital for an elective laparoscopic cholecystectomy procedure and scheduled for surgery on the same day as the study group. The preoperative interdisciplinary team specialists were decided about eligibility of the patients for bariatric surgery. This interdisciplinary health care team included an anesthesiologist, surgeon, pulmonologist (a spirometry report was mandatory), cardiologist (an ergospirometry was mandatory), psychiatrist, endocrinologist, and for women, a gynecologist.

A cardiologists examination with electrocardiogram (ECG) and chest X-ray not older than one month were obligatory before cholecystectomy in the control group. Laboratory panel testing (complete blood count, biochemical and coagulation analyses) within 14 days before surgery was mandatory in both groups. All patients from both groups received antibiotic prophylaxis (cephazoline) 30 minutes before the operation. Low molecular weight heparin (LMWH) for deep vein thrombosis prophylaxis was administered 2 hours before surgery. In the present study, hypertension, cardiac arrhythmias, coronary heart disease (CHD), hyperlipidemia (HLP), diabetes mellitus (DM) type 1 or 2, chronic obstructive pulmonary disease (COPD), obstructive hypoventilation syndrome or obstructive sleep apnea (OSA), epilepsy, anemia, varicose veins of lower extremities were recorded. The other comorbidities were listed as additional comorbidities. The American Society of Anesthesiologists (ASA) status was used to rated patients conditions.

Difficult intubation was recorded according to definition from the latest the Difficult Airway Society guidelines [13]. All analyzed data were obtained from medical records of patients.

Premedication with benzodiazepines was not routinely used. Before induction of anesthesia, preoxygenation was performed in all patients. Anesthesia techniques - balanced anesthesia (BA), total intravenous anesthesia (TIVA) or target control infusion anesthesia (TCI) was chosen by attending anesthesiologist. Induction of BA was achieved with propofol 1.5-2 mg/kg according to TBW and fentanyl 2-4 mcg/kg LBW. For maintenance of anesthesia sevofluran was used, with minimal alveolar concentration (MAC) 0.8-1.2Vol% based on the patient's age. Analgesia was achieved with fentanyl 50-200 mcg/h as an intermittent intravenous bolus according to patient comfort. For total intravenous anesthesia (TIVA) propofol was used 10-16 mg/kg/h according to TBW and remifentanyl 0.25-1 mcg/kg/min according to IBW for induction. During maintenance of anesthesia, propofol was used 4-6

mg/kg/h of LBW and remifentanyl 0.2-0.5 *mcg/kg/min* of IBW. In case of using target control infusion anesthesia (TCI), Marsh or Schneider model was used with target concentrations of propofol 6-8 *mcg/ml* according to TBW or LBW, respectively for induction of anesthesia. Maintenance anesthesia doses were 2-4*mcg/ml* of LBW. Remifentanyl (target effect site) was used in the range 6-10 *ng/ml* according to IBW for analgesia. In both intravenous techniques the breathing mixture was a combination of oxygen and air. During induction, for laryngoscopy and intubation, succinylcholine was used in dose of 1-1.2 *mg/kg* according to TBW, or rocuronium in dose of 0.6-1.2 *mg/kg* according to IBW in all patients. For maintenance neuromuscular blockade rocuronium was used in all anesthesia techniques in dose of 0.3 *mg/kg* IBW. Reversal of the neuromuscular blockade was performed with prostigmine/atropine or sugammadex in all patients, depending on attending the anesthesiologist.

Intraoperative monitoring (heart rate, electrocardiogram, non-invasive blood pressure, peripheral saturation of oxygen (SpO₂) and end-tidal CO₂ concentration was performed in all patients from both groups. In patients in the study group, two peripheral venous lines were placed, while in the control group one peripheral venous line was placed. Urinary catheter was inserted and hourly urine output was monitoring in the study group. Bispectral Index™ (BIS™ Medtronic, Minneapolis, USA) was used for TIVA or TCI.

In all patients, trachea was extubated in the end of surgery. After the extubation, the majority of patients were transferred to the department after 1 hour staying in the recovery room. Patients who required non-invasive mechanical ventilation following surgery were admitted to the intensive care unit (ICU) and stayed overnight. Postoperative multimodal analgesia was achieved with nonsteroid anti-inflammatory drugs, paracetamol and metamizole. If patient needed additional analgesia, tramadol or morphine were administered intravenously. Intraoperative and postoperative complications was recorded in patient's medical records. Bronchospasm, pneumothorax, desaturation (defined as SpO₂ < 90%), hypotension (defined as

systolic pressure $< 90\text{mmHg}$), hypertension (defined as $> 20\%$ of initial arterial pressure), bradycardia (defined as $\text{HR} < 50$ per minute), tachycardia (defined as $\text{HR} > 100$ per minute), and cardiac arrhythmia were defined as intraoperative complications and were reported in anesthesia records.

Postoperative complications were registered and categorized according to the Clavien-Dindo (CD) classification of surgical complications [14]. Minor complications were defined as CD grade I and II, major complications were defined as CD grade III and IV.

Statistical analysis was performed in IBM SPSS, version 28.0.1.1. Data were collected from medical and anesthesia records of patients. Data were described and analyzed using descriptive statistics, mean and standard deviation for continuous variables and counts and percentages for categorical variables. For clinical outcomes, for categorical variables Chi-square test or Fisher's exact test was used. For parametric variables, Student t-test was used. For non-parametric test Mann-Whitney test was performed. Statistical significance was calculated at level of significance of $p < 0.05$.

RESULTS

Of the total number of patients (469), the study group included 235 of patients, while in the control group were 234 patients. There was a statistically significant difference in the BMI between study and control group (44.9 ± 6.2 vs. 27.5 ± 4.6 , $p < 0.001$) (Table 1). In the control group, 26% of patients had $\text{BMI} > 30 \text{ kg/m}^2$. There was a statically significant difference in the age, younger patients were in study group (40.75 ± 9.9 vs. 48 ± 13.6 , $p < 0.001$) (Table 1). There was a statistically significant difference in the ASA status between groups ($p < 0.001$), patients in the study group were rated with higher ASA status (Table 1). More comorbidities were reported in the study than in control group (84.6% vs. 63.2%, $p < 0.001$) (Table 2). There was statistically significant difference in prevalence of HTA, DM and COPD in the study group

(55.8% vs. 39.3%, $p < 0.001$; 58% vs. 12.8%, $p < 0.001$; and 19% vs. 7.7%, $p < 0.001$, respectively (Table 2). More cardiac arrhythmia was detected in the control group (2.6 % vs. 6.8%, $p = 0.047$), and additional comorbidity was more verified in the study group (38.1% vs. 25.6%, $p = 0.005$) (Table 2). Premedication was more delivered in the study compared to the control group (70.6% vs. 33.9%, $p < 0.001$) (Table 3.). There was statistically significant difference in using succinylcholine for intubation between the study and the control group (87.5% vs 71.7%, $p < 0.001$) (Table 3). Also, there was statistically significant difference in using TIVA and TCI between study and control group (74% vs. 1.1%, $p < 0.001$; 7.2% vs. 1.7%, $p = 0.004$) (Table 3.). BA was the technique of choice in control group, and was statistically more performed (97.7% vs. 11.9, $p < 0.001$) (Table 3). The reversion of neuromuscular blockade was used in both groups, statistically significantly more often used in the study group difference (99.1% vs. 96.2%, $p = 0.032$) (Table 3.).

For neuromuscular reversion, sugamadex was used more in the study group (82.8% vs. 0.4%, $p < 0.001$) (Table 3.). There was no statistically significant difference in the occurrence in the total number of intraoperative complications between study and control group (42.6% vs. 43.2%, $p = 0.894$) (Table 4). Difficult intubation was more documented in the control group in compare to the study group (0.9% vs. 5.6%, $p = 0.004$) (Table 4). Incidence of intraoperative desaturation occurred significantly more in the study than in the control group (9.8% vs. 2.1%, $p < 0.001$) (Table 4). Hypotension episodes was statistically significant more documented in study compared to control group (14.5% vs. 2.1%, $p < 0.001$) (Table 4). There was statistical significance in occurrence of tachycardia between study and control group, more tachycardia was registered in control group (8.1% vs. 16.7%, $p = 0.005$) (Table 4). Postoperative complications according Clavien-Dindo classification gradus I was significantly more documented in the study compared to the control group (18.3% vs. 4.7%, $p < 0.001$). There was

statistically significant difference in occurrence of minor postoperative complications (CD grade I and II) between the study and control group (Table 5).

DISCUSSION

Obesity is a chronic disease of modern ages. Obesity itself already represents sufficient severity and these patients often have two or more comorbidities. [2,5,6,]. In the present study, more comorbidities were reported in obese patients than in patients for elective cholecystectomy. TIVA and TCI were the most common choice in bariatric patients. There were statistically significant more minor postoperative complications in the study group.

In the study group, an average BMI was 44.9 kg/m^2 , the mean age was 40 and 68.9% of patients were female which is in agreement with results from The International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) and in the research of the North-Western Europe countries. IFSO and North-Western Europe countries reported that bariatric patients have an average BMI $40\text{-}45 \text{ kg/m}^2$, the ages of forties with highest percentage of female patients [11, 15].

According to the results of the National Institute for Public Health in Serbia, 20.8% of the general population is obese [3]. In the control group, 26% patients were obese. It seems that a significant number of obese patients are going on the elective surgery on a daily basis.

Comorbidity frequency in bariatric patients varies significantly among countries, according to the population studies - hypertension (up to 83.2 %), HLP (up to 82.1%), DMT2 (up to 47.4%), and musculoskeletal pain (43.7%) [11,15,16,17]. In our study, bariatric patients were suffered from hypertension (55.8%), DMT2 (58%), and HLP (12.6%). The IFSO reported a large variation of the OSA incidence from 49.5% in Canada - Ontario to the lowest rates in Russia (2.7%), and 40% in UK [15,18]. In the current study OSA was found in 4.3% patients. The reason for a large disparity in the OSA incidence between observed centers may be found

that experts conducting polysomnography studies are needed [15]. COPD was documented in 19% of patients in the study group with a statistical difference compared to control group. Verberne et al. reported that one third of patients presenting with COPD have an average BMI of 33.7 kg/m^2 [6]. There was a statistically significant difference between the ASA status in study and control group. The majority of patients were rated with ASA score ≥ 2 in the study group, while in the control group the most of patients were rated with ASA status 1. According to literature, regardless whether obese patients have comorbidities, they will be rated with a higher ASA status. Patients with a BMI between 30 and 40 will be rated with ASA status 2, and for BMI over 40 with ASA status 3 [19].

The choice of anesthesia technique in obese patients depends of the excessive volume of distribution. These patients are often under- or over-dosed with anesthetic drugs [10,20].

TIVA and TCI with current pharmacokinetic models represent safe and precise anesthetic techniques, but definitely necessary combustible dose titration in obese patients. The use of BIS monitoring is mandatory, but clinical effects are also important [20,21]. In the present study TIVA and TCI with mandatory BIS monitoring were statistically significant more delivered during bariatric surgery compared to cholecystectomy where BA was used more frequently. Research show that opioid-free anesthesia in bariatric surgery is also a safe technique [22].

A difficult intubation is expected in bariatric patients. De Jong et al. showed that succinylcholine was the most common choice for muscle relaxation for intubation in the ICU (in 70% of cases), while in the operating room succinylcholine was used in only 19% of obese patients [23]. For intubation, atracurium and cisatracurium were the main choice in 73% of patients, whereas rocuronium was used in only 1% of patients in the operating room, and 11% of patients in the ICU [23]. The frequency of difficult intubations in obese patients was 8.2% in the operating theater and 16.3% in ICU [23]. In our study, difficult intubation was observed

only in 0.9% of patients in the study group, in contrast to 5.6% patients in the control group. This may be explained by the fact that we expected a difficult intubation in the study group, and the anesthesiologist was prepared for it. Every patient in the study group was positioned according to the recommendations (Rapid Airway Management Positioner- „RAMP-up “).

During the induction of anesthesia, the main complication was desaturation - 9.8% in the study group compared to 2.1% in the control group. De Jong et al. reported severe hypoxemia in ICU obese patients (50%), while in the operating theatre no severe hypoxemia was occurred [23,24]. In our study, all desaturations during apnea time were lasting less than 90 seconds in both groups and did not affect patient's safety. Reduced oxygen reserve due to lung restriction is the reason for desaturation during apnea time. An adequate patient positioning and nasopharyngeal insufflation of oxygen during the apnea period is sufficient to prevent desaturation in almost 100% of morbid obese patients [25]. In the present study, only in morbid obese patients with BMI>55 nasopharyngeal insufflation of oxygen was used during the apnea period.

The literature data favor the reversion of the complete neuromuscular blockade. [7, 12, 26]. Gaszynski et al. [26] showed benefit of using sugammadex, the train-of-four ratio was 3.5 times faster, than in the group receiving neostigmine for decurarization [26]. In our study, 82.8% of patients in the study group received sugammadex in compared 0.4% of patients in the control group for faster and safer reversal of neuromuscular block.

According to a multinational study of North -Western European countries, complications after bariatric intervention occurred in 6.5% of patients. The most common were bleeding, anastomotic leakage, gastrointestinal perforations and postoperative ileus [11]. In our research, the most common complications were CD grade 1 and 2 in study group. There was a statistically significant difference in minor complications between groups, but with no

significant difference in major complications. More minor complications were documented in the study group.

Limitations

The limitation of this study is that only elective laparoscopic gallbladders were observed in the control group. The reason is that it is the most common elective laparoscopic surgery, and patients are discharged home on the first or second postoperative day.

CONCLUSION

This study showed 1/3 of the patients in the elective program are obese. Bariatric patients are younger with more comorbidities compared to non-bariatric patients. In order to increase the safety of anesthesia in bariatric patients a multidisciplinary approach is required. TIVA and TCI are safe anesthesia technique in bariatric surgery. This study showed that bariatric patients have the same incidence of major postoperative complications as patients after elective cholecystectomy. Further research is needed to determine the clinical significance of our findings, in particular in the safety of the anesthesia technique and incidence of perioperative complications.

First-co-authors Marija Đukanović and Maja Maksimović-Mandić have contributed equally to this work and share first authorship.

Conflict of interest: None declared.

REFERENCES

1. WHO European Regional Obesity Report 2022. Copenhagen: WHO Regional Office for Europe; 2022. Licence: CC BY-NC-SA 3.0 IGO. (updated 2.May 2022; Cited 10.mart 2024). Available from: WHO European Regional Obesity Report 2022. ISBN: 9789289057738
2. Mahmoud I, Sulaiman N. Significance and agreement between obesity anthropometric measurements and indices in adults: a population-based study from the United Arab Emirates. *BMC Public Health*. 2021;21(1):1605. DOI: 10.1186/s12889-021-11650-7. PMID: 34465314;
3. Milić N, Stanisavljević D, Krstić M, Jovanović V, Brcanski J, Kilibarda B, et al. THE 2019 SERBIAN NATIONAL HEALTH SURVEY. Belgrade: Statistical Office of the Republic of Serbia; 2021.142p.
4. Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health*. 2009;25(9):88. DOI: 10.1186/1471-2458-9-88. PMID: 19320986.
5. Maksimović M, Vlajinac H, Radak D, Marinković J, Maksimović J, Jorga J. Relationship between Abdominal Obesity and Other Cardiovascular Risk Factors: Cross Sectional Study of Patients with Symptomatic Carotid Disease.*Srp Arh Celok Lek*. 2013 Jul Aug;141(7-8):460-465. DOI: 10.2298/sarh1308460m. PMID: 24073551.
6. Verberne LDM, Leemrijse CJ, Swinkels ICS, van Dijk CE, de Bakker DH, Nielen MMJ. Overweight in patients with chronic obstructive pulmonary disease needs more attention: a cross-sectional study in general practice. *NPJ Prim Care Respir Med*. 2017;22,(1):63. DOI: 10.1038/s41533-017-0065-3. PMID: 29167434.
7. Seyni-Boureima R, Zhang Z, Antoine MMLK, Antoine-Frank CD. A review on the anesthetic management of obese patients undergoing surgery. *BMC Anesthesiol*. 2022; 22(1):98. DOI: 10.1186/s12871-022-01579-8. PMID: 35382771.
8. Wynn-Hebden A, Bouch DC. Anaesthesia for the obese patient. *BJA Educ*. 2020;20(11):388-395. DOI: 10.1016/j.bjae.2020.07.003. PMID: 33456923.
9. Lee S, Jang EA, HongM, Bae HB, Kim J.Ramped versus sniffing position in the videolaryngoscopy-guided tracheal intubation of morbidly obese patients: a prospective randomized study. *Korean J Anesthesiol* 2023;76(1):47-55. DOI:10.4097/kja.22268 PMID:35912427
10. Ingrande J, Lemmens HJ. Dose adjustment of anaesthetics in the morbidly obese. *Br J Anaesth*. 2010;105(1):i16-23. DOI: 10.1093/bja/aeq312. PMID: 21148651.
11. Poелеmeijer, Y.Q.M., Liem, R.S.L., Våge, V. et al. Perioperative Outcomes of Primary Bariatric Surgery in North-Western Europe: a Pooled Multinational Registry Analysis.*OBES SURG*. 2018;28: 3916–3922. DOI: 10.1007/s11695-018-3408-4. PMID: 30027332.
12. Stenberg E, Dos Reis Falcão LF, O’Kane M, Liem R, Pournaras DJ, Salminen P, et al. Guidelines for Perioperative Care in Bariatric Surgery: Enhanced Recovery After Surgery (ERAS) Society Recommendations: A 2021 Update. *World J Surg*. 2022;46(4):729-751. DOI: 10.1007/s00268-021-06394-9. PMID: 34984504.
13. Apfelbaum JL, Hagberg CA, Connis RT, Abdelmalak BB, Agarkar M, Dutton RP, et al. 2022 American Society of Anesthesiologists Practice Guidelines for Management of the Difficult Airway. *Anesthesiology*. 2022 Jan 1;136(1):31-81. DOI: 10.1097/ALN.0000000000004002. PMID: 34762729.
14. Domenghino A, Walbert C, Birrer DL, Puhan MA, Clavien PA & The Outcome4Medicine consensus group. Consensus recommendations on how to assess the quality of surgical interventions.*Nat Med*.2023 Apr;29(4):811-822. DOI: 10.1038/s41591-023-02237-3.PMID: 37069361.
15. The International Federation for the Surgery of Obesity and Metabolic Disorders. 8th IFSO Global Registry Report 2023. IFSO 2023. (Cited 6 april 2024).Available from: <https://www.ifso.com/pdf/8th-ifso-registry-report-2023.pdf>
16. Pedersen MH, Bøgelund M, Dirksen C, Johansen P, Jørgensen NB, Madsbad S, et al. The prevalence of comorbidities in Danish patients with obesity - A Danish register-based study based on data from 2002 to 2018. *Clin Obes*. 2022;12(5):e12542. DOI: 10.1111/cob.12542. PMID: 35768944.
17. Castanha CR, Tcbc-Pe ÁABF, Castanha AR, Belo GQMB, Lacerda RMR, Vilar L. Evaluation of quality of life, weight loss and comorbidities of patients undergoing bariatric surgery. *Rev Col Bras Cir*. 2018;45(3):e1864. DOI: 10.1590/0100-6991e-20181864. PMID: 30020323.
18. The National Bariatric surgery registry of British Obesity & Metabolic Surgery Society. Third Registry Report. 2020. (Cited 10.01.2024) Available from: [NBSR2020.pdf](https://www.nbsr2020.pdf) (e-dendrite.com)
19. Guerry, C., Butterworth, J.F. A 25-year retrospective analysis of the American Society of Anesthesiologists physical status classification: did we “up-code” young obese patients when obesity was not yet considered a disease?. *Can J Anesth/J Can Anesth* .2018;65: 776–785. DOI: 10.1007/s12630-018-1096-0. PMID: 29572720
20. Coetzee JF. Total intravenous anaesthesia to obese patients: largely guesswork? *Eur J Anaesthesiol*. 2009 May;26(5):359-61. DOI: 10.1097/EJA.0b013e328329c6e2. PMID: 19359991.

21. Nimmo AF, Absalom AR, Bagshaw O, Biswas A, Cook TM, Costello A, et al. Guidelines for the safe practice of total intravenous anaesthesia (TIVA): Joint Guidelines from the Association of Anaesthetists and the Society for Intravenous Anaesthesia. *Anaesthesia*. 2019 Feb;74(2):211-224. DOI: 10.1111/anae.14428. PMID: 30378102.
22. Clanet M, Touihri K, El Haddad C, Goldsztejn N, Himpens J, Fils JF, et al. Effect of opioid-free versus opioid-based strategies during multimodal anaesthesia on postoperative morphine consumption after bariatric surgery: a randomised double-blind clinical trial. *BJA Open*. 2024, 23;9:100263. DOI: 10.1016/j.bjao.2024.100263. PMID: 38435809.
23. De Jong A, Molinari N, Pouzeratte Y, Verzilli D, Chanques G, Jung B, et al. Difficult intubation in obese patients: incidence, risk factors, and complications in the operating theatre and in intensive care units. *Br J Anaesth*. 2015;114(2):297-306. DOI: 10.1093/bja/aeu373. PMID: 25431308.
24. Yan L, Wang X, Du K, Liang Y. Effect of inspiratory muscle training on hypoxemia in obese patients undergoing painless gastroscopy: protocol for a single-center, double-blind, randomized controlled trial. *Front. Med (Laussane)*. 2023;10:1269486. DOI: 10.3389/fmed.2023.1269486. PMID: 37790126.
25. Baraka AS, Taha SK, Siddik-Sayyid SM, Kanazi GE, El-Khatib MF, Dagher CM, et al. Supplementation of pre-oxygenation in morbidly obese patients using nasopharyngeal oxygen insufflation. *Anaesthesia*. 2007; 62(8):769-73. DOI: 10.1111/j.1365-2044.2007.05104.x. PMID: 17635423.
26. Gaszynski T, Szewczyk T, Gaszynski W. Randomized comparison of sugammadex and neostigmine for reversal of rocuronium-induced muscle relaxation in morbidly obese undergoing general anaesthesia. *Br J Anaesth*. 2012;1(2):236-9. DOI: 10.1093/bja/aer330. PMID: 22012861.

Table 1. Demographic characteristics of patients

Parameters	Study group (n = 235)	Control group (n = 234)	p-value
Sex, female, n (%)	162 (68.9)	145 (62.2)	0.127**
Ages, mean \pm SD	40.75 \pm 9.9	48 \pm 13.6	< 0.001*
BMI (kg/m ²), mean \pm SD	44.9 \pm 6.2	27.5 \pm 4.6	< 0.001*
Body weight (kg), mean \pm SD	138 \pm 6.7	83.04 \pm 1.1	< 0.001*
ASA status			
ASA 1, n (%)	2 (0.9)	72 (30.8)	< 0.001**
ASA 2, n (%)	177 (76)	148 (63.2)	
ASA 3, n (%)	53 (22.7)	14 (6)	
ASA 4, n (%)	1 (0.4)	0 (0)	

*Student t test, ** Pearson's χ^2 test, p < 0.05 statistically significant difference

BMI – body mass index; ASA status – American Society of Anesthesiologists status

Table 2. Comorbidities

Parameters	Study group (n = 235)	Control group (n = 234)	p-value
Overall comorbidities, n (%)	198 (84.6)	148 (63.2)	< 0.001*
Hypertension, n (%)	130 (55.8)	92 (39.3)	< 0.001*
DM (type 1 or 2), n (%)	134 (58)	30 (12.8)	< 0.001*
Cardiac arrhythmia, n (%)	6 (2.6)	16 (6.8)	0.047*
HLP, n (%)	27 (12.6)	20 (8.5)	0.160*
CHD, n (%)	5 (2.2)	8 (3.4)	0.408*
Epilepsy, n (%)	3 (1.3)	3 (1.3)	0.992**
COPD, n (%)	44 (19)	18 (7.7)	< 0.001*
Anemia, no (%)	5 (2.2)	5 (2.1)	0.989*
Varicose veins of the lower extremities, n (%)	13 (5.6)	12 (5.1)	0.840*
OSA, n (%)	10 (4.3)	3 (1.3)	0.053**
Additional comorbidity, n (%)	88 (38.1)	60 (25.6)	0.005*

*Pearson's χ^2 test, **Fisher's exact test, p < 0.05 statistically significant difference

DM – diabetes mellitus; HLP – hyperlipoproteinemia; CHD – chronic heart disease; COPD – chronic obstructive pulmonary disease; OSA – obstructive sleep apnea

Table 3. Anesthesia techniques

Anesthesia techniques	Study group n = 235	Control group n = 234	p-value
Premedication, n (%)	166 (70.6)	79 (33.9)	< 0.001*
Neuromuscular relaxant for intubation			
Succinylcholine, n (%)	203 (87.5)	167 (71.7)	< 0.001*
Rocuronium, n (%)	27 (11.6)	62 (26.6)	
Cisatracurium, n (%)	2 (0.9)	4 (1.7)	
TIVA, n (%)	174 (74)	2 (1.1)	< 0.001**
TCI, n (%)	17 (7.2)	4 (1.7)	0.004**
BA, n (%)	28 (11.9)	229 (97.7)	< 0.001*
Reversion neuromuscular blockade, n (%)	233 (99.1)	225 (96.2)	0.032*
Neostigmine, n (%)	40 (17.2)	221 (98.7)	< 0.001*
Sugammadex, n (%)	193 (82.8)	1 (0.4)	
Neostigmine and sugammadex, n (%)	0 (0)	2 (0.9)	

*Pearson's χ^2 test, ** Fisher's exact test, $p < 0.05$ statistically significant difference
 TIVA – total intravenous anesthesia; TCI – target-controlled infusion; BA – balance anesthesia

Table 4. Intraoperative complications

Parameters	Study group n = 235	Control group n = 234	p-value
Total complications, n (%)	100 (42.6)	101 (43.2)	0.894*
Difficult intubation, n (%)	2 (0.9)	13 (5.6)	0.004**
Bronchospasm, n (%)	3 (1.3)	3 (1.3)	0.999**
Pneumothorax, n (%)	2 (0.9)	0 (0)	0.49**
Desaturation, n (%)	23 (9.8)	5(2.1)	< 0.001*
Hypotension, n (%)	34 (14.5)	5 (2.1)	< 0.001*
Hypertension, n (%)	49 (20)	67 (28.6)	0.055*
Bradycardia, n (%)	20 (8.5)	19 (8.1)	0.999*
Tachycardia, n (%)	19 (8.1)	39 (16.7)	0.005*
Cardiac arrhythmia, n (%)	0 (0)	2 (0.9)	0.248 **

*Pearson's χ^2 test, ** Fisher's exact test, $p < 0.05$ statistically significant difference

Table 5. Postoperative complications according Clavien-Dindo classification

Clavien- Dindo classification	Study group n = 235	Control group n = 234	p-value
Grade I, n (%)	43 (18.3)	11 (4.7)	< 0.001*
Grade II, n (%)	6 (2.6)	1 (0.4)	0.13**
Grade IIIa, n (%)	0 (0)	0 (0)	0.999**
Grade IIIb, n (%)	1 (0.4)	0 (0)	0.988**
Grade IV (a and b), n (%)	0 (0)	0 (0)	0.999**
Grade V, n (%)	0 (0)	0 (0)	0.999**
Minor complications, n (%) #	49 (20.8)	12 (5.1)	< 0.001*
Major complications, n (%) #	1 (0.4)	0 (0)	0.999**

*Pearson's χ^2 test, ** Fisher's exact test, p < 0.05 statistically significant difference

#Minor complications – Clavien–Dindo grade I and II; #Major complications – Clavien–Dindo grade III, IV and V