



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Case Report / Приказ болесника

Dejana Bajić^{1,*}, Zoran Komazec^{2,3}

Congenital tongue base cyst as uncommon cause of laryngeal stridor in an infant

Урођена циста базе језика као неуобичајен узрок ларингеалног стридора одојчета

¹University of Novi Sad, Faculty of Medicine, Department of Biochemistry, Novi Sad, Serbia;

²Clinical Center of Vojvodina, Clinic for Otorhinolaryngology and Head and Neck Surgery, Novi Sad, Serbia;

³University of Novi Sad, Faculty of Medicine, Department of Otorhinolaryngology, Novi Sad, Serbia

Received: January 11, 2024

Revised: July 11, 2024

Accepted: July 12, 2024

Online First: July 15, 2024

DOI: <https://doi.org/10.2298/SARH240111059B>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

* **Correspondence to:**

Dejana BAJIĆ

University of Novi Sad, Faculty of Medicine, Department of Biochemistry, Hajduk Veljkova 3, 21137 Novi Sad, Serbia

Email: dejana.bajic@mf.uns.ac.rs

Congenital tongue base cyst as uncommon cause of laryngeal stridor in an infant

Урођена циста базе језика као неуобичајен узрок ларингеалног стридора одојчета

SUMMARY

Introduction Congenital tongue base cysts are uncommon in everyday clinical practice and they could be asymptomatic or cause stridor, respiratory distress and asphyxia. We are presenting a case of two-month-old infant with the symptoms of stridor and the acute respiratory insufficiency caused by congenital tongue base cyst.

Case outline A two-month-old afebrile male infant was admitted to hospital with the symptoms of severe dyspnea, nonproductive cough and stridor. Due to severe respiratory insufficiency the infant was on mechanical ventilation. Computed tomography scans of thorax and neck was performed along with the application of the intravenous contrast where a cystic formation was shown. The depicted formation narrowed the lumen of oropharynx which is the same as the size of valleculas (3 mm). The formations pressed both valleculas, more significantly the left one. The same day the marsupialization of the cyst was done and the material was sent to the pathohistological analysis (the report of the pathologist indicates the cyst of the thyroid channel).

Conclusion Clinical manifestations of cyst depend on the level of obstruction and can be presented as inspiratory stridor, apnea, cyanosis, chronic coughing and feeding difficulty. The flexible nasopharyngeal laryngoscopy or bronchoscopy, CT and MR imaging help consider the differential diagnosis. The symptoms of stridor were removed completely after applied marsupialization of the cyst.

Keywords: airway obstruction; thyroglossal cyst; stridor; respiratory insufficiency

САЖЕТАК

Увод Урођене цисте базе језика не срећу се свакодневно у клиничкој пракси, а могу бити асимптоматске или изазивати стридор, отежано дисање или асфиксију. Приказујемо случај двомесечног одојчета са стридором и развијеном акутном респираторном инсуфицијенцијом услед притиска цисте на бази језика.

Приказ болесника Мушко афебрилно одојче, узраста два месеца је примљено у болницу због отежаног дисања, непродуктивног кашља и стридора. Услед тешког погоршања респираторне функције дете је прикључено на механичку вентилацију. Скенер главе и врата са применом контраста је детектовао присуство цисте у пределу врата тачније у корену језика. Описана формација је сузила лумен орофаринкса на 3 мм, колико је и лумен валекула. Циста је притискала обе валекуле, више леву, услед чега је настала опсежна опструкција дисајног пута. Након радиолошког налаза, урађена је марсупијализација цисте која је послата на патохистолошку анализу. Извештај патолога иде у прилог цисте тиреоглосног канала.

Закључак Клиничке манифестације цисте зависе од нивоа и обима опструкције, а могу да се испоље као инспираторни стридор, апнеа, цијаноза, хроничан кашаљ и потешкоће са храњењем. Флексибилна назофарингеална ларингоскопија, бронхоскопија, компјутеризована томографија и магнетно снимање могу да помогну у диференцијалној дијагностици. Симптоми стридора и отежаног дисања су потпуно нестали након марсупијализације цисте.

Кључне речи: опструкција дисајних путева; тиреоглосална циста; стридор; респираторна инсуфицијенција

INTRODUCTION

Laryngomalacia is considered to be the most common cause of stridor (noisy breathing) in the neonatal period and infancy [1]. The obstructions of respiratory airways are common, while congenital tongue base cysts are not frequent in everyday clinical practice [2]. A tongue base cyst may cause stridor, respiratory distress or be totally asymptomatic [3]. Owing to their specific position, tongue base cysts may cause perilous complications through mass effects on

the hypopharynx as well as by displacing the epiglottis. In the most serious cases cyst can lead to the fatal outcome due to asphyxia [4]. The mortality rate among the patients with the diagnosed congenital tongue cyst in the infant period vary, and in some studies goes to 40% [5]. Surgical treatment has been verified as the most efficient method with a significant improvement of the symptoms generated by the compressive effects of the cyst [6]. We are presenting a case of two-month-old infant with the symptoms of laryngeal stridor and the acute respiratory insufficiency caused by congenital tongue base cyst.

CASE REPORT

A two-month-old apyretic male infant was admitted to hospital with the symptoms of severe dyspnea, nonproductive cough and noisy breathing (stridor). He had been examined four days before admission by a pediatrician and the initial treatment included inhalation of short acting agonist beta 2 bronchodilators (fenoterol ipratropium bromide) and parenteral corticosteroids (methylprednisolone). The difficulties were persistent, and on the admission day the child was inconsolable crying for hours, breathing heavily and had cyanotic attacks. The child had been previously hospitalized for seven days due to bronchial obstruction and dyspnea caused by whooping cough (*Diagnosis Pertussis per Bordetella Pertussis*, confirmed by PCR technique) when he was only one month old. *DTP* (*Diphtheria*, *Tetanus* and *Pertussis*) vaccine should be given at the age of two months, and because the infant was only a month old he was given solely *Euvax-B*[®] (prevent hepatitis B) and *BSG* (*Bacillus Calmette Guerin*) vaccine (prevents *tuberculosis*) after labor.

On admission he was conscious and alert, pale, tachypneic with the respiratory rate (RR) of fifty breaths per minute, oxygen saturation of 97% on room air, heart rate of 150 beats per minute and the normal body weight of 4,800 g. In physical examination lung auscultation

revealed wheezing, bilateral early-inspiratory crackles on the lungs and inspiratory stridor. All other physical findings were normal. The initial analyses were within normal ranges (CRP 3.8 mg/L, WBC $10.9 \times 10^9/L$, RBC $3.77 \times 10^{12}/L$, HGB 101 g/L, HCT 29.0%, PLT $361 \times 10^9/L$). Testing of his arterial blood gases showed combined metabolic and respiratory acidosis (pH 7.291, pCO_2 4.42 kPa, pO_2 7.8 kPa, standard bicarbonate HCO_3 -std 16.6 mmol/L, base excess BE -9.7 mmol/L). Due to repeating noisy breathing and wheezing, alpha-1 antitrypsin (1.51 g/L) and total immunoglobulin E ($IgE < 30.00 \mu g/L$) were estimated and results were within referent values. Chest X-ray showed bilateral decreased transparency of lungs.

After admission, the inhalations with bronchodilator (ipratropium bromide/fenoterol hydrobromide) every four hours and corticosteroid (budesonid) every 12 hours were included. Respiratory physiotherapy (postural drainage aspiration) was applied. On the second day of the treatment around 13:00 p.m the deterioration of the respiratory status and symmetrically weakened breathing sound were noted. The noisy high tonal wheezing with the saturation drops up to 70 and heartbeat of over 200/min with occasional apneas lasting over 10 seconds occurred. The infant was tachypneic (RR/60 min), with ash gray skin, with visible intercostal and xiphoid retraction, nares dilatation (flaring) using additional intercostal musculature, with seesaw respiration. The deteriorating inspiratory stridor was present. The oxygen therapy through nasal canils initially 2L per min up to 6L per min was applied and systemic corticosteroid methylprednisolone was given. Because of the general status deterioration empiric antibiotic ceftazidime was included parenterally. The gas analyses were urgently done (pH 6.946, pCO_2 7.21 kPa, pO_2 7.6 kPa, BE(B)-20.9 mmol/ L, HCO_3 std 9.3 mmol/ L) indicating the non-compensative critical metabolic and respiratory acidosis. The inflammatory markers in the laboratory analyses were still normal, not showing the acute infection (CRP 2.2 mg/L, WBC $15.9 \times 10^9/L$, RBC $3.44 \times 10^{12}/L$, HGB 90 g/L, HCT 26.8%). The progressive deterioration of the respiratory status was caused by the obstruction of the upper

airways. Applied measures did not give signs of improvement. Due to the development of the respiratory insufficiency, the infant was intubated and on Synchronized Intermittent-Mandatory Ventilation (SIMV mechanical ventilation). During the evening hours of the same day the increase of the inflammatory parameters occurred (CRP 38.3 mg/L) so the therapy switched another antibiotic vancomycin. The parameters of the mechanical ventilation were gradually decreasing, so the infant was extubated the next day and initially transferred to HFNC (high flow nasal cannula) and then to oxygen therapy through nasal cannulas. Because the gas exchange was stable, the oxygen saturation satisfactory, the needs for additional oxygen ceased on the fourth day of hospitalization. Further on, the infant was in good general condition, while the parameters of acute inflammation were dropping.

The dual antibiotic therapy of ceftazidime and vancomycin continued during XIV days in total. At the same time, bronchodilator inhalations on every 6h went on, while the interval between the inhalations was gradually prolonged. The infant was put in the high headboard bed with his head tilted back because of the laryngeal stridor in order to relieve airways. The cardiologist and pulmonologist were consulted, all potential causes of stridor and difficult breathing in infancy were considered, so the directoscopy of larynx and rigid bronchoscopy were recommended. The otolaryngologist was consulted and unusual formation was detected. A computed tomography (CT) of thorax and thoracic neck was performed along with the application of the intravenous contrast where a cystic formation on the base of the tongue was shown (Figure 1). On the posterior tongue base in the middle section right above epiglottis and between valleculas and piriform recessuses, the clearly visible cystic formation with the dimensions of 12x9x9,4 mm was positioned, (AP *anterior posterior*, CC *cranial-caudal*, LL *latero-lateral*) liquid density without central, post contrast imbibition (Figure 2). The depicted formation narrows the lumen of oropharynx which is the same as the AP diameter of valleculas (3 mm). The formation compressed both valleculas, more significantly the left one.

Otorhinolaryngologist indicated the operative removal of the cyst. The directoscopy of the larynx and rigid bronchoscopy were performed and on the left side of the tongue base an oval whitish formation leaning on the left entrance of the larynx was spotted causing the difficult intubation. The trachea had normal ring-shaped form, with the membrane wall and preserved lumen. Other results of the bronchoscopy were normal. The same day the marsupialization of the cyst was done and the material was sent to the pathohistological analysis. After the operation parenteral antibiotic therapy with cefepime (cephalosporin of IV generation) was applied along with bronchodilator and corticosteroid. A few days later child was in a good general condition, with regular gas exchange and peroral nutrition (adapted milk formula), with gradually increased intake. At the control ORL examination the result after marsupialization of the cyst was normal, without stridor. The pathohistological analysis of the tissue showed that it was formed of smooth laminar layer of epithelium with mucosal glands. In the deeper layer there was transverse muscle tissue and edematous connective tissue. The report of the pathologist described the cyst of the thyroid channel.

Ethical standards: We confirm that we have read the journal's position on issues involving ethical publication and affirm that this work is consistent with those guidelines. Written consent to publish all shown material was obtained from parents.

DISCUSSION

Causes of heavy breathing in infants are numerous and can be the result of innate states, congenital anomalies or inherited diseases. Clinical manifestations depend on the level of obstruction (choanal atresia for instance, larynx) and the degree of the obstruction (different size of the cyst or some other formation that obstruct the airway [7]). The common cause of bronchial obstruction and vising in infancy is the atopic constitution of a child and the

tendencies of later development of allergic diseases (asthma, allergic rhinitis, conjunctivitis and eczemas) [8]. Acute airway obstruction can be life threatening condition which requests prompt reaction and treatment. Seldom, the deterioration of respiratory status can be that serious to require around the clock monitoring at the intensive care wards with the applied intubation and ventilation.

In medical literature, a variety of terms have been used for tongue base cysts, such as epiglottic cyst, lingual cyst, vallecular cyst, or laryngeal cyst [9]. Two major hypotheses to explain the pathogenesis of these cysts are the ductal obstruction of mucus glands or an embryological malformation [10]. Most affected infants have symptoms during the first week of life [11]. Clinical manifestations consist of various degrees of upper airway obstruction such as inspiratory stridor, chest retraction, apnea, cyanosis and feeding difficulty. Stridor is the most common symptom [12]. In neonatal stridor, evaluation of the airway anatomy and differential diagnosis from other causes of stridor are important to prevent any mortality and morbidity from these sources [10, 11]. Clinical presentation is usually related to upper respiratory tract. Obstruction and stridor are the most frequently encountered symptoms of vallecular cyst cases. Other symptoms include dyspnea, feeding difficulties, voice changes, chronic coughing and cyanotic attacks.

Definitive diagnosis can be obtained from bronchoscopy or laryngoscopy [13]. Primary diagnostic approach to laryngeal or vallecular cyst should be a flexible nasopharyngeal laryngoscopy or bronchoscopy. CT and magnetic resonance imaging (MRI) help narrow the differential diagnosis, and note differences between lingual, thyroid, proximal cystic dilatation of the thyroid duct, lymphangioma or hemangioma, dermoid cyst, lipoma, fibroma or carcinoma [14].

Timely differential diagnosis of laryngeal stridor and its consequences is crucial in critically ill patients with severe respiratory failure. Laryngomalacia is the most common cause of noisy breathing in infancy [15]. While having deteriorations of the respiratory status, our patient had the episodes of apnea (breathing breaks longer than 10 seconds followed by desaturation and bradycardia). It is significant to differ periodical breathing from apnea, the type of apnea (central, obstructive or mixed), as well as to determine if its cause is certain metabolic disorder which is manifested by apnea [16]. Subsequent management consists of determining the underlying etiology and instituting specific targeted therapy to the identified cause.

Congenital base cysts are rare and can be confused with laryngomalacia due to their nonspecific symptoms. They present a bulging mass at the base of tongue, obstructing the upper airway and are responsible for severe respiratory distress that sometimes can be fatal. Direct laryngoscopy, neck CT scan and pathological analysis were key for diagnosis. Marsupialization was the treatment of choice. The symptoms of heavy breathing were removed completely after applied marsupialization of the cyst. The incidence of recidivism (relapse) within the monitoring period is low.

ACKNOWLEDGMENT

We would like to express special thanks of gratitude to the radiologist Nikola Eić who helped us gathering graphical data and interpret them from other perspective.

Conflict of interest: None declared.

REFERENCES

1. Pokharel A. Prevalence of laryngomalacia among young children presenting with stridor in a tertiary care hospital. *J Nepal Med Assoc.* 2020;58(230):712–6. [DOI: 10.31729/jnma.5244] [PMID: 34504366]
2. Al-Mahboob A, Hajr E, Alammam A. Tongue Cyst in a Neonate: Unusual Presentation. *Cureus.* 2020;12(7):e9252. [DOI: 10.7759/cureus.9252] [PMID: 32821598]
3. Patnaik S, Zacharias G, Jain MK, Samantaray KK, Surapaneni SP. Etiology, Clinical Profile, Evaluation, and Management of Stridor in Children. *Indian J Pediatr.* 2021;88(11):1115–20. [DOI: 10.1007/s12098-021-03722-8] [PMID: 33728566]
4. Alharbi SM, Alahmari M, Al Hamoud M, Al Malwi K, Abujamilah H. Successful Management of a Symptomatic Vallecular Cyst in an Adult: A Case Report. *Cureus.* 2023;15(7):13–6. [DOI: 10.7759/cureus.41829] [PMID: 37575698]
5. Varlas VN, Parlatescu I, Epistatu D, Neagu O, Varlas RG, Bălănescu L. Mixed Heterotopic Gastrointestinal/Respiratory Oral Cysts in Newborns: From Prenatal Diagnosis to Histopathological and Therapeutic Management: A Case Report and Literature Review. *Diagnostics.* 2024;14(3):339. [DOI: 10.3390/diagnostics14030339] [PMID: 38337855]
6. Sato A, Kako E, Okuni N, Kusama N, Kamimura Y, Sento Y, et al. Perioperative management of thyroglossal duct cystectomy in a pediatric patient: A case report. *Clin Case Reports.* 2021;9(2):673–6. [DOI: 10.1002/ccr3.3607] [PMID: 33598223]
7. Barker N, Thevasagayam R, Ugonna K, Kirkby J. Pediatric Dysfunctional Breathing: Proposed Components, Mechanisms, Diagnosis, and Management. *Front Pediatr.* 2020;8(July):1–10. [DOI: 10.3389/fped.2020.00379] [PMID: 32766182]
8. Altman M, Calatroni A, Ramratnam S, Jackson D, Presnell S, Rosasco M, et al. Endotype of allergic asthma with airway obstruction in urban children. *The Journal of Allergy and Clinical Immunology.* 2021;148(5):1198–1209. [DOI: 10.1016/j.jaci.2021.02.040] [PMID: 33713771]
9. Lee EY, Juliano AF. Pediatric Tongue Lesions: Am J Roentgenol. 2020;(May):1–11. [DOI: 10.2214/AJR.19.22121] [PMID: 32069081]
10. Cunha JLS, Roza ALOC, Cruz VMS, Ribeiro JL, Cavalcante IL, Cavalcante RB, et al. Oral Lymphoepithelial Cyst: A Collaborative Clinicopathologic Study of 132 Cases from Brazil. *Head Neck Pathol.* 2022;16(1):268–77. [DOI: 10.1007/s12105-021-01352-2] [PMID: 34185247]
11. Neonatal Vallecular Cyst: Unusual Presentation with Near Catastrophic Respiratory Distress and Successful Treatment Using Endoscopy-assisted Plasma Ablation: A Case Report. *Journal of Postgraduate Medicine Education and Research.* 2020;54(1). [DOI: 10.5005/jp-journals-10028-1349]
12. Alnaimi A, Abushahin A. Vallecular Cyst: Reminder of a Rare Cause of Stridor and Failure to Thrive in Infants. *Cureus* 2021;13(11):e19692. [DOI: 10.7759/cureus.19692] [PMID: 34934567]
13. Shufen X, Shan S, Zhen L. Characteristics of lingual thyroglossal duct cysts under flexible laryngoscopy and treatment approaches. *Am J Otolaryngol - Head Neck Med Surg.* 2022;43(2):103343. [DOI: 10.1016/j.amjoto.2021.103343] [PMID: 35033924]
14. Wang Y. Case Report: A Rare Cause of Stridor and Hoarseness in Infants—Congenital Laryngeal Web. *Front Pediatr.* 2022;10(April):8–11. [DOI: 10.3389/fped.2022.875137] [PMID: 35463881]
15. Bredun S, Kotowski M, Mezydło J, Szydłowski J. Characteristics of Patients with Laryngomalacia: A Tertiary Referral Center Experience of 106 Cases. *Diagnostics.* 2023;13(20). [PMID: 37892001] [DOI: 10.3390/diagnostics13203180]
16. Thomas S, Patel S, Gummalla P, Tablizo MA, Kier C. You Cannot Hit Snooze on OSA: Sequelae of Pediatric Obstructive Sleep Apnea. *Children.* 2022;9(2):1–16. [DOI: 10.3390/children9020261] [PMID: 35204981]



Figure 1. Cystic formation on the base of the tongue



Figure 2. Sagittal computed tomography scan of neck with detected cystic formation at the base of tongue which narrow respiratory airway