



**СРПСКИ АРХИВ**  
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО  
**SERBIAN ARCHIVES**  
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: [office@srpskiarhiv.rs](mailto:office@srpskiarhiv.rs), Web address: [www.srpskiarhiv.rs](http://www.srpskiarhiv.rs)

**Paper Accepted\***

**ISSN Online 2406-0895**

**Case report / Приказ болесника**

Svetlana Kašiković-Lečić<sup>1,2</sup>, Milan Popović<sup>1,3,\*</sup>, Bosiljka Krajnović<sup>1,2</sup>, Milorad Bijelović<sup>1,2</sup>,  
Dragana Tegeltija<sup>1,2</sup>

**Nodular amyloidosis of the lung presenting as lung malignancy**

Нодуларна амилоидоза плућа презентована као малигни тумор плућа

<sup>1</sup>University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia;

<sup>2</sup>Institute for Pulmonary Diseases of Vojvodina, Sremska Kamenica, Serbia;

<sup>3</sup>Oncology Institute of Vojvodina, Sremska Kamenica, Serbia

**Received: September 15, 2022**

**Revised: November 2, 2022**

**Accepted: November 23, 2022**

**Online First: November 29, 2022**

**DOI: <https://doi.org/10.2298/SARH220915112K>**

\***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. Srp Arh Celok Lek. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

**\*Correspondence to:**

Milan POPOVIĆ

University of Novi Sad, Faculty of Medicine

Hajduk Veljkova 3, 2100 Novi Sad, Serbia

e-mail: [milan.popovic@mf.uns.ac.rs](mailto:milan.popovic@mf.uns.ac.rs)

## Nodular amyloidosis of the lung presenting as lung malignancy

### Нодуларна амилоидоза плућа презентована као малигни тумор плућа

#### SUMMARY

**Introduction** Amyloidosis is a disease associated with the extracellular deposition of insoluble protein material called amyloid. It can be acquired or hereditary, systemic or organ-limited. Nodular pulmonary amyloidosis is defined as one or more tumefactive amyloid deposits in the lungs.

**Case outlines** This study presents two cases that were hospitalized at the Institute for Pulmonary Diseases to clarify the origin of lesions detected on CT scans of lung parenchyma. In the first case, in a 78-year-old woman, numerous non-calcified nodules were described on the chest CT. The patient died during hospitalization, and the autopsy revealed diffusely distributed greyish-yellow nodular lesions in the upper and middle parts of the right lung, as well as lesions in the form of partially calcified nodules in both lungs. Histological analysis of samples from macroscopically described nodules confirmed nodular amyloidosis. The patient from the second presented case is a male, 58 years old, who was operated on for rectal adenocarcinoma three years ago. A CT scan of the lung parenchyma shows a tumor nodule localized in the lower lobe and a nodular lesion localized in the upper lobe of the right lung. Histological analysis confirmed that the lesion from the lower lobe corresponds to the metastasis of colorectal cancer, while in the lesion from the upper lobe amyloid deposits were found.

**Conclusion** Pulmonary nodular amyloidosis is a rare condition, and because of the imaging similarities it is difficult to distinguish it from malignant nodules in the lung parenchyma. Therefore, as a part of routine practice, a definitive diagnosis of amyloidosis needs to be confirmed by tissue biopsy.

**Keywords:** amyloidosis; amyloid; lung malignancy; autopsy; Congo red

#### САЖЕТАК

**Увод** Амилоидоза је болест повезана са ванћелијским таложењем нерастворљивог протеинског материјала који се назива амилоид. Обољење може бити стечено или наследно, системско или ограничено на органе. Нодуларна плућна амилоидоза се дефинише као један или више тумефактивних депозита амилоида у плућном паренхиму.

**Прикази случајева** Приказана су два случаја која су хоспитализована на Институту за плућне болести ради разјашњења етиологије, на ЦТ-у детектованих патолошких промена у плућном паренхиму. У првом описаном случају код особе женског пола, старости 78 година, на ЦТ-у грудног коша описују се бројни не калцификовани нодуси, међутим током хоспитализације дошло је до леталног исхода. На аутопсији се на плућима уочавају сивкасто-жућкасте нодулиране промене дифузно распоређене у горњим и средњим партијама десног плућног крила, као и промене у виду делимично калцификованих чворића локализоване у оба плућна крила. Хистолошким анализом узорака из макроскопски описаних нодуса доказана је нодуларна амилоидоза. Други пацијент је мушког пола, старости 58 година, који је пре три године оперисао аденокарцином ректума. На ЦТ-у плућног паренхима описује се туморски нодус који је локализован у доњем делу десног плућног крила као и нодулирана промена локализована у горњем режњу десног плућног крила. Хистолошким анализом постоперативних узорака утврђено је да нодус из доњег режња одговара метастази колоректалног карцинома, док се у узорку из горњег режња десног плућног крила налазе депозити амилоида.

**Закључак** Плућна нодуларна амилоидоза је ретко стање, а због сличности на дијагностичком имџингу тешко ју је разликовати од малигну тумора плућа, па се у рутинском раду коначна дијагноза амилоидозе мора поставити на биоптичком узорку ткива.

**Кључне речи:** амилоидоза; амилоид; плучни малигнитети; аутопсија; Congo red

## INTRODUCTION

Amyloidosis is a disease associated with the extracellular deposition of insoluble protein material called amyloid in various tissues and organs. This condition can be found in humans

as well as in other vertebrates. The term amyloid was used in medicine for the first time by Rudolf Virchow in 1854. Amyloid deposits are composed of fibril proteins (95%), and the remaining part is amyloid P and glycoproteins. Up to now, 36 amyloid fibril proteins and their precursors have been identified in humans, and ten in other vertebrates [1, 2].

Amyloidosis can be acquired or hereditary, systemic or organ-limited. Pulmonary amyloidosis can be seen in three distinct forms: diffuse alveolar-septal amyloidosis, nodular pulmonary amyloidosis, and tracheobronchial amyloidosis [3].

Nodular pulmonary amyloidosis is defined as one or more tumefactive amyloid deposits in the lungs and usually represents incidental findings on chest imaging [4]. The imaging similarities of multiple pulmonary nodules (with calcification and cavitation) with malignant lung tumours make differential diagnosis particularly difficult. For that reason, histological confirmation should be mandatory. In this case study, we report two cases of pulmonary nodular amyloidosis that were suspected of lung malignancy.

### **Case No. 1**

A 78-year-old woman was admitted to the Institute for Pulmonary Diseases of Vojvodina for cough, fever, haemoptysis and nausea. Upon admittance, at initial examination, there was not any palpable lymphadenopathy. Results of periphery blood showed anaemia and discrete leucocytosis, while the analysis of bio-humoral tests showed elevated values of inflammatory markers such as urea and LDH. In addition, a blood gas test showed hypercapnic respiratory failure.

Computed tomography (CT) scan of the chest has shown multiple non-calcified nodules in the right lung (between segments S2/S6 measuring 30×50 mm and S5 38x12 mm), and in the upper lobes of both lungs, and in the lower lobe of the left lung there were partly calcified nodules with a diameter of up to 24 mm. Cytological analysis showed that tracheobronchial

aspirate contained mucopurulent exudates.

Despite the whole therapy, intensive monitoring and care, on the 12<sup>th</sup> day of hospitalization the patient died. At the autopsy, the weight of the left lung was 670 g, while the weight of the right lung was 990 g. Both lungs were pinkish coloured on the cross-section, and a large amount of pale pink foamy fluid flew from its cut surface. In the upper and middle lobes of the right lung and the upper lobe of the left lung, individual nodules of greyish-yellow colour, less than 2 cm in diameter, were found. In the lower lobes on both sides of the lungs, there were greyish-yellow nodules, up to 4.6 cm in diameter (Fig. 1). Some of them had the appearance of grapes.

Histologically, nodules are composed of homogeneous, densely eosinophilic amorphous material. Deposits are orange-red on Congo red stain (fig. 2) and under polarized light show apple-green birefringence (fig. 2). In the surrounding lung tissue, during the pathohistological examination, haemorrhagic infarcts, chronic catarrhal bronchitis, emphysema, and pulmonary oedema (which was established as the cause of death) were revealed. Elements of malignant tissue were not seen.

## Case No. 2

The male patient, 58 years old, was admitted to the Institute for Pulmonary Diseases of Vojvodina for identification of aetiology of pathologic lesions in lung parenchyma which were found on CT imaging. His past medical history included arterial hypertension and dilated cardiomyopathy, and three years ago, the patient was diagnosed with adenocarcinoma of the rectum, which had been surgically removed.

During the surgery, whitish lesions were noted in the upper lobe of a lung. Tissue fragments from the lower and upper lobe of the right lung were sent for histopathological analysis.

Upon gross examination of lung parenchyma of the lower lobe of the lung, we found a clearly defined whitish tumour nodule, with 3.7x3 cm in size. In tissue samples taken from that area, we found a tumour tissue that, due to histological characteristics, corresponds to the metastasis of colorectal cancer.

In the histological samples from the upper lobe of the right lung in alveolar parenchyma there were nodular acellular homogeneous deposits that were eosinophilic by standard histological staining. Around described deposits, a lymphocyte infiltrate as well as giant cells were noted. Upon Congo red histochemical staining, acellular deposits were coloured orange, and under polarized light showed apple-green birefringence (fig. 2). Described findings correspond to nodular amyloidosis.

The study was approved by the Ethics Committee of the Institute for Pulmonary Diseases of Vojvodina.

## DISCUSSION

Amyloidosis is a rare disease of unclear aetiology, and its most common clinical forms are amyloid A amyloidosis (AA), immunoglobulin light chain amyloidosis (AL), transthyretin amyloidosis (ATTR), and Beta - 2 microglobulin amyloidosis ( $A\beta 2M$ ) [5,6].

The exact incidence of amyloidosis is unknown, but estimates show that 6-10 cases per million are diagnosed annually [7]. Amyloid deposits commonly affect males (2:1) of the middle age group (50–60 years) [8].

Amyloid can even affect pleura, pulmonary arteries, lymph nodes, and diaphragm [9, 10]. Deposition of amyloid results in destruction of tissue structure and impairment of organ function. The diagnosis of amyloidosis can only be confirmed by the histopathologic section of tissue stained with Congo Red, and the amyloid will show green apple birefringence under polarized light, as in this case. In addition, amyloid can be detected when stained with

metachromatic dye, such as crystal violet, or with thioflavin T as a fluorescent marker [11, 12].

Nowadays, the use of immunohistochemistry in combination with clinical testing can help identify precursor proteins in amyloid deposits and their subtyping [13, 14].

Mass spectrometry-based proteomic analysis can be performed on formalin-fixed, paraffin-embedded tissue, and due to its high sensitivity and specificity, this method can be applied in clinical biopsy specimens and it will be a useful tool in amyloid subtyping [5].

Clinically, nodular pulmonary amyloidosis is asymptomatic and on CT is seen as multiple nodules of varying size with sharp or lobulated margins, usually peripherally located in lower lobes of the lungs [4, 15]. Calcifications are seen in 50% of cases, and one of the characteristics is slow nodule growth with no regression [16, 17]. That is very similar to localisation of deposits in our first case, where we had amyloid nodules in lower lobes of both lungs, but the uncommon presentation was finding amyloid deposits in the middle lobe of the right lung.

For pulmonary nodular amyloidosis, differential diagnosis includes malignancy of the lung (primary or metastatic) and granulomatous diseases [18]. Previous study reported that during 18F-FDG PET / CT imaging of patients with amyloid deposits, most of them showed increased FDG uptake, but no FDG uptake has been mentioned [19]. Therefore, 18F-FDG PET / CT is not a useful diagnostic tool for distinguishing amyloidosis from lung malignancy, but in the future, the use of some more specific tracers such as C-PIB and 18F-florbetapir might be a better solution [20].

Pathohistological examination is essential for a definitive diagnosis. Differential diagnoses of nodular pulmonary amyloidosis on histological tissue specimens include pulmonary hyalinising granuloma and amyloid-like nodules in light-chain deposition disease. In light-chain deposition disease, amyloid-like nodules in lung parenchyma are histologically very similar to amyloid. However, they are composed of  $\kappa$ -light chain fragments which are

Congo red negative and do not form fibrils, but present as granular material under electron microscope [5].

Pulmonary hyalinising granuloma is a rare benign disease, and on chest imaging, shows unilateral or bilateral well-defined nodules. In histological slides, hyalinising granulomas are different from amyloid deposits. They are presented as fibrotic nodules composed of hyalinized collagen bundles with lamellar arrangement, so they are not homogeneous and the Congo red staining is negative [21]. Chronic inflammation is localised both around and within them. The inflammatory infiltrate predominantly contains lymphocytes and plasma cells, but it can also hold neutrophils and histiocytes [22].

Studies have shown that most cases of nodular pulmonary amyloidosis are associated with B-cell lymphomas [23] and MALT lymphoma (mucosa-associated lymphoid tissue lymphoma) [24], and association with pulmonary amyloidosis and lymphomas has been found in Sjögren's disease [25-27]. In the second case, we reported nodular amyloidosis in a patient with metastatic adenocarcinoma in the same lung.

Pulmonary nodular amyloidosis is a rare condition, but because of imaging similarities it is difficult to distinguish it from malignant nodules in the lung parenchyma. For that reason, it should be added to the differential diagnosis. In routine practice, a definitive diagnosis of amyloidosis needs to be confirmed by tissue biopsy and, in addition to standard H&E staining, histochemical Congo red should be used.

**Conflicts of interest:** None declared.

## REFERENCES

1. Benson MD, Buxbaum JN, Eisenberg DS, Merlini G, Saraiva MJM, Sekijima Y, et al. Amyloid nomenclature 2020: update and recommendations by the International Society of Amyloidosis (ISA) nomenclature committee. *Amyloid*. 2020;27(4):217-22. doi:10.1080/13506129.2020.1835263. PMID: 33100054.
2. Virchow R. Ueber eine im Gehirn und Rückenmark des Menschen aufgefunden Substanz mit der chemischen Reaction der Cellulose. *Arch für Pathol Anat und Physiol und für Klin Med*. 1854;6(1):135-8.
3. Gandham AK, Gayathri AR, Sundararajan L. Pulmonary amyloidosis: a case series. *Lung India*. 2019;36(3):229-232. doi:10.4103/lungindia.lungindia\_205\_18. PMID: 31031344.
4. Al-Umairi RS, Al-Lawati F, Al-Busaidi FM. Nodular pulmonary amyloidosis mimicking metastatic pulmonary nodules: a case report and review of the literature. *Sultan Qaboos Univ Med J*. 2018;18(3):e393-e396. doi: 10.18295/squmj.2018.18.03.023. PMID: 30607286.
5. Picken MM. The pathology of amyloidosis in classification: a review. *Acta Haematol*. 2020;143(4):322-34. doi: 10.1159/000506696. PMID: 32392555.
6. Ke PC, Zhou R, Serpell LC, Riek R, Knowles TPJ, Lashuel HA, et al. Half a century of amyloids: past, present and future. *Chem Soc Rev*. 2020;49(15):5473-509. doi: 10.1039/c9cs00199a. PMID: 32632432.
7. Kyle RA, Larson DR, Kurtin PJ, Kumar S, Cerhan JR, Therneau TM, et al. Incidence of AL amyloidosis in Olmsted County, Minnesota, 1990 through 2015. *Mayo Clinic Proc*. 2019;94(3):465-71. doi: 10.1016/j.mayocp.2018.08.041. PMID: 30713046.
8. Tanrıverdi E, Özgül MA, Uzun O, Gül Ş, Çörtük M, Yaşar Z, et al. Tracheobronchial amyloidosis mimicking tracheal tumor. *Case Rep Med*. 2016;2016:1-4. doi: 10.1155/2016/1084063. PMID: 27594885.
9. Khoor A, Colby TV. Amyloidosis of the Lung. *Arch Pathol Lab Med*. 2017 Feb;141(2):247-254. doi: 10.5858/arpa.2016-0102-RA. PMID: 28134587.
10. Lavatelli F, di Fonzo A, Palladini G, Merlini G. Systemic amyloidoses and proteomics: The state of the art. *EuPA Open Proteom*. 2016;11:4-10. doi: 10.1016/j.euprot.2016.02.003. PMID: 29900105.
11. Yakupova EI, Bobyleva LG, Vikhlyantsev IM, Bobylev AG. Congo red and amyloids: history and relationship. *Biosci Rep*. 2019;39(1):BSR20181415. doi: 10.1042/BSR20181415. PMID: 30567726.
12. Guselnikova VV, Antimonova OI, Fedorova EA, Shavlovsky MM, Safray AE, Rukavishnikova AA, et al. A novel method for amyloid detection in human tissue load using a fluorescent dye - Congo red analogue. *Sovrem Tekhnologii Med*. 2020;12(1):65-70. doi: 10.17691/stm2020.12.1.08. PMID: 34513039.
13. Barreca A, Bottasso E, Veneziano F, Giarin M, Nocifora A, Martinetti N, et al. Immunohistochemical typing of amyloid in fixed paraffin-embedded samples by an automatic procedure: Comparison with immunofluorescence data on fresh-frozen tissue. *PLoS One*. 2021;16(8):e0256306. doi: 10.1371/journal.pone.0256306. PMID: 34428239.
14. Menter T, Bachmann M, Grieshaber S, Tzankov A. A more accurate approach to amyloid detection and subtyping: combining in situ Congo red staining and immunohistochemistry. *Pathobiology*. 2017;84(1):49-55. doi: 10.1159/000447304. PMID: 27487341.
15. Rech JS, Arnulf B, de Margerie-Mellon C, Talbot A, Malphettes M, Vignon M, et al. Lower respiratory tract amyloidosis: presentation, survival and prognostic factors. A multicenter consecutive case series. *Am J Hematol*. 2019;94(11):1214-26. doi: 10.1002/ajh.25608. PMID: 31396978.
16. Nikzad S, Al-Arnawoot AA, Cabaero M, Rogalla P. An unusual case of nodular pulmonary amyloidosis. *Clin Case Rep*. 2021 Aug 7;9(8):e04562. doi: 10.1002/ccr3.4562. PMID: 34401154.
17. Polo-Nieto JF, Quiroga-Dussan MDP, Castañeda-González JP, Fierro-Rodríguez DM, Durán-Acuña R, Carrillo-Bayona JA. Perilymphatic micronodular pattern as a manifestation of pulmonary amyloidosis on high-resolution computed tomography. *Radiol Case Rep*. 2021;16(4):850-4. doi: 10.1016/j.radcr.2021.01.027. PMID: 33552338.
18. Baqir M, Schwegman AR, Maldonado F, Johnson TF, Ryu JH. Airway amyloidosis: a retrospective analysis of 43 patients. *J Bronchology Interv Pulmonol*. 2022. doi: 10.1097/LBR.0000000000000836. Epub ahead of print. PMID: 34984992.
19. Albano D, Premi E, Peli A, Camoni L, Bertagna F, Turrone R, Borroni B, Calhoun VD, Rodella C, Magoni M, Padovani A, Giubbini R, Paghera B. Correlation between brain glucose metabolism (18F-FDG) and cerebral blood flow with amyloid tracers (18F-Florbetapir) in clinical routine: Preliminary evidences. *Rev Esp Med Nucl Imagen Mol (Engl Ed)*. 2022;41(3):146-52. doi: 10.1016/j.remnie.2021.03.016. PMID: 35577490
20. Prato FS, Wisenberg G. Reproducibility and repeatability of assessment of myocardial light chain amyloidosis burden using 18F-florbetapir PET/CT. *J Nucl Cardiol*. 2021;28(5):2011-3. doi: 10.1007/s12350-019-01971-x. PMID: 31797317.
21. Rahi MS, Gunasekaran K, Amoah K, Chowdhury F, Kwon J. Pulmonary hyalinizing granuloma: a rare cause of a benign lung mass. *Clin Pract*. 2021;11(1):37-42. doi: 10.3390/clinpract11010007. PMID: 33572722.
22. Tang I, Moore AJ, Fryer E, Sykes A. Pulmonary hyalinising granuloma: a rare and elusive cause of multiple lung nodules. *BMJ Case Rep*. 2020;13(3):e233327.



doi: 10.1136/bcr-2019-233327. PMID: 32217660.


23. Ohara S, Tomizawa K, Shimizu S, Suda K, Fujino T, Hamada A, et al. Primary pulmonary mucosa-associated lymphoid tissue lymphoma with amyloid light chain-type amyloidosis. *Surg Case Rep.* 2019;5(1):105. doi: 10.1186/s40792-019-0663-0. PMID: 31243589.

24. Lantuejoul S, Moulai N, Quetant S, Brichon PY, Brambilla C, Brambilla E, et al. Unusual cystic presentation of pulmonary nodular amyloidosis associated with MALT-type lymphoma. *Eur Respir J.* 2007;30(3):589–92. doi: 10.1183/09031936.00136605. PMID: 17766635.


25. Kinoshita Y, Ikeda T, Miyamura T, Ueda Y, Yoshida Y, Ueda M, et al. Pulmonary amyloidosis associated with Sjögren's syndrome. *Intern Med.* 2022;61(6):877-881. doi: 10.2169/internalmedicine.8169-21. PMID: 34471033.

26. Wu CW, Chien YC. Sjögren's syndrome with nodular pulmonary amyloidosis. *Respirol Case Rep.* 2021;9(1):e00699. doi: 10.1002/rcr2.699. PMID: 33343903.

27. Kim J, Kim YS, Lee HJ, Park SG. Pulmonary amyloidosis and multiple myeloma mimicking lymphoma in a patient with Sjogren's syndrome: A case report. *World J Clin Cases.* 2022;10(3):1016-23. doi: 10.12998/wjcc.v10.i3.1016. PMID: 35127915.



**Figure. 1.** Gross images of lung with amyloid nodules



**Figure. 2.** Congo red stained lung parenchyma with amyloid deposit, examined under bright field and polarized light microscopy,  $\times 4$