

ORIGINAL ARTICLE / ОРИГИНАЛНИ РАД

Cardiac surgery in patients with chronic renal failure

Ksenija Babović-Stanić^{1,2}, Jadranka Dejanović^{1,2}, Aleksandra Vulin^{1,2}, Lazar Velicki^{1,2}, Aleksandar Redžek^{1,2}¹University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia²Vojvodina Institute of Cardiovascular Diseases, Sremska Kamenica, Serbia;**SUMMARY**

Introduction/Objective Patients with chronic renal failure (CRF) undergoing cardiac surgery are believed to have more postoperative complications and significantly higher mortality rate.

The aim of the paper was to determine preoperative predictors of exacerbation of CRF and the outcome in patients with CRF submitted to cardiac surgery.

Methods A retrospective study included 169 patients hospitalized from 2012 to 2015 (age 67.71 ± 8.46 years, 72.3% male). The analysis included numerous perioperative characteristics.

Results Preoperative stage I CRF was present in 62 (37%), stage II in 77 (46%), and stage III–V in 30 (17%) patients. Exacerbation of CRF was registered in 37 (21.9%), and the lethal outcome in 16 (9.5%) patients. Stage II of CRF (odds ratio [OR] 4.76; 95% confidence interval [CI] 1.31–17.28; $p = 0.018$) and stage III–V of CRF (OR 11.39; 95% CI 2.87–45.14; $p = 0.001$) were designated as predictors for exacerbation of CRF following cardiac surgery. In patients with CRF stage I and II, multivariate analysis designated previous cerebrovascular insult (OR 3.36; 95% CI 1.04–10.93; $p = 0.044$) and ejection fraction $\leq 35\%$ (OR 5.35; 95% CI 1.83–15.64; $p = 0.02$) as predictors for the exacerbation of CRF. The only predictor of postoperative dialysis requirement was higher stage of CRF (OR 5.81; 95% CI 1.22–27.81; $p = 0.028$). CRF stage III–V was a predictor of lethal outcome (OR 7.64; 95% CI 1.49–39.27; $p = 0.015$).

Conclusion Higher stage of CRF in patients submitted to cardiac surgery is a predictor of exacerbation of renal failure and the lethal outcome.

Keywords: chronic renal failure; cardiac surgeries; morbidity; mortality

INTRODUCTION

There is a significant increase in the number of patients with chronic renal failure (CRF) who require cardiac surgery. Heart diseases are more frequent in this specific population compared to the general population [1]. Terminal renal failure is recognized as a significant risk factor for the outcome of cardiac surgery. The prognosis of patients diagnosed with cardiovascular disease and CRF is much worse than in patients without this associated morbidity. In patients with impaired renal function, cardiac surgery may cause aggravation of pre-existing renal failure or irreversible renal damage [1]. The problems these patients have are further complicated during open heart surgery when extracorporeal circulation is used.

In order to reduce perioperative risk, an optimal perioperative strategy is required. Most published studies emphasize the importance that preoperative clinical status has on postoperative mortality and morbidity in this category of patients, but little is known about the predictive factors for long-term survival. In patients with mild to moderate reduction of glomerular filtration rate, there is quite consistent evidence that surgical revascularization is a better therapeutic option compared to percutaneous coronary intervention (PCI) [2]. This especially applies to patients with CRF caused by diabetes

mellitus, who are recommended “off pump” surgery [3, 4]. In patients with the terminal phase of the renal disease, there is no such clear evidence in favor of surgical approach. Better long-term results in these patients are being achieved with the surgical approach, but with the higher rate of intra-hospital events and complications, whereas vice versa is true for PCI comparing to coronary artery bypass grafting (CABG).

Only 15 years ago, patients with CRF were believed to have unacceptably high operative risk and only rarely underwent cardiac surgery [5, 6]. Perioperative treatment strategy of these patients in intensive care units is constantly improving. Based on the research results and improving clinical practice results, the attitude towards candidates for cardiac surgery with CRF has been dramatically changed.

In our study, we sought to determine the following:

- 1) Present preoperative clinical CRF patient profile;
- 2) Examine occurrence of postoperative morbidity and mortality in patients with CRF who underwent cardiac surgery.

METHODS

This retrospective analysis included 169 patients with CRF who underwent cardiac sur-

Примљено • Received:

October 25, 2016

Ревизија • Revised:

March 21, 2017

Прихваћено • Accepted:

March 31, 2017

Online first: March 31, 2017

Correspondence to:

Lazar VELICKI
Institute of cardiovascular
diseases Vojvodina
Put doktora Goldmana 4, 21204
Sremska Kamenica, Serbia
lazar.velicki@mf.uns.ac.rs

gery (coronary, valvular, and combined) during the 2012–2015 period, at the Vojvodina Institute of Cardiovascular Diseases. Patients with CRF were divided into three groups according to the creatinine clearance (CC) values [1]. The first group comprised patients with $CC \geq 90$ mm/min./1.73 m², the second group – patients with CC 60–89 mm/min./1.73 m², and the third group – patients with CC 15–59 mm/min./1.73 m². The third group included all the patients with CRF stage III, IV, and V, considering the small number of patients in each stage separately. The following comorbidities and patient characteristics were assessed: high blood pressure (> 140/90 mmHg), diabetes, blood lipid levels, chronic obstructive pulmonary disease, peripheral arterial occlusive disease, cerebrovascular insult, myocardial infarction, heart failure (HF) class using New York Heart Association (NYHA) III/IV classification, left ventricle ejection fraction (EF) \leq 35%. The following postoperative complications were assessed: new onset of myocardial infarction, heart rhythm disturbances, cerebrovascular insult, sepsis, sternal infection, return to the intensive care unit, respiratory insufficiency, pericardial effusion, exacerbation of renal failure, and lethal outcome.

Heart surgeries were performed using extracorporeal circulation, in moderate hypothermia and perfusion pressure > 50 mmHg. Heart was stopped using the cold antegrade St Thomas' Hospital cardioplegia (with 40 mmol/l K⁺). Intraoperative and postoperative monitoring included electrocardiography, transesophageal echocardiography, oxygen saturation (pulse oximetry), diuresis, continual measurement of arterial, central venous, and pulmonary arterial pressure.

Criteria for the diagnosis of HF were defined as the need to ensure hemodynamic support with the inotropes – adrenaline, dobutamine (> 5 mg/kg), and/or high doses of dopamine for longer than 24 hours, and the value of cardiac index < 2.0 l/min./m² [2].

Postoperative myocardial infarction was defined as > 10-fold increase of cardiac enzymes (troponin and CK-MB) following the surgery compared to reference levels [5]. Heart rhythm disturbances were defined as the new onset of arrhythmia – atrial fibrillation or ventricular tachycardia. Cerebrovascular event was defined as an isolated neurological deficit after the surgery. Respiratory failure was defined as a requirement for mechanical ventilation for longer than 24 hours. Septic attack was diagnosed on the basis of sepsis criteria (white blood cell count, body temperature, respiratory rate, inflammatory mediators: CRP, fibrinogen, procalcitonin, and positive blood cultures). Pericardial effusion was defined as the presence of abnormal amount of fluid in pericardial cavity – more than 100 ml. Exacerbation of renal failure was defined as significant increase of nitrogenous substances in blood comparing to preoperative values [4]. Hospital mortality was defined as any death that occurred during index hospitalization.

Statistical analysis was performed using IBM SPSS Statistics for Windows, Version 19.0 (IBM Corp., Armonk, NY, USA). A p-value of < 0.05 was considered statistically




Figure 1. Representation of patients with chronic renal failure based on creatinine clearance; 1 – patients with creatinine clearance ≥ 90 mm/min./1.73 m²; 2 – patients with creatinine clearance of 60–89 mm/min./1.73 m²; 3 – patients with creatinine clearance of 15–59 mm/min./1.73 m²

significant. The Kolmogorov–Smirnov test was used for the determination of quantitative data distribution. Differences of mean values were tested by the independent samples t-test or Mann–Whitney U-test and the results are presented as mean (standard deviation). The relations between categorical variables were tested using the χ^2 test and the results are presented as frequencies and percentages. Univariate and multivariate binary logistic regression were performed to determine the effects of all the factors on the dependent variable. Only variables designated as significant by a univariate analysis were entered into multivariate regression analysis.

RESULTS

The study was conducted on 169 CRF patients, mean age 67.71 ± 8.46 years, of whom 125 patients (73.9%) were males. The number and percentage of patients according to the CRF stage are presented in Figure 1. The majority of patients (45%) were classified as CRF stage II.

High blood pressure was detected in 166 patients (98.2%). There was no statistically significant difference between the number of patients with high blood pressure within different stages of renal failure ($p > 0.05$). Diabetes mellitus was present in 64 (37.8%) patients. There was no statistically significant difference in the incidence of diabetes mellitus throughout renal failure stages ($p > 0.05$).

Significant difference was noted in the incidence of patients with HF ($p = 0.018$), NYHA III/IV stage ($p = 0.032$), and $EF \leq 35\%$ according to the renal failure stage. Comparing patients with stage II CRF to those with stage I, there was statistically significant difference in the incidence of patients with NYHA III/IV stage ($p = 0.017$) and $EF \leq 35\%$ ($p = 0.013$). There was statistically significant difference in the incidence of patients with HF ($p = 0.005$), NYHA III/IV ($p = 0.022$), and $EF \leq 35\%$ ($p = 0.026$) when comparing patients with stage III–IV CRF to those with stage I.

Previous myocardial infarction (MI) was present in 83 (49.1%) patients. There was no significant difference in the incidence of MI throughout the stages of renal failure ($p > 0.05$) (Table 1).

Table 1. Preoperative characteristics of patients with CRF

Variable	Renal failure stage			p
	1	2	3	
	n (%)	n (%)	n (%)	
SEX (male)	49 (79%)	51 (66.2%)	25 (73.5%)	0.242
HBP	60 (96.8%)	76 (98.7%)	34 (100%)	0.473
DM	19 (30.6%)	32 (41.6%)	13 (38.2%)	0.410
HLP	45 (72.6%)	56 (72.7%)	25 (73.5%)	0.995
COPD	12 (19.4%)	11 (14.3%)	6 (18.2)	0.712
PAOD	4 (6.5%)	3 (3.9%)	1 (2.9%)	0.677
CVI	7 (11.3%)	16 (20.8%)	5 (14.7%)	0.309
MI	28 (45.2%)	36 (46.8%)	19 (55.9%)	0.579
HF	19 (30.6%)	32 (41.6%)	20 (60.6)	0.018
NYHA III/IV	8 (12.9%)	23 (29.9%)	11 (32.4%)	0.032
EF ≤ 35	7 (11.3%)	22 (28.6%)	10 (29.4%)	0.030

HBP – high blood pressure; DM – diabetes mellitus; HLP – hyperlipoproteinemia; COPD – chronic obstructive pulmonary disease; PAOD – peripheral arterial occlusive disease; CVI – cerebrovascular insult; MI – myocardial infarction; HF – heart failure; NYHA – New York Heart Association; EF – ejection fraction




Figure 2. Representation of patients with postoperative heart failure according to chronic renal failure stage; 1 – patients with creatinine clearance ≥ 90 mm/min./1.73 m²; 2 – patients with creatinine clearance of 60–89 mm/min./1.73 m²; 3 – patients with creatinine clearance of 15–59 mm/min./1.73 m²

The number of patients who underwent coronary surgery was 90 (53.2%), valvular surgery 36 (21.3%), and combined surgery 43 (25.5%). Mean duration of extracorporeal circulation was 88.23 ± 35.52 min.

There was no significant difference in the incidence of patients with the new onset of MI in regard to the CRF stage ($p > 0.05$). A new onset of MI was a rare complication after surgery in patients with CRF. There was no significant difference in the incidence of patients with arrhythmias, CVI and sepsis rate in regard to the CRF stage ($p > 0.05$).

Significant difference in the incidence of respiratory failure with regard to the CRF stage ($p = 0.049$) was noted, and it was most notable between CRF stage I and stages III–IV ($p = 0.016$). Postoperative occurrence of HF – defined as the need to ensure hemodynamic support with the inotropes (adrenaline, dobutamine > 5 mg/kg, and/or high doses of dopamine for periods longer than 24 hours), and the value of cardiac index < 2.0 l/min./m² – is presented in Figure 2 ($p < 0.0001$).

A highly significant difference was present in the incidence of acute exacerbation of renal failure with regard to the CRF stage ($p < 0.01$) – between the CRF stages I and II ($p = 0.011$), II and III–V ($p = 0.016$), and between the

Table 2. Postoperative complication in patients with CRF

Postoperative complications	Renal failure stage			p
	1	2	3	
	n (%)	n (%)	n (%)	
NOMI	0 (0%)	2 (2.6%)	2 (5.9%)	0.185
NOHA	24 (38.7%)	35 (45.5%)	15 (44.1%)	0.716
NOCVI	2 (3.2%)	4 (5.2%)	0 (0%)	0.383
SEPSIS	2 (3.2%)	3 (3.9%)	1 (2.9%)	0.690
STERINF	2 (3.2%)	4 (5.2%)	1 (2.9%)	0.788
RET to ICU	6 (9.7%)	10 (13%)	7 (20.6%)	0.320
RESP FAIL	4 (6.5%)	9 (11.7%)	8 (23.5%)	0.049
PER EFF	5 (8.1%)	12 (15.6%)	5 (14.7%)	0.387
AC RF	3 (4.8%)	15 (19.5%)	14 (41.2%)	0.000
Mortality	2 (3.3%)	7 (9.1%)	7 (20.6%)	0.021

NOMI – new onset of myocardial infarction; NOHA – new onset of heart arrhythmias; NOCVI – new onset of cerebrovascular insult; STERNINF – sternal infection; RET to ICU – return to the intensive care unit; RESP FAIL – respiratory failure; PER EFF – pericardial effusion; AC RF – acute exacerbation of renal failure

CRF stages I and III–V ($p < 0.01$). Also, there was a significant difference in the mortality rate in regard to the CRF stage ($p = 0.021$) – between the CRF stages I and III–V ($p = 0.006$) (Table 2).

A total of 16 deaths were registered. Thereof, seven patients underwent coronary surgery, six underwent combined surgery and three patients underwent valve surgery. Seven patients were in CRF stage III–V, seven patients were in CRF stage II, and two patients were in CRF stage I.

Univariate analysis designated CRF stage as a predictor of the occurrence of respiratory failure after surgery (OR = 2.1; 95% CI 1.12–4.07; $p = 0.021$) and CRF stage III–V (OR = 4.46; 95% CI 1.23–16.15; $p = 0.023$). CRF stage was also the predictor of in-hospital mortality (OR = 2.72; 95% CI 1.27–5.08; $p = 0.009$) and CRF stage III–V (OR = 7.64; 95% CI 1.49–39.27; $p = 0.015$). Preoperative CRF stage was the predictor of acute exacerbation of renal failure following surgery (OR = 3.14; 95% CI 1.69–5.79; $p < 0.01$), stage II (OR = 4.76; 95% CI 1.31–17.28; $p = 0.018$), and CRF stage III–V (OR = 11.39; 95% CI 2.87–45.14; $p = 0.001$).

In patients with CRF stage I and II, the univariate predictors of acute exacerbation of CRF were preoperative CVI (OR = 3.06; 95% CI 1.01–9.25; $p = 0.048$), NYHA III/IV (OR = 4.5; 95% CI 1.60–12.64; $p = 0.004$), EF $\leq 35\%$ (OR = 5.05; 95% CI 1.78–14.30; $p = 0.002$), and CRF stage (OR = 4.76; 95% CI 1.31–17.28; $p = 0.018$). In patients with CRF stage I and II, the multivariate predictors of acute exacerbation of CRF were previous CVI (OR = 3.36; 95% CI 1.04–10.93; $p = 0.044$) and EF $\leq 35\%$ (OR = 5.35; 95% CI 1.83–15.64; $p = 0.02$).

DISCUSSION

The global number of cardiac surgery procedures performed in patients with CRF is constantly increasing [2, 3]. According to the available data, adverse cardiovascular events are the most important cause of death in patients with CRF. The development of exacerbation of CRF is correlated with substantial short- and long-term morbidity

and mortality. The pathogenesis is multifactorial. Hemodynamic, inflammatory, metabolic and nephrotoxic factors are involved and overlap each other, which leads to further aggravation of renal failure. Preoperative risk factors include advanced age, impaired left ventricular function or congestive HF, diabetes, chronic obstructive pulmonary disease, and the urgency of the operation [7, 8].

In large studies that have addressed this issue, the most commonly described postoperative morbidity included: respiratory failure, arrhythmias, acute exacerbation of renal insufficiency, pericardial effusion. Most of these complications might be explained, to an extent, through hypervolemic state developed early after the surgery (inadequate fluid management). Contrary to our findings, the need for dialysis was over 15% [9].

The high frequency of postoperative arrhythmias can be compared with the results of other studies as a result of electrolyte imbalance in the perioperative period [10–15]. The tendency to infection is a common feature for these patients, due to reduced chemotaxis, lymphopenia, decreased cell-mediated immunity, and reduction functions of monocytes. In our study, patients underwent antibiotic protection so no significant occurrence of infection was recorded compared to other studies where the presence of sternal infection was around 8% [16–19].

A retrospective analysis of patients undergoing CABG at the Cleveland Clinic found that the level of creatinine over 168 mmol/l results in a higher perioperative morbidity (2.8%) and mortality (3.7%). Hospital mortality in patients with CRF submitted to cardiac surgery goes up

to 36.7% as reported in large multicenter studies [20–23]. Our results are in line with the reports of cardiovascular centers around the world, and also with the largest multicenter study performed so far, in which the mortality was 12.5% [24]. Our analyses designated the following factors as predictors of postoperative mortality of CRF patients: COPD and diabetes. Isolated coronary surgery is associated with the lowest incidence of acute exacerbation of CRF, followed by valvular and combined operations as is the case with our group of patients [25].

In terms of prevention of acute exacerbation of renal failure after surgery, a proper preoperative patient management is required, especially for those on chronic dialysis, patients with pulmonary diseases and metabolic disorders (diabetes). As our understanding of the pathogenesis of renal injury following cardiac surgery grows, better preventive and therapeutic strategies will arise. Current approaches include deferring elective surgery, until there is adequate recovery following pre-existing renal injury, careful pre-operative risk stratification of patients, and consideration of less invasive procedures in those at greatest risk [26].

CONCLUSION

Preoperative chronic renal failure leads to an increased morbidity and mortality in patients submitted to cardiac surgery. Therefore, a careful preoperative evaluation is warranted as well as optimal perioperative management and treatment strategy for the purpose of risk reduction.

REFERENCES

- National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Am J Kidney Dis.* 2002; 39(2 Suppl 1): S1–266.
- Witczak B, Hartmann A, Svennevig JL. Multiple risk assessment of cardiovascular surgery in chronic renal failure patients. *Ann Thorac Surg.* 2005; 79(4):1297–302.
- Niyahara K, Maeda M, Sakurai H, Nakayama M, Murayama H, Hasegawa H. Cardiovascular surgery in patients on chronic dialysis: Effect of intraoperative hemodialysis. *Interact Cardiovasc Thorac Surg.* 2004; 3(1):148–52.
- Herzog CA, Ma JZ, Collins AJ. Long-term survival of dialysis patients in the United States with prosthetic heart valves: Should ACC/AHA practice guidelines on valve selection be modified? *Circulation.* 2002; 105(11):1336–41.
- Horai T, Fukui T, Tabata M, Takanashi S. Early and midterm results of off-pump coronary artery bypass grafting in patients with end stage renal disease: Surgical outcomes after achievement of complete revascularization. *Interact Cardiovasc Thorac Surg.* 2008; 7(2):218–21.
- Zanardo G, Michielon P, Paccagnella A, Rosi P, Caló M, Salandin V, et al. Acute renal failure in the patient undergoing cardiac operation: Prevalence, mortality rate, and main risk factors. *J Thorac Cardiovasc Surg.* 1994; 107(6):1489–95.
- Gultekin B, Ozkan S, Uguz E, Atalay H, Akay T, Arslan A, et al. Valve Replacement surgery in patients with end-stage renal disease: Long-term results. *Artif Organs.* 2005; 29(12):972–5.
- Taggart DP, D'Amico R, Altman DG. Effect of arterial revascularisation on survival: a systematic review of studies comparing bilateral and single internal mammary arteries. *Lancet.* 2001; 358(9285):870–5.
- Glasscock R, Winearls C. An epidemic of chronic kidney disease: fact or fiction. *Nephrol Dial Transplant.* 2008; 23(4):1117–21.
- Liu J, Lin C, Chua C, Chiang S, Hung H, Lu M, et al. Outcome of off-pump coronary artery bypass in renal dialysis patients. *Thorac Cardiovasc Surg.* 2008; 56(7):412–7.
- Cooper WA, O'Brien SM, Thourani VH, Guyton RA, Bridges CR, Szczech LA, et al. Impact of renal dysfunction on outcomes of coronary artery bypass surgery: Results from the Society of Thoracic Surgeons National Adult Cardiac Database. *Circulation.* 2006; 113(8):1063–70.
- Hannan EL, Wu C, Walford G, Culliford AT, Gold JP, Smith CR, et al. Drug-eluting stents vs. coronary-artery bypass grafting in multivessel coronary disease. *N Engl J Med.* 2008; 358(4):331–41.
- Nicolini F, Beghi C, Muscari C, Agostinelli A, Budillon A, Spaggiari I, et al. Myocardial protection in cardiac surgery: Current options and future challenges. *Eur J Cardiothorac Surg.* 2003; 24(6):986–92.
- Durmar I, Bukit S, Atay Y, Yagch T, Orburam M, Boga M, et al. Cardiac surgery with cardiopulmonary bypass in patients with chronic renal failure. *J Thorac Cardiovasc Surg.* 1999; 118(2):306–15.
- Bechtel MJF, Detter C, Fischlein T, Krabatsch T, Osswald BR, Riess F, et al. Cardiac surgery in patients on dialysis: Decreased 30-day mortality, unchanged overall survival. *Ann Thorac Surg.* 2008; 85(1):147–53.
- Krabatsch T, Yeter R, Hetzer R. Coronary surgery in patients requiring chronic hemodialysis. *Kidney Blood Press Res.* 2005; 28(5-6):270–4.
- Taylor J. Third Universal definition of myocardial infarction. *Eur Heart J.* 2012; 33(20):2506–7.
- Levey AS, Jong PE, Coresh J, El Nahas M, Astor BC, Matsushita K, et al. The definition, classification and prognosis of chronic kidney disease: a KDIGO Controversies Conference report. *Kidney Int.* 2011; 80(1):17–28.
- Rahmanian PB, Adams DH, Castillo JG, Vassalotti J, Filsoufi F. Early and late outcome of cardiac surgery in dialysis-dependent patients: Single-center experience with 245 consecutive patients. *J Thorac Cardiovasc Surg.* 2008; 135(4):915–22.
- Horst M, Mehlhorn U, Hoerstrup SP, Suedkamp M, de Vivie ER. Cardiac surgery in patients with end-stage renal disease: 10-year experience. *Ann Thorac Surg.* 2000; 69(1):96–101.

21. Boku N, Masuda M, Eto M, Nishida T, Morita S, Tominaga R. Risk evaluation and midterm outcome of cardiac surgery in patients on dialysis. *Asian Cardiovasc Thorac Ann.* 2007; 15(1):19–23.
22. Kato W, Tajima K, Terasawa S, Tanaka K, Usui A, Ueda Y. Results of isolated valve replacement in hemodialysis patients. *Asian Cardiovasc Thorac Ann.* 2007; 15(5):386–91.
23. Unić-Stojanović D, Miličić M, Vuković P, Babić S, Jović M. Kardiohirurške intervencije kod bolesnika na hroničnom programu hemodijalize. *Med Pregl.* 2013; 66(1-2):64–9.
24. Luciani N, Nasso G, D'alessandro C, Testa F, Glieda F, Gaudino M, et al. Heart surgery interventions in chronic dialysis patients: short- and long-term results. *Ital Heart J Suppl.* 2002; 3(7):746–52.
25. Kolh P. Renal insufficiency after cardiac surgery: a challenging clinical problem. *Eur Heart J.* 2009; 30(15):1824–7.
26. Filsoufi F, Rahmanian PB, Castillo JG, Silvey G, Carpentier A, Adams DH. Predictors and early and late outcomes of dialysis-dependent patients in contemporary cardiac surgery. *J Cardiothorac Vasc Anesth.* 2008; 22(4):522–9.

Кардиохирургија код болесника са хроничном бубрежном инсуфицијенцијом

Ксенија Бабовић-Станић^{1,2}, Јадранка Дејановић^{1,2}, Александра Вулин^{1,2}, Лазар Велици^{1,2}, Александар Реџек^{1,2}

¹Универзитет у Новом Саду, Медицински факултет, Нови Сад, Србија;

²Институт за кардиоваскуларне болести Војводине, Сремска Каменица, Србија

САЖЕТАК

Увод/Циљ Сматра се да болесници са хроничном бубрежном инсуфицијенцијом (ХБИ) после кардиохируршких операција имају више оперативних компликација и знатно повећану смртност.

Циљ рада је био да се одреде преоперативни показатељи погоршања ХБИ и смртог исхода код болесника са ХБИ подвргнутих кардиохируршким операцијама.

Методе Ретроспективно су анализирани периоперативне карактеристике код 169 болесника (старости $67,71 \pm 8,46$ година, 72,3% мушкараца) лечених од 2012. до 2015.

Резултати Преоперативно, у I стадијуму ХБИ било је 62 (37%), у II – 77 (46%), а III–V стадијуму 30 (17%) болесника. Погоршање ХБИ регистровано је код 37 (21,9%), а смртни исход код 16 болесника (9,5%). II стадијум ХБИ (OR 4,76; 95%

CI 1,31–17,28; $p = 0,018$) и III–V стадијум ХБИ (OR 11,39; 95% CI 2,87–45,14; $p = 0,001$) били су предсказатељи погоршања ХБИ. Код болесника са I и II стадијумом ХБИ, мултиваријантни предсказатељи погоршања ХБИ били су претходни показатељи удара (OR 3,36; 95% CI 1,04–10,93; $p = 0,044$) и ејекциона фракција $\leq 35\%$ (OR 5,35; 95% CI 1,83–15,64; $p = 0,02$). Једини предсказатељ постоперативне дијализе је виши стадијум ХБИ (OR 5,81; 95% CI 1,22–27,81; $p = 0,028$). III–V стадијум ХБИ био је предсказатељ смртог исхода (OR 7,64; 95% CI 1,49–39,27; $p = 0,015$).

Закључак Тежи стадијум ХБИ код болесника подвргнутих кардиохируршким операцијама јесте предсказатељ погоршања бубрежне инсуфицијенције и смртог исхода.

Кључне речи: хронична бубрежна инсуфицијенција; кардиохирургија; морбидитет; mortalитет