



REVIEW ARTICLE / ПРЕГЛЕДНИ РАД

Invasive diagnostic procedures for early-stage lung cancer – the clinical significance of novel navigational techniques in interventional bronchoscopy

Spasoje Popević^{1,2}, Nensi Lalić^{3,4}, Marko Bojović^{3,5}, Rade Milić^{6,7}, Ivica Lalić⁸, Branislav Ilić^{1,2}, Ivana Sekulović-Radovanović², Dane Krtnić^{9,10}

¹University of Belgrade, Faculty of Medicine, Belgrade, Serbia;

²University Clinical Center of Serbia, Clinic of Pulmonology, Belgrade, Serbia;

³University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia;

⁴Institute for Pulmonary Diseases of Vojvodina, Clinic for Pulmonary Oncology, Novi Sad – Sremska Kamenica, Serbia;

⁵Oncology Institute of Vojvodina, Clinic for Radiation Oncology, Novi Sad – Sremska Kamenica, Serbia;

⁶University of Defence, Medical Faculty of the Military Medical Academy, Belgrade, Serbia;

⁷Military Medical Academy, Pulmonology Clinic, Belgrade, Serbia;

⁸University Business Academy in Novi Sad, Faculty of Pharmacy, Novi Sad, Serbia;

⁹University of Niš, Faculty of Medicine, Niš, Serbia;

¹⁰University Clinical Center of Niš, Oncology Clinic, Niš, Serbia

SUMMARY

Long-term statistical data worldwide on lung cancer (LC) show an overall 34% reduction in mortality compared to 1991. The primary reasons for this decline include a reduced smoking rate, earlier diagnosis, advancements in invasive diagnostic methods, and the introduction of low-dose computed tomography screening. These factors have contributed to detecting LC at earlier stages of the disease and improving timely treatment. The diagnostic sensitivity of conventional bronchoscopy for peripheral pulmonary lesions (PPL), representing early-stage LC, has historically been relatively low, ranging 30–60%. Over the past two decades, diagnostic sensitivity for PPL has improved with the development of advanced navigational techniques, such as virtual bronchoscopic navigation, electromagnetic navigation bronchoscopy, radial endobronchial ultrasound, cone-beam computed tomography, and ultrathin bronchoscopy. In the past two to three years, robotic-assisted bronchoscopy has further enhanced diagnostic navigation capabilities to their current maximum potential.

Keywords: bronchoscopy; early detection of cancer; lung neoplasms; smoking; video-assisted techniques and procedures

INTRODUCTION

Global data on lung cancer (LC) incidence and mortality have become increasingly refined in recent years and demonstrate significant geographical heterogeneity [1–4]. In the period following the COVID-19 pandemic, studies reported an increase in LC incidence and mortality worldwide, largely due to delayed diagnostic evaluation and detection at more advanced stages, which negatively affected treatment outcomes. Overall cancer incidence in 2020 was 9% lower than in 2019, with the most substantial decrease observed in asymptomatic (*in situ* and localized) disease, attributable to reductions in screening and incidental detection during routine medical visits [5]. However, when long-term LC statistical data are considered, there is a continued decrease in mortality, amounting to 34% compared with 1991 (US data). The main reasons for the reduced LC mortality are the decline in the smoking rate, earlier detection, advances in invasive

diagnostic modalities, and the implementation of low-dose computed tomography (LDCT) screening, all contributing to earlier-stage diagnosis and timely treatment [6]. Obtaining an adequate bronchial or pulmonary tissue sample is crucial not only for establishing a diagnosis of LC but also for determining the molecular and immunologic profile of the lung tumor [7]. When the objective of invasive pulmonary diagnostics is early-stage diagnosis, sampling typically involves small peripheral pulmonary lesions (PPL), where flexible bronchoscopy has replaced rigid bronchoscopy completely. New image-guided bronchoscopic techniques have increased diagnostic yield and reduced complication rates compared with conventional bronchoscopy. These include virtual bronchoscopic navigation (VBN), electromagnetic navigation bronchoscopy (ENB), radial endobronchial ultrasound (R-EBUS), cone-beam computed tomography (CBCT), ultrathin bronchoscopy (UTB), and robotic bronchoscopy [8]. Advanced bronchoscopic imaging techniques

Received • Примљено:
December 12, 2025

Accepted • Прихваћено:
January 19, 2026

Online first: January 21, 2026

Correspondence to:

Marko BOJOVIĆ
University of Novi Sad
Faculty of Medicine
Department of Oncology
Hajduk Veljkova 3
21000Novi Sad
Serbia
marko.bojovic@mf.uns.ac.rs



Figure 1. Radial endobronchial ultrasound [16]

such as autofluorescence imaging and narrow-band imaging can also detect changes in the bronchial epithelium, including carcinoma *in situ*, 40% of which may progress to invasive carcinoma. Early detection allows these lesions to be treated endoscopically, for example, by endobronchial brachytherapy or photodynamic therapy [9].

Transthoracic needle aspiration and biopsy (TTNA/TTNB)

TTNA/TTNB are well-established and safe diagnostic techniques for obtaining cytologic or histologic samples from PPL. Over time, several imaging modalities have been used to guide needle placement, including plain radiography, fluoroscopy, computed tomography (CT), ultrasound, and electromagnetic-navigated TTNA [10]. These TTNA/TTNB methods have become procedures of choice for diagnosing peripheral pulmonary nodules due to their high diagnostic yield. A sample can be obtained via needle biopsy or fine needle aspiration for cytology and cell-block preparation, and both sample types are suitable for molecular analysis. The CT-guided TTNA and TTNB demonstrate diagnostic yields ranging from 64% to 97% [11]. The most common complications include pneumothorax (rate to ~25%, with ~5–6% requiring intervention) and hemoptysis (5% prevalence) [12]. Thoracic oncologists generally avoid TTNA/TTNB for PPL because these lesions are potentially resectable. However, this approach is preferred for suspected small cell lung carcinoma, for which surgery is not indicated. When surgery is contraindicated for any reason, TTNA/TTNB method is strongly recommended.

Bronchoscopy – new diagnostic procedures

Interventional pulmonology has expanded rapidly since the first groundbreaking studies in 2004 which demonstrated the usefulness of R/L-EBUS for diagnosing pulmonary lesions and LC staging. The development of interventional pulmonology has significantly improved the management of LC, particularly early-stage non-small cell LC [12, 13].

R-EBUS with and without guide sheath (GS)

Although conventional flexible bronchoscopy can be used for diagnosing PPL, its diagnostic yield is variable and generally lower than that of CT-guided TTNA/TTNB. This limitation facilitated the development of newer bronchoscopic techniques, such as R/L-EBUS [14]. Several studies have confirmed that a multimodal approach that combines R-EBUS with VBN or ENB has an improved diagnostic yield when compared to using only R-EBUS, as navigational systems compensate for instances in which R-EBUS fails to reach the lesion [15]. The key limitation of R-EBUS is the lack of real-time tissue sampling, as the radial probe (RP) must be removed from the bronchoscope's working channel before biopsy instruments are introduced. Maintaining a consistent position of the RP withdrawal and instrument insertion is challenging (Figure 1) [16]. GS use can partly help mitigate this issue. Diagnostic sensitivity of R-EBUS for PPL, especially for LC diagnosis, depends on lesion size, location, presence of a bronchus sign, lesion type, and the availability of rapid on-site evaluation [17]. Diagnostic sensitivity is reduced for upper lobe lesions because the RP cannot navigate sharply angled segments. The diagnostic sensitivity is greatly influenced by




Figure 2. Electromagnetic navigation bronchoscopy [23]

the position of PPL when compared to RP, with the highest diagnostic sensitivity when the R-EBUS probe is centrally positioned within the lesion (82.6%), lower when in the adjacent position (56.8%), and the lowest when the probe is outside the lesion (17.3%) (24). A meta-analysis of 46 studies (2002–2022) included 7252 patients with PPL, out of which 5173 patients were successfully diagnosed by R-EBUS with an overall diagnostic sensitivity of 73.4% (95% CI: 69.9–76.7%) [18].

UTB in the diagnosis of PPL

UTB uses bronchoscopes with a working channel diameter of 1.8–2.2 mm, enabling navigation into smaller bronchi and facilitating access and biopsy of PPL. When combined with R-EBUS, UTB demonstrates better diagnostic sensitivity compared with thin bronchoscopy (TB), particularly for lesions in the upper segments of the lower lobes and the upper lobes [19, 20].

Multiple studies have shown higher sensitivity with UTB compared to TB (70.1% vs. 58.7%). In a study by Nishii et al. [21], patients first underwent TB with R-EBUS; if the probe did not enter the lesion, TB was replaced with UTB. Positive or negative bronchus sign presence or absence had an impact on statistically significant difference between diagnostic and nondiagnostic bronchoscopies, and both procedures were performed with VBN assistance [21].

New navigational techniques in bronchoscopy of PPL

ENB is a type of bronchoscopic navigation that enables guided sampling of PPL using electromagnetic field-based reconstruction of a three-dimensional pathway (3D) to the target lesion [22]. During the procedure, the bronchoscope

is navigated through the airways using real-time electromagnetic navigation. Virtual 3D mapping laid over the live bronchoscopic imaging allows the operator to lead the bronchoscope with precision to the lesion. Biopsy instruments are then introduced into the working channel to obtain diagnostic biopsy samples (Figure 2) [23].

In the multicenter NAVIGATE study, ENB achieved a diagnostic yield of 73%, with the R-EBUS-assisted group demonstrating superior sensitivity compared with the fluoroscopy group [24].

CBCT in bronchoscopy

CBCT provides real-time airway imaging and confirms navigational accuracy through specialized software. This enables the bronchologist to localize with PPL that are difficult or impossible to reach using conventional bronchoscopy and would require fluoroscopy to guide the instruments into the lesion [25]. The use of radiation or fluoroscopy is highly reduced. Limitations of the procedure are the following: the need for specialized equipment and infrastructure, including a complete CBCT system. The interpretation of CBCT also requires advanced operator training, and there are also significant system and maintenance costs. All this makes CBCT less accessible for institutions with limited resources [25, 26].

Bhadra et al. [27] published a study of 200 patients who underwent CBCT bronchoscopy with a multimodal approach which incorporated both conventional and UTB, and with a smaller group of patients also underwent cryobiopsy. The diagnostic sensitivity was 90%, with 60% malign lung lesions, 30% benign lung lesions and 10% undiagnosed. By using the cryoprobe, the authors increased the diagnostic sensitivity from 86.4% to 90.1% [27].



Figure 3. Application of robotic assisted bronchoscopy; the bronchologist guides the procedure via the display; on the left in the image of the monitor is a real-time bronchoscopic procedure, the right side shows a virtual bronchoscopic view [28]

Robotic-assisted bronchoscopy (RAB)

RAB enables the bronchologist to visualize and access previously unreachable peripheral lung regions. Although system cost remains a significant barrier for RAB (Figure 3), it currently offers the highest diagnostic yield for PPL [28]. This method integrates three navigational modalities – electromagnetic guidance, optical pattern recognition, and robotic kinematic feedback – providing highly accurate localization on high-resolution monitors. Initially, target lesions are mapped on CT, imported into planning software, and then via robotic platform the target lesions are accessed with precision by guiding the bronchoscope through the airways [29].

The robotic platform includes an innovative telescoping endoscope mounted on flexible robotic arms, enabling superior maneuverability, reach, and stability. Combining advanced imaging, improved biopsy tools, and robotic precision has enabled diagnostic sensitivity for small PPL to exceed 90% [30]. These results are comparable to CT-guided TTNA/TTNB, which has proven sensibility of 84–96%. However, RAB has a better safety profile, with the 2.3% pneumothorax rate and 0.6% bleeding risk [31]. Current evidence suggests that, with appropriate expertise

and technology, bronchoscopy-based biopsy of PPL can be recommended as preferable method with minimal selection bias.

CONCLUSION

Bronchoscopy has undergone profound transformation since its inception, evolving from a simple tool for removing foreign bodies into a sophisticated diagnostic and therapeutic platform. Modern navigational technologies have markedly improved the safety and accuracy of diagnosing PPL. RAB represents the most advanced development to date. As these technologies continue to evolve, they hold promise not only for improving peripheral lesion diagnosis but also for enabling future bronchoscopic therapeutic interventions in peripheral lung malignancies.

Ethics: The authors declare that the article was written in accordance with the ethical standards of the Serbian Archives of Medicine as well as the ethical standards of medical facilities for each author involved.

Conflict of interest: None declared.

REFERENCES

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J Clin.* 2022;72(1):7–33. [DOI: 10.3322/caac.21708] [PMID: 35020204]
2. Conic I, Nedovic B, Zivadinovic R, Zivadinovic R, Petric A, Stojnev S, et al. Analysis of progression-free and overall survival in ovarian cancer: Bevacizumab treatment outcomes using historical cohort. *Pharmazie.* 2024;79(9):195–201. [DOI: 10.1691/ph.2024.4570] [PMID: 39407422]
3. Popevic S, Maric N, Ilic B, Belic S, Sekulovic Radovanovic I, Dimic-Janjic S, et al. Hyperglycemia and Lung Cancer-A Possible Relationship. *Diagnostics (Basel)* 2025;15(6):651. [DOI: 10.3390/diagnostics15060651] [PMID: 40149994]
4. Osarogiagbon RU, Liao W, Faris NR, Fehnel C, Goss J, Shepherd CJ, et al. Evaluation of Lung Cancer Risk Among Persons Undergoing Screening or Guideline-Concordant Monitoring of Lung Nodules in the Mississippi Delta. *JAMA Netw Open.* 2023;6(2):e230787. [DOI: 10.1001/jamanetworkopen.2023.0787] [PMID: 36848089]

5. Negoita S, Chen H-S, Sanchez PV, Sherman RL, Henley SJ, Siegel RL, et al. Annual Report to the Nation on the Status of Cancer, part 2: Early assessment of the COVID-19 pandemic's impact on cancer diagnosis. *Cancer*. 2024;130(1):117–27. [DOI: 10.1002/cncr.35026] [PMID: 37755665]
6. Siegel RL, Kratzer TB, Giaquinto AN, Sung H, Jemal A. Cancer statistics, 2025. *CA Cancer J Clin*. 2025;75(1):10–45. [DOI: 10.3322/caac.21871] [PMID: 39817679]
7. Jovanović D, Roksandić-Milenković M, Kotur-Stevuljević J, Čeriman V, Vukanić I, Samardžić N, et al. Soluble sPD-L1 and Serum Amyloid A1 as Potential Biomarkers for Lung Cancer. *J Med Biochem*. 2019;38(3):332–41. [DOI: 10.2478/jomb-2018-0036] [PMID: 31156344]
8. Pinsky P, Miller E, Faris N, Osarogiagbon R. Pulmonary Nodules, Lung Cancer Screening, and Lung Cancer in the Medicare Population. *Chest*. 2023;163(5):1304–13. [DOI: 10.1016/j.chest.2022.12.006] [PMID: 36529155]
9. Lalić N, Bojović M, Ivanov O, Ličina J, Popević S, Stjepanović M, et al. Effective Differences between 2D and 3D Planned Brachytherapy in Lung Cancer: An Institutional Retrospective Study. *Medicina (Kaunas)*. 2024;60(3):452. [DOI: 10.3390/medicina60030452] [PMID: 38541177]
10. Xu C, Yuan Q, Chi C, Zhang Q, Wang Y, Wang W, et al. Computed tomography-guided percutaneous transthoracic needle biopsy for solitary pulmonary nodules in diameter less than 20 mm. *Medicine (Baltimore)*. 2018;97(14):e0154. [DOI: 10.1097/MD.00000000000010154] [PMID: 29620630]
11. Guzinski M, Puła M, Zdanowicz A, Kacała A, Dudek K, Lipiński A, et al. Safety, feasibility, and effectiveness of a CT-guided transthoracic lung and pleural biopsy - a single-centre experience with own low-dose protocol. *Pol J Radiol*. 2023;88:e546–51. [DOI: 10.5114/pjr.2023.133309] [PMID: 38125814]
12. Lalić N, Lovrenski A, Ilić M, Ivanov O, Bojović M, Lalić I, et al. Invasive Diagnostic Procedures from Bronchoscopy to Surgical Biopsy-Optimization of Non-Small Cell Lung Cancer Samples for Molecular Testing. *Medicina (Kaunas)*. 2023;59(10):1723. [DOI: 10.3390/medicina59101723] [PMID: 37893442]
13. Lalic N, Bursac D, Bojovic M, Nemet M, Ergelasev I. The importance of re-biopsy in the era of molecular therapy for lung cancer. *Srp Arh Celok Lek*. 2024;152(3–4):209–13. [DOI: 10.2298/SARH231114015L]
14. Leong TL, Steinfort DP. Contemporary Concise Review 2023: Advances in lung cancer and interventional pulmonology. *Respirology*. 2024;29(8):665–73. [DOI: 10.1111/resp.14789] [PMID: 38960450]
15. Bae S, Lim S, Ahn JJ, Jegal Y, Seo KW, Ra SW, et al. Diagnosing peripheral lung lesions using endobronchial ultrasonography with guide sheath: A prospective registry study to assess the effect of virtual bronchoscopic navigation using a computed tomography workstation. *Medicine (Baltimore)*. 2020;99(17):e19870. [DOI: 10.1097/MD.00000000000019870] [PMID: 32332652]
16. Zarogoulidis P, Huang H, Chen W, Petridis D, Matthaios D, Hohenforst-Schmidt W, et al. Radial Endobronchial Ultrasound for Lung Cancer Diagnosis: Tips and Tricks. *J Cancer*. 2022;13(4):1307–12. [DOI: 10.7150/jca.67113] [PMID: 35281877]
17. Boonsarngsuk V, Petnak T, So-Ngern A, Saksitthichok B, Kanoksil W. Comparison of different transbronchial biopsy sampling techniques for the diagnosis of peripheral pulmonary lesions with radial endobronchial ultrasound-guided bronchoscopy: A prospective study. *Respir Invest*. 2020;58(5):381–6. [DOI: 10.1016/j.resinv.2020.03.004] [PMID: 32284223]
18. Tian S, Li X, Liu J, Wang X, Chen H, Dai Z, et al. Radial endobronchial ultrasound – guided bronchoscopy for the diagnosis of peripheral pulmonary lesions: A systematic review and meta-analysis of prospective trials. *Heliyon*. 2024;10(8):e29446. [DOI: 10.1016/j.heliyon.2024.e29446] [PMID: 38660275]
19. Nguyen P, Shah PL. Contemporary Concise Review 2024: New Techniques in Interventional Pulmonology. *Respirology*. 2025;30(8):697–704. [DOI: 10.1111/resp.70084] [PMID: 40629539]
20. Oki M, Saka H, Asano F, Kitagawa C, Kogure Y, Tsuzuku A, et al. Use of an Ultrathin vs Thin Bronchoscope for Peripheral Pulmonary Lesions: A Randomized Trial. *Chest*. 2019;156(5):954–64. [DOI: 10.1016/j.chest.2019.06.038] [PMID: 31356810]
21. Nishii Y, Nakamura Y, Fujiwara K, Ito K, Sakaguchi T, Suzuki Y, et al. Use of Ultrathin Bronchoscope on a Need Basis Improves Diagnostic Yield of Difficult-to-Approach Pulmonary Lesions. *Front Med (Lausanne)*. 2020;7:588048. [DOI: 10.3389/fmed.2020.588048] [PMID: 33385003]
22. Fu Y, Yuan X, Pan W, Lu Y, Zeng D, Jiang J. Comparative analysis of the efficacy and safety of electromagnetic navigation bronchoscopy combined with x-ray or radial endobronchial ultrasound biopsy in the diagnosis of small peripheral pulmonary nodules. *Thorac Cancer*. 2023;14(15):1348–54. [DOI: 10.1111/1759-7714.14879] [PMID: 37037512]
23. Yutaka Y, Sato T, Isowa M, Murata Y, Tanaka S, Yamada Y, et al. Electromagnetic navigation bronchoscopy versus virtual bronchoscopy navigation for improving the diagnosis of peripheral lung lesions: analysis of the predictors of successful diagnosis. *Surg Today*. 2022;52(6):923–30. [DOI: 10.1007/s00595-021-02398-z] [PMID: 34705111]
24. Folch EE, Pritchett MA, Nead MA, Bowling MR, Murgu SD, Krimsky WS, et al. Electromagnetic Navigation Bronchoscopy for Peripheral Pulmonary Lesions: One-Year Results of the Prospective, Multicenter NAVIGATE Study. *J Thorac Oncol*. 2019;14(3):445–58. [DOI: 10.1016/j.jtho.2018.11.013] [PMID: 30476574]
25. Aboudara M, Katsis J. Cone-Beam CT Scan-Guided Peripheral Bronchoscopy. *CHEST Pulmonary*. 2024;2(2):100045. [DOI: 10.1016/j.chp.2024.100045]
26. Popević S, Ilić B, Grujić M, Zečević A, Uskoković Stefanović Ž, Belić S, et al. Hemoptizije. In: Selected chapters from pulmonology. Beograd: Medical Faculty in Belgrade; 2023.
27. Bhadra K, Setser RM, Condra W, Bader BA, David S. A Cone Beam CT Bronchoscopy Study of the Ultrathin Cryoprobe for Biopsy of Peripheral Lung Lesions. *J Bronchology Interv Pulmonol*. 2024;31(2):117–25. [DOI: 10.1097/LBR.0000000000000936] [PMID: 37450607]
28. Chan JWY, Chang ATC, Yu PSY, Lau RWH, Ng CSH. Robotic Assisted-Bronchoscopy With Cone-Beam CT ICG Dye Marking for Lung Nodule Localization: Experience Beyond USA. *Front Surg*. 2022;9:943531. [DOI: 10.3389/fsurg.2022.943531] [PMID: 35836599]
29. Wijma IN, Casal RF, Cheng GZ, Einsiedel PF, Fantin A, Hall DJ, et al. Radiation Principles, Protection, and Reporting for Interventional Pulmonology: A World Association of Bronchology and Interventional Pulmonology White Paper. *Respiration*. 2024;103(11):707–22. [DOI: 10.1159/000540102] [PMID: 39033746]
30. Chambers J, Knox D, Leclair T. O-arm CT for Confirmation of Successful Navigation During Robotic Assisted Bronchoscopy. *J Bronchology Interv Pulmonol*. 2023;30(2):155–62. [DOI: 10.1097/LBR.0000000000000894] [PMID: 36066326]
31. Ali MS, Ghori UK, Wayne MT, Shostak E, De Cardenas J. Diagnostic Performance and Safety Profile of Robotic-assisted Bronchoscopy: A Systematic Review and Meta-Analysis. *Ann Am Thorac Soc*. 2023;20(12):1801–12. [DOI: 10.1513/AnnalsATS.202301-075OC] [PMID: 37769170]

Инвазивне дијагностичке процедуре за рак плућа у раном стадијуму – клинички значај нових навигационих техника у интервентној бронхоскопији

Спасоје Попевић^{1,2}, Ненси Лалић^{3,4}, Марко Бојовић^{3,5}, Раде Милић^{6,7}, Ивица Лалић⁸, Бранислав Илић^{1,2}, Ивана Секуловић-Радовановић², Дане Кртинић^{9,10}

¹Универзитет у Београду, Медицински факултет, Београд, Србија;

²Универзитетски клинички центар Србије, Клиника за пулмологију, Београд, Србија;

³Универзитет у Новом Саду, Медицински факултет, Нови Сад, Србија;

⁴Институт за плућне болести Војводине, Клиника за плућну онкологију, Нови Сад – Сремска Каменица, Србија;

⁵Институт за онкологију Војводине, Клиника за радијацијску онкологију, Нови Сад – Сремска Каменица, Србија;

⁶Универзитет одбране, Медицински факултет Војномедицинске академије, Београд, Србија;

⁷Војномедицинска академија, Клиника за пулмологију, Београд, Србија;

⁸Универзитет Привредна академија у Новом Саду, Фармацеутски факултет, Нови Сад, Србија;

⁹Универзитет у Нишу, Медицински факултет, Ниш, Србија;

¹⁰Универзитетски клинички центар Ниш, Клиника за онкологију, Ниш, Србија

САЖЕТАК

Дугорочни статистички подаци о раку плућа показују укупно смањење морталитета за 34% у поређењу са 1991. годином. Главни разлози за овај пад укључују смањену стопу пушења, ранију дијагнозу, напредак у инвазивним дијагностичким методама и увођење скрининга нискодозном компјутеризованом томографијом. Ови фактори су допринели откривању рака плућа у ранијим стадијумима болести и побољшању благовременог лечења. Дијагностичка осетљивост конвенционалне бронхоскопије за периферне плућне лезије, које представљају рани стадијум болести, била је релативно ниска, између 30% и 60%. Током протекле две

деценије, дијагностичка осетљивост за периферне плућне лезије побољшала се развојем напредних навигационих техника, као што су виртуелна бронхоскопска навигација, електромагнетна навигациона бронхоскопија, радијални ендобронхијални ултразвук, конусна компјутеризована томографија и ултратанка бронхоскопија. У протекле две до три године, роботски потпомогнута бронхоскопија додатно је побољшала могућности дијагностичке навигације до њиховог тренутног максималног потенцијала.

Кључне речи: бронхоскопија; рано откривање карцинома; неоплазме плућа; пушење; видеоасистирани технике и процедуре