

ORIGINAL ARTICLE / ОРИГИНАЛНИ РАД

Optical coherence tomography angiography microvascular changes in the macular region in open-angle glaucoma

Rada Zečević¹, Sunčica Srećković^{2,3}, Marko Kontić¹¹Belgrade Ophthalmology Center Special Ophthalmology Hospital, Belgrade, Serbia;²Kragujevac University Clinical Center, Clinic of Ophthalmology, Kragujevac, Serbia;³University of Kragujevac, Faculty of Medical Sciences, Department of Ophthalmology, Kragujevac, Serbia

SUMMARY

Introduction/Objective Glaucoma is the leading cause of irreversible blindness worldwide. It is hypothesized that glaucomatous microvascular changes in the superficial capillary plexus (SCP) may lead to early structural damage and correlate with visual field defects. The aim of this study was to quantitatively analyze microvascular characteristics of the SCP in the macular region of healthy eyes and in eyes with varying severity of open-angle glaucoma (OAG) using optical coherence tomography angiography (OCT-A).

Methods A total of 144 eyes were included in this cross-sectional study, including 109 eyes with confirmed OAG and 35 healthy eyes. Based on the Hodapp–Anderson–Parrish classification, patients were categorized into early, moderate, and severe stage glaucoma. All subjects underwent visual field, optical coherence tomography, and OCT-A examinations.

Results Significant reductions were observed in macular vessel density across glaucoma stages. Foveal vessel density decreased from $23 \pm 1.9\%$ in normal eyes to $13.9 \pm 1.8\%$ in severe glaucoma eyes. Foveal and parafoveal vessel densities were significantly reduced even in early-stage glaucoma. OCT parameters progressively decreased with glaucoma severity. Total ganglion cell complex thickness decreased from $108.5 \pm 5.6 \mu\text{m}$ in healthy eyes to $61.2 \pm 6.9 \mu\text{m}$ in severe glaucoma eyes. Total peripapillary retinal nerve fiber layer thickness decreased from $106.2 \pm 7.9 \mu\text{m}$ in healthy eyes to $48.7 \pm 7.5 \mu\text{m}$ in severe glaucoma.

Conclusion Our findings support the hypothesis that macular vessel density decreases progressively with glaucoma severity. These results reinforce the potential clinical utility of OCT-A in detecting early glaucoma and monitoring disease progression.

Keywords: glaucoma; vascular density; intraocular pressure; optical coherence tomography

INTRODUCTION

Glaucoma is the leading cause of irreversible blindness worldwide. It is a multifactorial optic neuropathy characterized by progressive loss of retinal ganglion cells (RGCs) and their axons in both the macular and peripapillary regions, with characteristic structural changes in the optic nerve head (ONH) and corresponding functional visual field defects [1]. Among the many risk factors for glaucoma, intraocular pressure (IOP) is the only modifiable one. It is estimated that in 2020, approximately 76 million people were affected by glaucoma, with projections indicating a 74% increase by 2040 [2].

The primary goal of glaucoma treatment is to preserve patients' visual function and quality of life at an acceptable cost [1]. Lowering IOP is currently the only proven treatment to slow disease progression [3]. However, some patients continue to experience deterioration despite achieving target IOP, suggesting that optic nerve ischemia and reduced ocular blood flow contribute to glaucoma pathogenesis and progression [4, 5].

Optical coherence tomography angiography (OCT-A) is a novel, non-invasive imaging modality that provides detailed visualization of

the retinochoroidal microvasculature, including blood flow around the ONH and macula, without the need for contrast dyes [6]. Because RGCs are predominantly supplied by the superficial macular vascular complex, it is hypothesized that glaucomatous microvascular changes in this layer may lead to early structural damage and correlate with visual field defects [7, 8]. Therefore, the macular region is considered a strategic location for detecting and monitoring glaucoma progression.

The aim of this study was to quantitatively analyze microvascular characteristics of the superficial capillary plexus (SCP) in the macular region of healthy eyes and in eyes with varying severity of open-angle glaucoma (OAG). Additionally, we sought to investigate correlations between macular vessel density and traditional structural and functional parameters of glaucoma.

Received • Примљено:

July 22, 2025

Revised • Ревизија:

January 25, 2026

Accepted • Прихваћено:

January 30, 2026

Online first: February 4, 2026

Correspondence to:

Rada ZEČEVIĆ
Belgrade Ophthalmology Center
Special Ophthalmology Hospital
27. marta 17–19
11000 Belgrade, Serbia
radiculjak_2901@yahoo.com

METHODS

Study design

Subjects

A total of 109 eyes from patients aged ≥ 40 years with confirmed OAG in at least one eye were included. The control group comprised 35 healthy eyes from individuals over 40 years of age.

All participants underwent a comprehensive ophthalmic examination, including: best-corrected visual acuity (BCVA) using the Snellen chart, slit-lamp biomicroscopy, IOP measurement with Goldmann applanation tonometry, central corneal thickness (CCT) measurement (Topcon Aladdin Optical Biometer, Oakland, CA, USA), gonioscopy using a Goldmann contact lens, dilated fundus examination, standard automated perimetry using Octopus 600 (Haag-Streit, Mason, OH, USA), OCT imaging of the ONH and macula, including retinal nerve fiber layer (RNFL) and ganglion cell complex (GCC) thickness measurements, and macular vessel density assessment via OCT-A (Topcon Maestro 2, IMAGENet6 software, Oakland, CA, USA).

Inclusion criteria

The inclusion criteria were age ≥ 40 years, a confirmed diagnosis of OAG for at least one year, and an open anterior chamber angle on gonioscopy.

The control group included eyes with IOP ≤ 21 mmHg, no history of elevated IOP, normal optic nerve appearance with intact neuroretinal rims and RNFL, and a normal visual field with no defects.

Exclusion criteria

Exclusion criteria for both healthy and glaucoma groups included: BCVA < 0.2 , spherical refraction ≥ 5 D or myopia > 4 D, pregnancy or breastfeeding, other types of glaucoma (angle-closure), age-related macular degeneration or other inherited/acquired macular diseases, uveitis, diabetic or hypertensive retinopathy/maculopathy, corneal diseases, history of ocular trauma or previous ocular surgeries (except cataract and glaucoma surgery), non-glaucomatous optic neuropathy, use of medications affecting retinal function (tamoxifen, antimalarials, phenothiazines, canthaxanthin, methoxyflurane), neurological disorders (Alzheimer's, Parkinson's, dementia, stroke).

Glaucoma classification

Based on the Hodapp–Anderson–Parrish classification, patients were categorized as follows:

- Early-stage glaucoma: Mean deviation (MD) > -6 dB;
- Moderate-stage glaucoma: MD between -6 and -12 dB;
- Severe-stage glaucoma: MD < -12 dB;
- Control group: Healthy eyes.

OCT angiography and structural measurements

OCT-A Measurements

OCT-A scans were performed using the Topcon Maestro 2 (IMAGENet6 software, Oakland, CA, USA). This device uses an 840 nm light source with an A-scan rate of 50,000 scans per second, providing high-resolution 3D visualization of retinal vasculature.

Superficial vessel density (SVD) was automatically calculated as the percentage of the scanned area occupied by blood vessels. Measurements focused on the SCP, spanning from the internal limiting membrane (ILM) to the inner plexiform layer (IPL).

A 3×3 mm² field centered on the fovea was analyzed, with density assessed in the following:

- Foveal zone (1 mm diameter);
- Parafoveal zones (1–3 mm): superior, inferior, nasal, and temporal sectors.

OCT and OCT-A image quality review was completed according to the Imaging Data Evaluation and Analysis Reading Center protocol on all scans using standard Topcon Maestro 2 software (IMAGENet6 software). Poor-quality scans were excluded from the study and analysis if one of the following criteria was met: signal strength index < 45 , poor clarity, residual motion artifacts, image cropping or local weak signal due to vitreous floaters, segmentation errors, or images with off-center fovea.

OCT structural measurements

All participants underwent ONH and macular imaging, with measurements including:

- ONH: Disc area, rim area, cup-to-disc (C/D) ratios, and RNFL thickness
- GCC: Thickness from ILM to inner nuclear layer, including nerve fiber, ganglion cell, and IPL layers

Statistical analysis

Results are presented as mean \pm standard deviation, median (25th–75th) and number (%). Graphical and mathematical methods were used to examine data distribution. The groups were compared using the χ^2 test. Data were analyzed using one-way ANOVA or the Kruskal–Wallis test when three groups were compared. For significant one-way ANOVA results, Tukey post hoc analysis was used. Afterward, for the Kruskal–Wallis test, the Mann–Whitney test was used as a post hoc test. All $p < 0.05$ values were considered statistically significant. Statistical analysis was performed using IBM SPSS Statistics, Version 26 (IBM Corp., Armonk, NY, USA).

Ethics: This cross-sectional study was approved by the Research Ethics Committee of the Belgrade Ophthalmology Center Special Eye Hospital on November 9, 2022 (approval number: 3/30/2022). The study was conducted in accordance with the tenets of the Declaration of Helsinki. Informed consent was obtained from all participants.

Table 1. Demographic and clinical characteristics of normal, early, moderate, and severe open-angle glaucoma eyes

| Parameter | Normal n = 35 | Early n = 44 | Moderate n = 34 | Severe n = 31 | p |
|----------------------|-------------------------|-------------------------|-------------------------|------------------------|---------|
| Age (y) | 55.8 ± 8.8 | 61 ± 10.4 | 64.4 ± 12.1 | 67.1 ± 10.7 | < 0.001 |
| Sex (female/male) | 20 (57.1%) / 15 (42.9%) | 27 (61.4%) / 17 (38.6%) | 19 (55.9%) / 15 (44.1%) | 7 (22.6%) / 24 (77.4%) | 0.005 |
| BCVA (Snellen) | 0.9 ± 0.1 | 0.9 ± 0.2 | 0.8 ± 0.3 | 0.8 ± 0.2 | 0.003 |
| CCT (µm) | 562.9 ± 17.7 | 552.7 ± 31.1 | 546.6 ± 24.4 | 541.9 ± 26 | 0.007 |
| Linear C/D ratio | 0.4 ± 0.1 | 0.8 ± 0.1 | 0.9 ± 0.1 | 0.9 ± 0 | < 0.001 |
| Vertical C/D ratio | 0.4 ± 0.1 | 0.8 ± 0.1 | 0.9 ± 0.1 | 1 ± 0 | < 0.001 |
| IOP (mmHg) | 16.4 ± 2.2 | 17.1 ± 2.8 | 16.5 ± 2.7 | 16.5 ± 3.2 | 0.68 |
| MD visual field (dB) | 0.5 ± 0.6 | -2.5 ± 1.8 | -8.8 ± 1.9 | -18.4 ± 3.6 | < 0.001 |

BCVA – best corrected visual acuity; CCT – central corneal thickness; IOP – intraocular pressure; MD – mean deviation

Table 2. Optical coherence tomography structural characteristics of normal, early, moderate, and severe open-angle glaucoma eyes

| Parameter | Normal n = 35 | Early n = 44 | Moderate n = 34 | Severe n = 31 | P |
|--------------------------------|---------------|--------------|-----------------|---------------|---------|
| Total GCC thickness (µm) | 108.5 ± 5.6 | 89.6 ± 13.6 | 77.6 ± 6.6 | 61.2 ± 6.9 | < 0.001 |
| Superior GCC thickness (µm) | 107.1 ± 4.9 | 90.9 ± 12.8 | 79.2 ± 11.4 | 62.5 ± 7.5 | < 0.001 |
| Inferior GCC thickness (µm) | 110 ± 6.9 | 88 ± 16.3 | 76.1 ± 9.9 | 60.6 ± 6.5 | < 0.001 |
| Total pRNFL thickness (µm) | 106.2 ± 7.9 | 84 ± 16.4 | 63.1 ± 10.9 | 48.7 ± 7.5 | < 0.001 |
| Superior pRNFL thickness (µm) | 126.1 ± 13.2 | 98.1 ± 22.2 | 71.1 ± 19.8 | 55.4 ± 12.7 | < 0.001 |
| Inferior pRNFL thickness (µm) | 139 ± 13.6 | 100.8 ± 26.1 | 65.9 ± 13.1 | 50.9 ± 6.6 | < 0.001 |
| Average Macular thickness (µm) | 279.7 ± 11.1 | 266.1 ± 15.4 | 255.3 ± 14.1 | 237.3 ± 14.7 | < 0.001 |

GCC – ganglion cell complex; pRNFL – peripapillary retinal nerve fiber layer

RESULTS

A total of 35 healthy eyes, 44 early-stage glaucoma eyes, 34 moderate-stage glaucoma eyes, and 31 severe-stage glaucoma eyes met the inclusion criteria.

Demographic and clinical characteristics of normal, early, moderate, and severe OAG eyes are presented in Table 1. The mean age increased progressively from normal eyes (55.8 years) to severe glaucoma eyes (67.1 years). The proportion of males increased significantly with disease severity ($p = 0.005$), with the severe glaucoma group being predominantly male (77.4%). BCVA declined with increasing severity of glaucoma ($p = 0.003$). CCT decreased with increasing glaucoma severity ($p = 0.007$). The linear C/D ratio increased with disease severity ($p < 0.001$). Post hoc tests showed significant differences between all severity levels ($p < 0.001$). Similar to the linear C/D ratio, the vertical C/D ratio significantly increased with severity ($p < 0.001$), with all pairwise comparisons being statistically significant ($p < 0.001$). No statistically significant difference in IOP was observed between the groups ($p = 0.68$). MD worsened with increasing glaucoma severity ($p < 0.001$). Post hoc tests showed significant differences between all groups ($p < 0.001$), confirming progressive visual field loss across stages.

The data on structural characteristics of normal, early, moderate, and severe OAG eyes are presented in Table 2. Total GCC thickness progressively decreased with increasing glaucoma severity, from 108.5 ± 5.6 µm in healthy eyes to 61.2 ± 6.9 µm in severe glaucoma eyes ($p < 0.001$). Tukey HSD tests revealed significant differences between normal and moderate glaucoma (mean difference: -30.9 µm, $p < 0.001$) and between normal and severe glaucoma (mean difference: -47.3 µm, $p < 0.001$). Both superior and

inferior GCC thicknesses decreased significantly across all stages of glaucoma ($p < 0.001$). Pairwise comparisons showed significant differences between normal and early glaucoma, as well as between early and moderate glaucoma (all $p < 0.001$).

Regarding peripapillary RNFL (pRNFL) thickness, a progressive reduction was observed with increasing disease severity ($p < 0.001$). Total pRNFL thickness decreased from 106.2 ± 7.9 µm in healthy eyes to 48.7 ± 7.5 µm in severe glaucoma eyes. Pairwise comparisons showed significant differences between normal and early ($p < 0.001$), early and moderate ($p < 0.001$), and moderate and severe glaucoma groups ($p < 0.001$). Similar trends were observed for superior and inferior pRNFL thickness ($p < 0.001$). The average macular thickness decreased significantly with increasing glaucoma severity ($p < 0.001$), from 279.7 ± 11.1 µm in healthy eyes to 237.3 ± 14.7 µm in severe glaucoma.


The data on superficial macular vessel density in different retinal regions are presented in Table 3 and Figure 1. Foveal vessel density progressively decreased from $23 \pm 1.9\%$ in healthy eyes to $13.9 \pm 1.8\%$ in severe glaucoma ($p < 0.001$). Post hoc analysis showed significant reductions between normal and early glaucoma (mean difference: 4.89% , $p < 0.001$), normal and moderate glaucoma (6.81% , $p < 0.001$), and normal and severe glaucoma (9.13% , $p < 0.001$). Differences between early and moderate glaucoma (1.92% , $p < 0.001$) and early and severe glaucoma (4.24% , $p < 0.001$) were also statistically significant.

Parafoveal vessel density in the superior sector declined significantly from $44.3 \pm 2.5\%$ in healthy eyes to $37.2 \pm 3.5\%$ in severe glaucoma ($p < 0.001$). Parafoveal vessel density in the inferior sector exhibited a similar pattern, decreasing from $45.4 \pm 3.2\%$ in healthy eyes to $36.7 \pm 2.5\%$ in severe glaucoma ($p < 0.001$). Parafoveal

Table 3. Superficial macular vessel density measurements in normal, early, moderate, and severe open-angle glaucoma eyes

| Parameter | Normal | Early | Moderate | Severe | p |
|----------------------------|------------|------------|------------|------------|---------|
| Foveal VD (%) | 23 ± 1.9 | 18.1 ± 1.6 | 16.2 ± 2.2 | 13.9 ± 1.8 | < 0.001 |
| Parafoveal VD Superior (%) | 44.3 ± 2.5 | 40.7 ± 4.2 | 39.6 ± 3.7 | 37.2 ± 3.5 | < 0.001 |
| Parafoveal VD Inferior (%) | 45.4 ± 3.2 | 40.3 ± 5.3 | 37.8 ± 4.3 | 36.7 ± 2.5 | < 0.001 |
| Parafoveal VD Temporal (%) | 44.8 ± 2.8 | 44 ± 4.6 | 42.4 ± 4.5 | 39 ± 2.9 | < 0.001 |
| Parafoveal VD Nasal (%) | 45.1 ± 2.8 | 43.2 ± 3.9 | 42 ± 4.5 | 41.3 ± 3.1 | < 0.001 |

VD – vessel density

**Figure 1.** Superficial macular vessel density measurements in normal, early, moderate, and severe open-angle glaucoma eyes; SCP – superficial capillary plexus

vessel density in the temporal sector was significantly lower in glaucoma eyes, decreasing from $44.8 \pm 2.8\%$ in normal eyes to $39 \pm 2.9\%$ in severe glaucoma ($p < 0.001$). Post hoc comparisons revealed a significant difference between normal and severe glaucoma (5.75% , $p < 0.001$), while differences between early and moderate glaucoma were not statistically significant. Parafoveal vessel density in the nasal sector also declined with disease progression, from $45.1 \pm 2.8\%$ in healthy eyes to $41.3 \pm 3.1\%$ in severe glaucoma ($p < 0.001$). Post hoc analysis between normal and early glaucoma showed significant differences for all

parameters except for temporal and nasal parafoveal vessel density.

The correlations between OCT-A vessel density measurements and structural parameters in the glaucoma group are presented in Table 4. A strong positive correlation was observed between total GCC thickness and macular vessel density measurements in all parafoveal sectors, with the highest correlation found in the foveal region ($r = 0.602$, $p < 0.001$) followed by the temporal sector ($r = 0.557$, $p < 0.001$). Significant correlations were also noted in the superior, inferior, and nasal parafoveal regions ($p < 0.001$; $p < 0.001$; $p = 0.025$). Conversely, linear and vertical C/D ratios exhibited strong negative correlations with vessel density parameters. The strongest negative correlation for the linear C/D ratio was in the foveal region ($r = -0.607$, $p < 0.001$), and for the vertical C/D ratio, the most pronounced correlation was also in the foveal region ($r = -0.579$, $p < 0.001$). Total pRNFL thickness was positively correlated with vessel density, with the strongest correlation observed in the foveal region ($r = 0.536$, $p < 0.001$).

DISCUSSION

Both the mechanical and vascular theories are considered to play a crucial role in the pathogenesis and progression of glaucoma. While mechanical factors, such as increased IOP and ONH deformation, have been extensively studied, the vascular hypothesis remains an area of growing interest. The vascular influence, particularly in the macular region, warrants further investigation, as this area is frequently affected in glaucoma [9]. The irreversible loss of RGCs is believed to be caused, at least in part, by insufficient blood supply. Given their high metabolic demand, RGCs are heavily dependent on the SCP for perfusion. Their unequal distribution within the retina, with the highest concentration in the macular region, suggests that macular vessel density changes may play a role in glaucomatous disease progression [10].

To further explore these vascular changes, we measured and compared macular SVD across different stages of OAG using the Topcon Maestro 2 OCT and OCT-A device. We also assessed the correlations between capillary vessel density and structural parameters.

Our study confirmed that total GCC thickness progressively decreased with increasing glaucoma severity, consistent with findings from the study by Srivastava et al. [11]. Both superior and inferior GCC thickness decreased significantly across glaucoma stages, which aligns with findings from Soares et al.

Table 4. Correlation between optical coherence tomography angiography (OCT-A) vessel density measurements and structural parameters in glaucoma group

| N = 109 | | OCT-A SCP fovea | OCT-A SCP parafovea superior | OCT-A SCP parafovea inferior | OCT-A SCP parafovea temporal | OCT-A SCP parafovea nasal |
|-------------------------------------|---|-----------------|------------------------------|------------------------------|------------------------------|---------------------------|
| GCC thickness total μm | R | 0.602** | 0.457** | 0.435** | 0.557** | 0.214* |
| | P | < 0.001 | < 0.001 | < 0.001 | < 0.001 | 0.025 |
| linear c/d | R | -0.607** | -0.415** | -0.351** | -0.471** | -0.243* |
| | P | < 0.001 | < 0.001 | < 0.001 | < 0.001 | 0.011 |
| vertical c/d | R | -0.579** | -0.347** | -0.330** | -0.424** | -0.224* |
| | P | < 0.001 | < 0.001 | < 0.001 | < 0.001 | 0.019 |
| pRNFL thickness total μm | R | 0.536** | 0.326** | 0.329** | 0.448** | 0.157 |
| | P | < 0.001 | 0.001 | < 0.001 | < 0.001 | 0.102 |

SCP – superficial capillary plexus; GCC – ganglion cell complex;

pRNFL – peripapillary retinal nerve fiber layer;

* $p < 0.05$;** $p < 0.001$

[12]. Moreover, pairwise comparisons confirmed significant differences between normal and early glaucoma, supporting results from Hassanen et al. [13].

These findings reinforce previous reports suggesting that glaucomatous damage primarily affects the inner retinal layers, leading to the thinning of these structures.

Regarding the pRNFL thickness, we found a significant reduction in all glaucoma stages. Similar trends were observed for superior and inferior RNFL thickness, aligning with findings from previous studies [14]. However, in contrast to our results, Bhat et al. [15] reported that inferior pRNFL thinning was the most sensitive marker for glaucoma detection across all disease stages.

Our results, in line with previous research, showed that macular SVD was significantly reduced in glaucoma eyes compared with healthy controls [16]. Specifically, foveal vessel density progressively decreased with glaucoma severity, consistent with the study conducted by Hwang et al. [17].

Parafoveal vessel density declined significantly across all disease stages, with notable reductions in superior, inferior, temporal, and nasal sectors. The nasal parafoveal vessel density also declined with disease progression, although post hoc analysis showed that differences between normal and early glaucoma were not significant for temporal and nasal parafoveal vessel density. This could be explained by the fact that these regions have higher vessel density in healthy eyes [18].

Kuryшева et al. [19] found that SVD was more sensitive than GCC thickness in distinguishing early glaucoma from healthy eyes, supporting the idea that vascular changes may precede structural thinning in some cases.

To further explore the relationship between vascular and structural changes, we analyzed correlations between macular vessel density and retinal structural parameters. Our results showed a strong positive correlation between total GCC thickness and macular vessel density across all parafoveal sectors, with the strongest correlation in the foveal region, followed by the temporal sector. However,

it remains unclear whether RGC loss is primarily driven by vascular insufficiency or whether vessel density reductions are a secondary consequence of structural damage.

Conversely, linear and vertical C/D ratios exhibited strong negative correlations with vessel density parameters, indicating that larger C/D ratios are associated with reduced vessel density. Total pRNFL thickness was positively correlated with vessel density, with the strongest correlation in the foveal region.

This study has several limitations that should be acknowledged. First, all patients were using different anti-glaucoma medications, which could potentially influence ocular blood flow and retinochoroidal vessel measurements. Second, we used a 3×3 mm² scan instead of a 6×6 mm² scan, which might have led to different conclusions. Considering that RGCs are more concentrated in the perifoveal rather than the foveal and parafoveal regions, a larger scan size may yield different results. Finally, this study does not provide longitudinal data on structural and functional changes over time. A longitudinal study would offer a more robust evaluation of the relationship between macular microvascular perfusion and glaucomatous optic neuropathy progression.

CONCLUSION

Our findings support the hypothesis that macular vessel density decreases progressively with glaucoma severity and that vascular and structural parameters are closely correlated. These results reinforce the potential clinical utility of OCT-A in detecting early glaucoma and monitoring disease progression. Future studies should aim to clarify the causal relationship between vascular insufficiency and RGC loss and further explore the longitudinal changes in macular microvascular perfusion in glaucoma.

Conflict of interest: None declared.

REFERENCES

- Pazos M, Traverso CE, Viswanathan A; European Glaucoma Society; Guidelines Task Force; Guidelines Writers, Authors and Contributors. European Glaucoma Society - Terminology and guidelines for glaucoma, 6th Edition. *Br J Ophthalmol.* 2025;109(Suppl 1):1–212. [DOI: 10.1136/bjophthalmol-2025-egsguidelines] [PMID: 41026937]
- Allison K, Patel D, Alabi O. Epidemiology of glaucoma: the past, present, and predictions for the future. *Cureus.* 2020;12(11):e11686. [DOI: 10.7759/cureus.11686] [PMID: 33391921]
- Bengtsson B, Lindén C, Heijl A, Andersson-Geimer S, Aspberg J, Jóhannesson G. The glaucoma intensive treatment study: interim results from an ongoing longitudinal randomized clinical trial. *Acta Ophthalmol.* 2022;100(2):e455–62. [DOI: 10.1111/aos.14978] [PMID: 34327833]
- Dinakaran S, Mehta P, Mehta R, Tilva B, Arora D, Tejwani S. Significance of non-intraocular pressure (IOP)-related factors particularly in normal tension glaucoma: looking beyond IOP. *Indian J Ophthalmol.* 2022;70(2):569–73. [DOI: 10.4103/ijo.IJO_861_21] [PMID: 35086239]
- Prinz J, Fuest M, Kotliar K, Walter P, Hollstein M, Plange N, et al. Vascular reaction parameters correlate with severe nerve fibre thickness loss and visual field defects in glaucoma. *Int Ophthalmol.* 2025;45(1):424. [DOI: 10.1007/s10792-025-03810-0] [PMID: 41117964]
- Hormel TT, Huang D, Jia Y. Advances in OCT angiography. *Transl Vis Sci Technol.* 2025;14(3):6. [DOI: 10.1167/tvst.14.3.6] [PMID: 40052848]
- Yoon J, Lee A, Song WK, Kim KE, Kook MS. Association of superficial macular vessel density with visual field progression in open-angle glaucoma with central visual field damage. *Sci Rep.* 2023;13(1):7190. [DOI: 10.1038/s41598-023-34000-6] [PMID: 37137927]
- Hohberger B, Lucio M, Schlick S, Wollborn A, Hosari S, Mardin C. OCT-angiography: regional reduced macula microcirculation in ocular hypertensive and pre-perimetric glaucoma patients. *PLoS One.* 2021;16(2):e0246469. [DOI: 10.1371/journal.pone.0246469] [PMID: 33571215]
- Higashide T, Udagawa S, Araie M, Saito H, Takemoto D, Sugiyama K, et al. Relationships between distance from the fovea to the disc and macular retinal layer thickness differ between normal and glaucomatous eyes. *Sci Rep.* 2025;15(1):11554. [DOI: 10.1038/s41598-025-96741-w] [PMID: 40185895]

10. Yoon J, Kim KE, Lee A, Song WK, Kook MS. Foveal avascular zone vessel density is associated with visual field progression in early-stage glaucoma eyes with central visual field damage. *Sci Rep.* 2023;13(1):18285. [DOI: 10.1038/s41598-023-45541-1] [PMID: 37880406]
11. Srivastava A, Sharma V, Kapoor G, Kamal VKBM, Baranwal VK, Singh A. Comparison of diagnostic capability of macular ganglion cell complex vs retinal fiber layer thickness in primary open angle glaucoma patients of Indian origin. *Indian J Ophthalmol.* 2024;72(Suppl 4):S628–33. [DOI: 10.4103/IJO.IJO_1768_23] [PMID: 38391143]
12. Soares RM, Braga JF, da Silva Fernandes J, Ferreira CC, Ribeiro L, Alves H, et al. Automated macular segmentation can distinguish glaucomatous from compressive optic neuropathy. *Graefes Arch Clin Exp Ophthalmol.* 2023;261(6):1701–12. [DOI: 10.1007/s00417-022-05962-6] [PMID: 36625929]
13. Hassanen RM, Sabry A, Mohamed A, Wahba J. Optical coherence tomography angiography changes of the macula in early primary open angle glaucoma compared to healthy eyes. *Minia Journal of Medical Research.* 2023;34(3):81–9. [DOI: 10.21608/MJMR.2023.209587.1393]
14. Acosta PCO, Leon JMS. Correlation of peripapillary retinal nerve fiber layer and macular ganglion cell-inner plexiform layer in early to moderate glaucoma using the Cirrus widefield analysis. *Indian J Ophthalmol.* 2024;72(3):412–6. [DOI: 10.4103/IJO.IJO_697_23] [PMID: 38099378]
15. Bhat KS, Reddy MV, Pai V. Correlation of retinal nerve fiber layer thickness with perimetric staging in primary open-angle glaucoma - A cross-sectional study. *Oman J Ophthalmol.* 2022;15(1):36–42. [DOI: 10.4103/ojo.ojo_345_20] [PMID: 35388245]
16. El-Nimri NW, Manalastas PIC, Zangwill LM, Proudfoot JA, Bowd C, Hou H, et al. Superficial and deep macula vessel density in healthy, glaucoma suspect, and glaucoma eyes. *J Glaucoma.* 2021;30(6):e276–84. [DOI: 10.1097/IJG.0000000000001860] [PMID: 33899812]
17. Hwang HS, Lee EJ, Kim H, Kim TW. Relationships of macular functional impairment with structural and vascular changes according to glaucoma severity. *Invest Ophthalmol Vis Sci.* 2023;64(12):5. [DOI: 10.1167/iovs.64.12.5] [PMID: 37669065]
18. Hashmani N, Hashmani S, Murad A, Baig N. Macular vascular density at the superficial capillary plexus using the optical coherence tomography angiography. *Clin Ophthalmol.* 2019;13:295–302. [DOI: 10.2147/OPHTH.S189515] [PMID: 30799915]
19. Kuryshva NI, Maslova EV, Zolnikova IV, Fomin AV, Lagutin MB. A comparative study of structural, functional and circulatory parameters in glaucoma diagnostics. *PLoS One.* 2018;13(8):e0201599. [DOI: 10.1371/journal.pone.0201599] [PMID: 30138396]

Микроваскуларне промене у жутој мрљи мерене оптичком кохерентном томографијом са ангиографијом код болесника са глаукомом отвореног угла

Рада Зечевић¹, Сунчица Срећковић^{2,3}, Марко Контић¹

¹Специјална очна болница „Београдски офталмолошки центар“, Београд, Србија;

²Универзитетски клинички центар „Крагујевац“, Клиника за офталмологију, Крагујевац, Србија;

³Универзитет у Крагујевцу, Факултет медицинских наука, Катедра за офталмологију, Крагујевац, Србија

САЖЕТАК

Увод/Циљ Глауком је водећи узрок ирверзибилног слепа у свету. Микроваскуларне глаукоматозне промене на нивоу површног капиларног плексуса у жутој мрљи могу да доведу до структуралних промена које корелирају са променама у видном пољу. Циљ ове студије је да се квантитативно анализира површни васкуларни плексус жуте мрље код здравих очију и очију са различитим стадијумима глаукома отвореног угла, употребом оптичке кохерентне томографије са ангиографијом.

Метод У студију су укључена 144 ока – 109 са дијагнозом глаукома отвореног угла и 35 здравих очију. Према Ходап–Андерсон–Паришовој класификацији учињена је даља подела на рани, средњи и узнатредовали стадијум глаукома. Свим болесницима је урађено испитивање видног поља, оптичка кохерентна томографија и оптичка кохерентна томографија са ангиографијом.

Резултати Уочена је статистички значајна редукција густине крвних судова жуте мрље у свим стадијумима глаукома. Фо-

веална густина крвних судова је редукована са $23 \pm 1,9\%$ код здравих очију на $13,9 \pm 1,8\%$ код узнатредовалог стадијума глаукома. Фовеална и парафовеална густина крвних судова значајно је редукована и у раном стадијуму болести. Параметри оптичке кохерентне томографије прогресивно су се смањивали са тежином обољења. Укупна вредност дебљине ганглијског ћелијског комплекса редукована је са $108,5 \pm 5,6 \mu\text{m}$ код здравих очију на $61,2 \pm 6,9 \mu\text{m}$ у узнатредовалом глаукому. Укупна дебљина перипапиларних нервних влакана ретине је смањена са $106,2 \pm 7,9 \mu\text{m}$ код здравих на $48,7 \pm 7,5 \mu\text{m}$ у узнатредовалом стадијуму глаукома.

Закључак Густина крвних судова жуте мрље прогресивно се смањује са напредовањем болести. Ови резултати подстичу потенцијалну употребу оптичке кохерентне томографије са ангиографијом у клиничкој пракси за детектовање раног стадијума глаукома, као и за праћење његове прогресије.

Кључне речи: глауком; жута мрља; интраокуларни притисак; густина крвних судова