



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Tatjana Novaković^{1,2,†}, Zlatica Mirković^{1,2}, Nenad Milošević^{1,2}, Zorica Živković^{1,2}, Dijana Mirić^{1,2}, Jana Mirković¹, Vladan Perić^{1,2}, Jovana Milošević¹

**Assessment of cardiovascular risk factors
in persons with impaired glucose tolerance**

Процена кардиоваскуларних фактора ризика
код особа са смањеном толеранцијом глукозе

¹University of Priština – Kosovska Mitrovica, Faculty of Medicine, Kosovska Mitrovica, Serbia;

²Priština Clinical-Hospital Center, Gračanica, Serbia

Received: February 12, 2018

Revised: April 25, 2019

Accepted: May 7, 2019

Online First: May 23, 2019

DOI: <https://doi.org/10.2298/SARH180212048N>

*Accepted papers are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

†Correspondence to:

Tatjana NOVAKOVIĆ

University of Priština – Kosovska Mitrovica, Filipa Višnjiča bb, 38220 Kosovska Mitrovica, Serbia

E-mail: sonjaziv@yahoo.com

Assessment of cardiovascular risk factors in persons with impaired glucose tolerance

Процена кардиоваскуларних фактора ризика код особа са смањеном толеранцијом глукозе

SUMMARY

Introduction/Objective The aim of the study was to determine the profile of cardiovascular risk factors in patients with impaired glucose tolerance (IGT) in comparison to patients with impaired fasting glucose (IFG).

Methods The study consisted of 222 adult participants with established fasting blood glucose values within the 5.6 - 6.9 mmol/L range. IGT was defined as blood glucose between 7.8 - 11.1 mmol/L in the second hour after the administration of 75g during oral glucose tolerance test (OGTT). IFG is the metabolic state between normal and impaired glucose tolerance, where fasting glucose levels are 5.6 - 6.9 mmol/L, and normal OGTT. IGT was confirmed in 142 of these individuals (107 females and 35 males; aged 54 ± 13 years). The remaining 80 participants (56 females and 24 males, $p=0.329$; aged 53 ± 13 years, $p=0.76$) were considered as IFG group. The following parameters were analyzed in both groups: body mass index (BMI), waist circumference, blood pressure, fasting glucose, fasting insulin levels, HOMA-IR (Homeostasis Model Assessment - Insulin Resistance), C-reactive protein, fibrinogen concentrations and lipid profile.

Results Participants in the IGT group were more obese than in the IFG group (BMI 30.8 ± 5.5 kg/m² vs 26.7 ± 3.8 kg/m², $p<0.001$), and with greater waist circumference (111 ± 12 cm vs 101 ± 6 cm; $p<0.001$). Glucose levels (6.02 ± 0.75 mmol/L vs 5.80 ± 0.62 mmol/L; $p<0.001$), and blood insulin levels (21.61 ± 3.46 vs 6.00 ± 2.8 mIU/L; $p<0.001$), as well as HOMA-IR (5.78 ± 2.68 mIU/L vs 1.54 ± 1.46 mIU/L; $p<0.001$) were also higher in IGT group. Median levels of HbA1c in IGT subjects were higher compared with those in the IFG group, but the difference was not statistically significant ($6.21 \pm 0.75\%$ vs $5.92 \pm 0.43\%$; $p=0.105$). Median hs-CRP levels in the IGT subjects (6.7 ± 4.88 mg/L) were higher than in IFG subjects (5.83 ± 6.47 mg/L), but without statistical significance ($p=0.76$).

Conclusion Our study indicates presence of a large number of cardiovascular risk factors in both groups. Still, obesity, hyperinsulinemia, hypercholesterolemia, hypertriglyceridemia, higher diastolic blood pressure, as well as sedentary lifestyle were statistical significantly more prevalent in patients with IGT.

Keywords: impaired glucose tolerance; impaired fasting glucose; cardiovascular risk factors; diabetes mellitus

САЖЕТАК

Увод/Циљ Циљ студије био је да се утврди који су кардиоваскуларни фактори ризика присутни код особа са смањеном толеранцијом глукозе (СТГ) у поређењу са особама са повишеном гликемијом наште (ПГН).

Методе У студију је било укључено укупно 222 одраслих испитаника са константним налазом концентрације глукозе у распону 5,6 - 6,9 mmol/L. СТГ је стање у којем глукоза у крви достиже ниво од 7,8 - 11,1 mmol/L у другом сату након примене 75g у оралном тесту толеранције глукозе (ОГТТ). ПГН је метаболичко стање између нормалне и оштећене толеранције глукозе, при чему нивои глукозе наште износе 5,6 - 6,9 mmol/L, и нормалним ОГТТ-ом. Групу са СТГ је чинило 142 испитаника (107 жена и 35 мушкараца; просечне старости 54 ± 13 година) код којих је орално глукозо толеранс тестом утврђена СТГ. Осталих 80 испитаника (56 жена и 24 мушкар

ца, $p=0.329$; просечне старости 53 ± 13 година, $p=0.76$) чинило је групу са ПГН. Код свих испитаника анализирали су следећи параметри: индекс телесне масе (ИТМ), обим струка, вредности крвног притиска, гликемије наште, базална инсулинемија, HOMA-IR (хомеостазни модел процене инсулинске резистенције), Ц-реактивни протеин, фибриноген и липидни статус.

Резултати Испитаници у групи са СТГ били су гојазнији него у групи са ПГН (ИТМ: $30,8 \pm 5,5$ kg/m² према $26,7 \pm 3,83$ kg/m²; $p<0,001$), са већим обимом струка (111 ± 12 cm према 101 ± 6 cm; $p<0,001$). Концентрације глукозе ($6,02 \pm 0,75$ mmol/L према $5,80 \pm 0,62$ mmol/L; $p<0,001$) и инсулина у крви ($21,61 \pm 3,46$ према $6,00 \pm 2,80$ mIU/L; $p<0,001$), као и вредности HOMA-IR ($5,78 \pm 2,68$ према $1,54 \pm 1,46$; $p<0,001$), такође су биле више у групи са СТГ. Средње вредности HbA1c код испитаника са СТГ биле су веће у поређењу са особама са ПГН, али није било статистички значајне разлике ($6,21 \pm 0,75$ према $5,92 \pm 0,43\%$, $p = 0,105$). Средњи нивои хс-ЦРП у групи са СТГ били су већи у поређењу са групом са ПГН ($6,70 \pm 4,88$ mg/L према $5,83 \pm 6,47$ mg/L), али без статистички значајне разлике ($p = 0,76$).

Закључак Наша студија указује на присуство великог броја кардиоваскуларних фактора ризика у обе групе. Додатно, потврђено је да су гојазност, хиперинсулинемија, хиперхолестеролемија, хипертриглицеридемија, виши дијастолни крвни притисак, као и седентарни начин живота, статистички значајно чешће заступљени код пацијената са СТГ.

Кључне речи: смањена толеранција глукозе; повишена глукоза наште; кардиоваскуларни фактори ризика; дијабетес мелитус

INTRODUCTION

Prediabetes is defined as a condition in which blood glucose levels are higher than normal but lower than the established thresholds for diagnosing diabetes [1]. Prediabetes includes impaired glucose tolerance (IGT), impaired fasting glucose (IFG), or glycated hemoglobin levels (HbA1c) in the range of 5.7 - 6.4% [1, 2]. Patients with isolated IFG can be distinguished from those with isolated IGT by their fasting and 2 h postload glucose values, as well as the shape of their glucose concentration curves obtained during the oral glucose tolerance test (OGTT). By definition, IGT is a condition in which blood glucose reaches levels between 7.8 and 11.1 mmol/L in the second hour after the administration of 75g of glucose during the OGTT, with the basal fasting glucose levels < 7.0 mmol/L. IFG, on the other hand, is defined as an intermediate metabolic state between normal and impaired glucose tolerance, where fasting glucose levels are between 5.6 and 6.9 mmol /L with normal levels of glucose during two hours in OGTT [1, 2, 3].

Prediabetes has started to receive considerable attention recently because individuals with impaired glucose regulation have been shown to be 4 - 6 times more likely to develop diabetes than those with normal glucose regulation [4].

Previous studies have shown that IGT when accompanied by other risk factors, such as age, gender, smoking, hypertension, obesity and dyslipidemia increases the risk of developing a cardiovascular disease (CVD) [5, 6]. While both IFG and IGT involve insulin resistance, these conditions are distinguished by the site of insulin resistance [5]. High hepatic insulin resistance is a typical finding in patients with IFG, with almost normal values in skeletal muscle. In patients with IGT, the main site of insulin resistance is muscle, with only small changes in liver insulin sensitivity [5]. IGT can easily progress to overt diabetes, whereby macrovascular changes are more pronounced. Thus, there is a relevant correlation with the metabolic syndrome and, consequently, a higher risk of developing CVD [6].

As individuals with IGT have slightly elevated blood glucose levels, which is a relatively weak risk factor for developing CVD, it is highly likely that other cardiovascular factors are also responsible for increased incidence of macrovascular diseases among individuals with IGT [7, 8, 9].

The aim of the present study was to determine which cardiovascular risk factors, other than elevated blood glucose levels, are present in individuals with IGT in comparison to patients with IFG.

METHODS

This prospective study was conducted with the approval from the Ethics Committee of the Faculty of Medicine, University of Pristina, located in Kosovska Mitrovica. Written informed consent was obtained from all subjects involved in the study. The research was conducted in accordance with the Ethical Principles for Medical Research Involving Human Subjects (WMA Declaration of Helsinki, 2013) [10].

Study participants

This prospective study included participants that were screened in the period from March 2016 to January 2017 at the Department of Endocrinology of the Clinical Hospital Centre in Pristina -Gracanica. Criteria for the inclusion into the study were established fasting glucose values in the range of 5.6 - 6.9 mmol/L based on two measurements and the OGTT findings. The exclusion criteria were: age < 40 years, overt diabetes, and history or presence of clinically significant cardiovascular, respiratory, hepatic, renal, gastrointestinal, neurological or infectious disorders capable of altering glucose metabolism. The study sample consisted of 222 patients, 142 (107 females and 35 males, mean age 54 ± 13 years) classified to IGT group, as they met the aforementioned criteria suggested by ADA in 2016 [1]. The remaining 80 subjects (56 females and 24 males, mean age 53 ± 13 years) were classified to IFG group, as they had isolated impaired fasting glucose and normal OGTT. Medical histories, family histories of cardiovascular disease, diabetes and obesity, reports on smoking status, and information about current diseases were obtained from all subjects. Alcohol intake was assessed using the CAGE questionnaire [11]. Physical activity was measured based on the World Health Organization (WHO) recommendations for healthy adults, and physically inactive participants (exercise < 300 min per week) were considered sedentary [12].

Anthropometric, clinical and biochemical measurements

Nutritional status was determined based on the body mass index (BMI) [13]. The specific distribution of adipose tissue, or the size of abdominal fat pad, was estimated by waist circumference, while the size of abdominal fat depots was determined in relation to the reference values provided by the World Health Organization [14]. Blood pressure was measured using the Riva-Rocci sphygmomanometer. Hypertension was diagnosed if systolic and diastolic blood pressure $\geq 140/90$ mmHg [15]. Metabolic syndrome (MetS) was diagnosed according to the American Heart Association National Heart, Lung and Blood Institute - modified Adult Treatment Panel III (ATP III) - criteria [16]. MetS was defined as the presence of least three of the following conditions: abdominal obesity presented as large waist circumference (men: ≥ 102 cm; women ≥ 88 cm), elevated triglyceride levels (≥ 1.7 mmol/L), low HDL cholesterol levels (men: < 1.0 mmol/L; women < 1.2 mmol/L), hypertension ($\geq 130/85$ mmHg or use of anti-hypertensive medication) or elevated fasting blood glucose level (≥ 5.6 mmol/L or use of glucose-lowering drugs).

All subjects were on normal diet prior to the biochemical assessments. On the day of testing, venous blood was collected after overnight fasting and was stored in vials with or without anticoagulant. Thereafter, a standardized oral glucose tolerance test (OGTT) was performed in all subjects following intake of 75 g of anhydrous glucose. Biochemical measurements, including concentrations of serum glucose, total cholesterol (TC), LDL-cholesterol (LDL), HDL-cholesterol (HDL) and triglycerides (TG), were performed by Olympus AU400 Chemistry Analyzer (Olympus, Tokyo, Japan) using commercial test reagent kits, according to the manufacturer's recommendations [15, 16]. Quality of glycemic control was assessed in terms of HbA1c levels, determined from anticoagulated blood samples by turbidimetric inhibition immunoassay method [17]. Fasting plasma insulin levels were measured by RIA, with a sensitivity of 2 mIU/L (normal range: 0.5 - 25.0 mIU/L) [18]. The insulin resistance index (homeostasis model assessment-insulin resistance HOMA-IR) was calculated using the HOMA model: $(\text{fasting insulin} \times \text{fasting glucose})/22.5$ [19]. High sensitivity C-reactive protein (hs-CRP) was measured by a high-sensitivity immunoturbidimetry method (latex) on a Hitachi 902 analyzer using Roche diagnostic reagents (normal range: 0 - 5.0 mg/L). The fibrinogen was measured by turbidimetric method (normal range: 2.0 - 4.0 g/L). The latest recommendations were adopted as target values [20].

Statistical analysis

The obtained data were analyzed using SPSS statistical software, version 19.0. Categorical variables were expressed as prevalence, in percentages, while continuous variables were expressed as mean values and standard deviations (SD) with a confidence interval (95% CI). To analyze the differences between the groups, the Chi-squared test or Fisher's exact test was adopted for testing the probability of the null hypothesis (when some of the expected frequencies were lower than five), while conducting the Student's t-test for the independent samples. Pearson correlation coefficient was used to investigate the association between the variables. The level of statistical significance was set at $p < 0.05$.

RESULTS

The study sample included 222 patients, 142 (107 females and 35 males) of whom fulfilled the criteria for IGT diagnosis. The average age of the subjects assigned to the IGT group was 54 ± 13 years. The remaining 80 subjects (56 females and 24 males; mean age 53 ± 13 years, $p=0.76$) were assigned to the IFG group, as the OGTT test indicated the isolated impaired fasting glucose diagnosis. Basic demographic, anthropometric and clinical data pertaining to these subjects are shown in Table 1.

The age, gender distribution, fasting blood glucose, HbA1c concentration and systolic blood pressure levels of the two groups were not statistically significantly different. In addition, no statistically significant differences in the hs-CRP levels and plasma fibrinogen were noted between the individuals with IGT and those diagnosed with IFG. However, the BMI, waist circumference, serum total cholesterol, LDL-cholesterol, triglycerides, fasting insulin levels, HOMA-IR and diastolic blood pressure were significantly higher, whereas HDL cholesterol concentration was significantly lower in the IGT group compared to the IFG group (Table 1). Also, there were more obese subjects ($BMI \geq 30$ kg/m²) in the IGT group than in the group with IFG, and the difference was statistically significant (Table 2).

In addition, we have found that central obesity, hyperinsulinemia, hypercholesterolemia, hypertriglyceridemia, higher diastolic blood pressure, as well as

sedentary lifestyle were statistically significantly more prevalent in patients with IGT, and report family history of type 2 diabetes mellitus (T2DM) (Table 3).

The correlation analysis showed a statistically significant positive correlation between body mass index (BMI), as well as waist circumference, and the current plasma insulin concentration, while the ratio between these nutritional status values and the current blood glucose levels was not statistically significant (Table 4).

DISCUSSION

The results obtained in the present study have shown a higher incidence of obesity, insulin resistance, dyslipidaemias, lower physical activity, positive history of smoking, and positive family history of obesity and T2DM among the subjects with IGT compared with the IFG group. These results confirm the existence of added risk factors for the development of micro and macrovascular complications in individuals with IGT. In a Danish population-based study of patients with IGT or IFG, the results of fasting laboratory measures and OGTT showed that hypertension, higher BMI, serum triglycerides and plasma glucose levels predicted development of T2DM during the 3.5-year follow-up period [21].

Most of the risk factors that were more common in the IGT group are considered modifiable, since they largely depend on the subject's lifestyle and habits. However, several risk factors cannot be modified, especially aging, which is an inevitable biological process associated with over-expression of some and under-expression of other genes. These changes lead to an unstable metabolic control and increased sensitivity to the effects of external factors, such as, for example, poor nutrition and other lifestyle habits [22, 23]. Consistent with previous studies, a strong association of cardiometabolic risk with increasing age in the IGT group was observed in our study. Our subjects were older than 50 years, which, according to some authors, significantly increases the likelihood of developing IGT and CVD risk [21, 22]. The mean age of our subjects with IGT was 53 ± 13.32 years, which is consistent with the findings of other authors [21], while the participants in the Garcia-Alcala's study were somewhat younger [24]. Our results suggest a slightly higher incidence of IGT in women, which is consistent with the data reported by other authors, who also found that its incidence continues to increase with age [5, 6].

Prediabetes carries some predictive power for macrovascular disease [9], but most of this association appears to be mediated through the metabolic syndrome. Elevated glucose levels are one component of the current consensus definition of the metabolic syndrome [16]. IGT and HbA1c appear to correlate more with the CVD risk than the IFG diagnosis [6, 25]. In the present study, patients with IGT (mean glucose 6.02 mmol/L and the HbA1c of 6.2%) had higher glucose levels than did the IFG group (mean glucose 5.8 mmol/L, HbA1c 5.9%), but the difference was not statistically significant.

Our findings further demonstrated that insulin resistance was significantly present in the IGT group, while it was absent in the group with IFG. The patients with IGT had significantly higher levels of fasting insulin, as well as the HOMA index of insulin resistance. An interesting post mortem analysis conducted by Butler et al. [26] revealed that individuals with IGT have 50% fewer β -cells, indicating that there is a significant loss of β -cell mass in individuals with IGT, which occurs long before the onset of T2DM. Accordingly, for the first time, the latest AACE recommendations [27] put emphasis on obesity, stressing that reduction of body weight, not hyperglycemia, should be the primary objective of prediabetes prevention and treatment. In fact, individuals with IGT, besides changing their lifestyles, have other options to achieve weight loss including drug therapy and surgical procedures, and consequently reduce insulin resistance and hyperglycemia, as well as successfully prevent progression to T2DM, while improving lipid status and blood pressure levels [28].

Our results not only revealed a high incidence of obesity in the IGT group, but also demonstrated that 38.03 % of the subjects had a BMI ≥ 30 kg/m², which was statistically significant in comparison to patients with impaired fasting glucose. Obesity plays an important part in the pathogenesis of insulin resistance, which is commonly seen in IGT patients [5]. In addition, our results show that people with IGT tend to exhibit central obesity, which is associated with higher insulin resistance. The risk for prediabetes increases with increasing BMI levels and is particularly significant in individuals with BMI values above 30 kg/m² [5, 28]. Importantly, a higher BMI level at baseline, but not BMI change, is associated with IFG/IGT to T2DM transition, as confirmed by the present study [28].

In addition, abdominal obesity is a clinical sign of excessive accumulation of visceral fat and is usually associated with a cluster of cardiovascular risk factors defined by the National Cholesterol Education Program IV Adult Treatment Panel III as the metabolic

syndrome [16]. We noted in the IGT group a significant presence of a similar cluster of cardiovascular risk factors to those associated with MetS.

Insulin resistance and visceral obesity are associated with high blood pressure. Both cause an increase in blood pressure by activating the sympathetic nervous system and the renin-angiotensin system, with the consequent retention of sodium and water, volume overload, endothelial dysfunction and impaired renal function [29]. Other than an increase in the number of individuals with IGT, the population-based, cross-sectional study of the prevalence of prediabetes in England that was conducted from 2003 to 2011 also revealed an upward trend in the blood pressure in this population [30].

Regarding serum lipid abnormalities in patients with IGT, in most extant studies, statistically significantly increased levels of total cholesterol, low-density cholesterol and/or triglycerides, and a decrease in high density lipoprotein were found compared to the group with IFG [31,32]. These findings further confirm atherogenic potential in our patients with IGT.

Some authors have reported that increased levels inflammatory cytokines, such as high-sensitivity C-reactive protein and tumor necrosis factor- α , are associated with an increased risk of progression from normoglycemia to prediabetes [33]. In obese individuals, an increase in adipose tissue and abnormal protein with hormone characteristics is common, causing infection of the systemic inflammation type and affecting the metabolic pathway in several processes, resulting in dysglycemia, IFG or IGT and abnormal blood pressure [34]. The median levels of hs-CRP were increased in both groups, but there was no statistically significant difference between their ($p=0.76$). Also, we did not find statistically significant difference in prevalence increased CRP between groups.

Finally, the impact of smoking and exposure to tobacco smoke on the development of prediabetes must be highlighted, even though it remains insufficiently explored. In fact, one of the few studies that have addressed the influence of smoking on the development of T2DM is the Multi-Ethnic Study of Atherosclerosis (MESA) [35].

This study has certain limitations, one of which is a relatively small sample size. In addition, no control group was included in the analyses, and the conclusions pertaining to the CV risks are largely based on extant empirical evidence rather than solely on the study results. Moreover, no information was available on whether the patients have taken some

measures to prevent CV risk factors. Patients with IGT have a significantly higher number of risk factors for the development of cardiovascular disease. Their association increases this risk.

CONCLUSION

Our results have shown that, beyond impaired glucose metabolism, individuals with IGT also exhibited obesity, lipoprotein imbalance, hyperinsulinemia, and higher diastolic blood pressure, as well as reported sedentary lifestyle, cigarette smoking, and family history of diabetes, more frequently than did patients with IFG, which may be implicated in the development of a cardiovascular disease. Thus, there is an evident need for thorough and systematic application of all preventive measures, including lifestyle changes in the first place followed by drugs and other treatment modalities, in reducing the risk for the development of T2DM and cardiovascular diseases.

Conflict of interest: None declared.

REFERENCES

1. American Diabetes Association. Strategies for improving care. Sec. 1. Standards of Medical Care in Diabetes 2016. *Diabetes Care* 2016; 39(1):6 - 12. DOI: 10.2337/dc16-S002.
2. American Diabetes Association, Classification and Diagnosis of Diabetes, *Diabetes Care* 2017;40(1):11 - 24. DOI: 10.2337/dc17-S001.
3. Buysschaert M, Medina JL, Buysschaert B, Bergman M. Definitions (and Current Controversies) of Diabetes and Prediabetes, *Current Diabetes Reviews*, 2016; 12:8-13. PMID: 25612821.
4. Abdul-Ghani MA, Williams K, DeFronzo R, Stern M. Risk of progression to type 2 diabetes based on relationship between postload plasma glucose and fasting plasma glucose. *Diabetes Care* 2006; 29: 1613–8. DOI: 10.2337/dc05-1711, PMID: 16801587.
5. Ferrannini E, Gastaldelli A, Iozzo P. Pathophysiology of prediabetes. *Med Clin North Am.* 2011; 95 (2): 327–39. doi: 10.1016/j.mcna.2010.11.005. PMID:21281836
6. Grundy SM. Pre-diabetes, metabolic syndrome, and cardiovascular risk. *J Am CollCardiol.* 2012; 59:635-43. PMID: 22322078 DOI: 10.1016/j.jacc.2011.08.080.
7. Abdul Ghani M, DeFronzo RA, Jayyousi A. Prediabetes and risk of diabetes and associated complications: impaired fasting glucose versus impaired glucose tolerance: does it matter? *Curr Opin Clin Nutr Metab Care.* 2016; 19:394–9. PMID: 27389083 DOI: 10.1097/MCO.0000000000000307.
8. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med.* 2002; 346(6):393–403. PMID: 11832527 DOI: 10.1056/NEJMoa012512.
9. Brannick B, Wynn A, Dagogo-Jack S. Prediabetes as a toxic environment for the initiation of microvascular and macrovascular complications. *Exp Biol Med (Maywood).* 2016; 241(12):1323-31. PMID: 27302176 PMID: DOI: 10.1177/1535370216654227
10. Helsinki Declaration, which was adopted in 1964 and supplemented on several occasions in 1975, 1983, 1989, 1996, 2000 and 2006. The text of the Declaration Available from: [http // www.wma.net / en / 30publications / 10policies / b3 / from 15:01 2012th](http://www.wma.net/en/30publications/10policies/b3/from15:012012th).
11. Ewing JA. Detecting alcoholism: the CAGE questionnaire. *JAMA.* 1984; 252(14):1905–7. PMID: 6471323.
12. World Health Organization. Global Recommendations on Physical Activity for Health; World Health Organization: Geneva, Switzerland, 2010.
13. National Institutes of Health. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults-The Evidence Report. *National Institutes of Health. Obes Res.* 1998; 6:51 - 209. PMID: 9813653.
14. Geneva: World Health Organization; 2008. World Health Organization (WHO). Waist Circumference and Waist-Hip Ratio. Report of WHO Expert Consultation.
15. Williams B, Mancia G, Spiering W, Agabiti-Rosei E, Azizi M, Burnier M, Clement D, Coca A, De Simone G, Dominiczak A, Kahan T, Mahfoud F, Redon J, Ruilope L, Zanchetti A, Kerins M, Kjeldsen S, Kreutz R, Laurent S, Lip GYH, McManus R, Narkiewicz K, Ruschitzka F, Schmieder R, Shlyakhto E, Tsioufis K, Aboyans V, Desormais I. 2018 Practice guidelines for the management of arterial hypertension of the European Society of Hypertension (ESH) and the European Society of Cardiology (ESC). *Blood Press.* 2018 Dec;27(6):314-340. PMID:30380928; doi: 10.1080/08037051.2018.1527177.
16. Grundy SM, Cleeman JJ, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation.* 2005; 112(17):2735–52. PMID: 16157765 DOI: 10.1161/CIRCULATIONAHA.105.169404.
17. Gosmanov AR, Wan J. Low positive predictive value of hemoglobin A1c for diagnosis of prediabetes in clinical practice. *Am J Med Sci.* 2014; 348(3):191–4. PMID: 24556928, DOI: 10.1097/MAJ.0000000000000223.
18. Melmed S, Polonsky KS, Larsen PR, Kronenberg HM. *Williams Textbook of Endocrinology.* 12th ed. Philadelphia: Elsevier Saunders; 2011. ISBN: 9781437703245
19. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.* 1985; 28(7):412–9. PMID: 3899825.
20. Catapano AL, Graham I, De Backer G, Wiklund O, Chapman MJ, Drexel H, et al. [2016 ESC/EAS Guidelines for the Management of Dyslipidaemias]. *Kardiol Pol.* 2016; 74 (11):1234–318. PMID: 27910077, doi: 10.5603/KP.2016.0157.
21. Rasmussen SS, Glümer C, Sandbaek A, Lauritzen T, Borch-Johnsen K. Determinants of progression from impaired fasting glucose and impaired glucose tolerance to diabetes in a high-risk screened population: 3

- year follow-up in the ADDITION study, Denmark. *Diabetologia*. 2008;51: (2):249–57. PMID: 18060659, DOI: 10.1007/s00125-007-0893-8.
22. Lyssenko V, Laakso M. Genetic screening for the risk of type 2 diabetes: worthless or valuable? *Diabetes Care*. 2013; 36(2):120–6. PMID: 23882036, doi: 10.2337/dcS13-2009.
 23. Yue J, Mao X, Xu K, Liu S, Chen F. & Wang, J, et al. Prevalence, Awareness, Treatment and Control of Diabetes Mellitus in a Chinese Population. *Veves A. ed.PLoS ONE*. 2016; 11: (4).doi:10.1371/journal.pone.0153791.
 24. García-Alcalá H, Genestier-Tamborero CN, Hiraes-Tamez O, Salinas-Palma J, Soto-Vega E. Frequency of diabetes, impaired fasting glucose, and glucose intolerance in high-risk groups identified by a FINDRISC survey in Puebla City, Mexico. *Diabetes MetabSyndrObes*. 2012; 5: 403 - 6. PMID: 23204848, DOI: 10.2147/DMSO.S35545
 25. Gurudevan S, Garg P, Malik S, Ramni K, Farhood S, Harvey H, et al. Impaired fasting glucose is associated with increased severity of subclinical coronary artery disease compared to patients with diabetes and normal fasting glucose: evaluation by coronary computed tomographic angiography. *BMJ Open*. 2016; 6(8):e005148. doi:10.1136/bmjopen-2014-005148.25.
 26. Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC. Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes*. 2003; 52(1):102–10. PMID: 12502499. <https://doi.org/10.2337/diabetes.52.1.102>.
 27. Garvey WT, Mechanick JI, Brett EM, Garber AJ, Hurley DL, Jastreboff AM, Nadolsky K, Pessah-Pollack R, Plodkowski R, Reviewers of the AACEOCPG. American Association of Clinical Endocrinologists and American College of Endocrinology Comprehensive Clinical Practice Guidelines for medical care of patients with obesity executive summary. *EndocrPract*. 2016; 22(7):842–84. 28. PMID: 27472012 DOI: 10.4158/EP161356.ESGL
 28. Ishola AF, Gerstein HC, Engert JC, Mohan V, Diaz R, Anand SS, Meyre D. Longitudinal relationships between glycemic status and body mass index in a multiethnic study: evidence from observational and genetic epidemiology. *Sci Rep*. 2016; 6:30744. PMID: 27480816; doi: 10.1038/srep30744.
 29. Sowers JR. Diabetes mellitus and vascular disease. *Hypertension*. 2013; 61: 943-7. PMID: 23595139, DOI:10.1161/HYPERTENSIONAHA.111.00612
 30. Mainous AG, Tanner RJ, Baker R, et al Prevalence of prediabetes in England from 2003 to 2011: population-based, cross-sectional study *BMJ Open* 2014;4:e005002. doi: 10.1136/bmjopen-2014-005002
 31. Stevens J, Erber-Oakkar E, Cui Z, Cai J, Virani SS, Di Angelantonio E, et al. Cardiovascular disease risk by assigned treatment using the 2013 and 1998 obesity guidelines. *Obesity (Silver Spring)*. 2016; 24(7):1554–60. PMID: 27184463, DOI:10.1002/oby.21496. 10.1161/JAHA.116.004413.
 32. Grøntved A, Koivula RW, Johansson I, Wennberg P, Østergaard L, Hallmans G. Bicycling to work and primordial prevention of cardiovascular risk: a cohort study among swedish men and women. *J Am Heart Assoc*. 2016; 5(11): e004413. DOI: 10.1161/JAHA.116.004413, PMID: 27799235
 33. Sabanayagam C, Shankar A, Lee J, Wong TY, Tai ES. Serum C-reactive protein level and prehypertension in two Asian populations. *J Hum Hypertens*. 2013;27(4):231-6. PMID: 2241874, doi: 10.1038/jhh.2011.117
 34. Ridker PM. Clinical application of C-reactive protein for cardiovascular disease detection and prevention. *Circulation* 2003; 107: 363-9. PMID: 12551853
 35. Keith RJ, Al Rifai M, Carruba C, De Jarnett N, McEvoy JW, Bhatnagar A, et al. Tobacco Use, Insulin Resistance, and Risk of Type 2 Diabetes: Results from the Multi-Ethnic Study of Atherosclerosis. *PLoS One*. 2016; 20:11 (6)e0157592. PMID: 27322410 PMID: DOI: 10.1371/journal.pone.0157592

Table 1. Anthropometric, clinical and laboratory data of studied participants

Parameters	IGT group (n = 142)	IFG group (n = 80)	P-value
Male/ Female (n)	35/107	24/56	0.329
Age (years)	54 ± 13	53 ± 13	0.7591
BMI (kg/m ²)	30.80 ± 5.50	26.70 ± 3.80	< 0.001
Waist circumference (cm)			
Men	111 ± 12	101 ± 6	0.007
Women	97.60 ± 21.70	86.00±11.30	0.019
Total cholesterol (mmol/L)	6.34 ± 0.93	4.52 ± 1.21	< 0.001
HDL-cholesterol (mmol/L)	1.31 ± 0.28	1.42 ± 0.17	0.032
LDL-cholesterol (mmol/L)	4.30± 0.92	2.85 ± 1.08	< 0.001
Triglycerides (mmol/L)	2.12 ± 1.25	1.25± 0.63	< 0.001
Fasting blood glucose (mmol/L)	6.02 ± 1.26	5.80 ± 0.62	0.852
HbA1c (%)	6.21 ± 0.75	5.92 ± 0.43	0.105
Fasting blood insulin (mIU/L)	21.61 ± 3,46	6.00 ± 2.80	< 0.001
HOMA IR	5.78 ± 2.68	1.54 ± 1.46	0.001
Blood pressure (mm Hg)			
Systolic	134.70 ± 17.60	129.80 ± 14.40	0.159
Diastolic	89.70 ± 10.00	82.90±9.00	<0.001
hs-C reactive protein (mg/L)	6.70±4.88	5.83±6.47	0.760
Fibrinogen (g/L)	3.67±4.88	3.36±0.95	0.900

Data are presented either as frequencies (n) or arithmetic mean ± SD; differences between groups were tested by Student's t-test;

BMI – body mass index; HOMA-IR – homeostasis model assessment insulin resistance;

HbA1c – glycated hemoglobin A1c; hs-C-reactive protein

Table 2. Distribution of anthropometric nutrition measures in studied groups

Parameters	IGT group (n=142)	IFG group (n=80)	P- value
BMI (kg/m ²)			
18.5 - 24.9 (%)	4.22	30	<0.001
25.0 - 29.9 (%)	57.75	60	0.835
≥ 30.0 (%)	38.03	10	<0.001
Waist circumference - women			
≤ 80.0 cm (%)	18.69	42.85	0.002
80.1 - 87.9 cm (%)	39.25	30.36	0.286
≥ 88.0 cm (%)	42.06	26.79	0.065
Waist circumference -men			
≤ 94.0 cm (%)	2.86	4.17	0.62
94.1 - 101.9 cm (%)	57.14	58.33	0.912
≥ 102.0 cm (%)	40	37.5	0.775

Differences between groups in anthropometric measures, body mass index (BMI) and waist circumference were tested by χ^2 test

Table 3. Distribution of cardiovascular risk factors in studied groups

Risk factors	IGT group (n = 142)	IFG group (n = 80)	P-value
Metabolic syndrome (%)	39.10	11.00	0.001
Alcohol intake (%)	12.80	13.00	0.964
Low physical activity (%)	45.30	51.20	0.001
Current smoker (%)	23.10	15.00	0.001
Family history of diabetes (%)	22.70	15.10	0.001
Hypertension (%)	23.10	20.00	0.63
LDL cholesterol > 3 mmol/L (%)	32.10	15.00	0.01
Triglycerides > 1.7 mmol/L (%)	31.00	12.00	0.001
hs-CRP 5 mg/L (%)	16.00	15.00	0.85
Fibrinogen > 4.0 g/L (%)	12.00	10.00	0.66

The presence of metabolic syndrome, alcohol intake, physical activity, smoking status and family history of diabetes were assessed as described in the Methods section; differences between groups were tested by χ^2 test;

LDL -- low density cholesterol, hs-CRP – high sensitivity C-reactive protein

Table 4. Relationships between body mass index, waist circumference and concentrations of glucose, insulin in group with IGT

	Fasting blood insulin (mIU/L)		Fasting plasma glucose (mmol/L)	
	Pearson's r	p-value	Pearson's r	p-value
BMI (kg/m ²)	0.365	0.010	-0.059	0.102
Waist circumference (cm)	0.402	0.005	-0.260	0.072

BMI – body mass index

Paper accepted