



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Zorica Popović^{1,†}, Mirjana Đuričković¹, Agima Ljaljević², Snežana Matijević¹,
Kosovka Obradović-Đuričić³

**Assessment of reliability and validity of Montenegrin version of the oral
health impact profile for use among the elderly in Montenegro**

Процена поузданости и валидности црногорске верзије профила утицаја
оралног здравља на квалитет живота старих особа у Црној Гори

¹University of Montenegro, Faculty of Medicine, Study Program for Dentistry, Podgorica, Montenegro;

²Institute of Public Health of Montenegro, Podgorica, Montenegro;

³University of Belgrade, School of Dental Medicine, Clinic for Dental Prosthetics, Belgrade, Serbia

Received: May 28, 2018

Revised: January 23, 2019

Accepted: May 7, 2019

Online First: May 23, 2019

DOI: <https://doi.org/10.2298/SARH180528049P>

***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

†**Correspondence to:**

Zorica POPOVIĆ

University of Montenegro, Faculty of Medicine, Department of Dentistry, Kruševac bb, 81000 Podgorica, Montenegro

E-mail: zorica.jankovicpopovic@gmail.com

Assessment of reliability and validity of Montenegrin version of the oral health impact profile for use among the elderly in Montenegro

Процена поузданости и валидности црногорске верзије профила утицаја оралног здравља на квалитет живота старих особа у Црној Гори

SUMMARY

Introduction/Objective The quality of life of elderly individuals has an active function in oral health; it is of great importance to learn that elders over the age of 65 years demonstrate an increase in seeking dental services. Oral Health Impact Profile-14 (OHIP-14) is especially suitable for use in the elderly. The aim of this study is to examine the reliability and validity of OHIP-14 in the Montenegrin population aged 65 and over and to determine the influence of oral health on the quality of their life in the central part of Montenegro.

Methods The research was conducted from September to December 2016 in the central region of Montenegro, at the Medical University in Podgorica and in the nursing homes of the elderly. The study covered 170 individuals, both genders, with an average age of 72.32 ± 6.85 . The research instrument is OHIP-14 index. Standard statistical tests were used. The statistical significance level is 0.05.

Results The Oral Health Impact Profile-14 is linguistically and culturally adapted for the Montenegrin population. The value of the Cronbach Alpha Index is 0.892. The relationship between correlations for individual issues and total correlations ranges from 0.21 to 0.69. The value of OHIP-14 is 19.24 ± 7.49 . Listed by domains: functional constraints 3.31 ± 1.75 ; physical pain 4.19 ± 1.31 ; psychological discomfort 2.52 ± 1.46 ; physical fitness 4.38 ± 1.40 ; mental incompetence 1.42 ± 1.23 ; social incapacity 1.18 ± 1.27 and handicap 2.21 ± 1.32 .

Conclusion The OHIP-14 index is reliable and valid and is recommended for use in the Montenegrin-speaking area, for the elderly. There is a significant impact of oral health on the quality of life of the elderly in the central part of Montenegro.

Keywords: quality of life; elderly; Montenegro

САЖЕТАК

Увод/Циљ Квалитет живота старих особа у функцији оралног здравља има велики значај с обзиром на повећање броја корисника стоматолошких услуга старијих од 65 година. *Oral Health Impact Profile-14 (OHIP-14)* нарочито је погодан за примену код старих особа. Циљ истраживања је да се испита поузданост и валидност *OHIP-14* код црногорског становништва старости 65 и више година и да утврди утицај оралног здравља на квалитет њиховог живота у средишњем региону Црне Горе.

Метод Истраживање је рађено од септембра до децембра 2016. у средишњем региону Црне Горе, на Медицинском факултету у Подгорици и у домовима за стара лица. Истраживањем је обухваћено 170 особа, оба пола, просечне старости $72,32 \pm 6,85$. Инструмент истраживања је индекс *OHIP-14*. Коришћени су стандардни статистички тестови. Ниво статистичке значајности је 0,05.

Резултати Индекс *OHIP-14* је језички и културолошки прилагођен за црногорско становништво. Вредност индекса Кронбахове алфе износи 0,892. Однос између корелација за поједина питања и укупне корелације креће се од 0,21 до 0,69. Вредност *OHIP-14* износи $19,24 \pm 7,49$. Исказано по доменима: функционална ограничења $3,31 \pm 1,75$; физички бол $4,19 \pm 1,31$; психолошка нелагодност $2,52 \pm 1,46$; физичка неспособност $4,38 \pm 1,40$; психичка неспособност $1,42 \pm 1,23$; социјална неспособност $1,18 \pm 1,27$ и хендикеп $2,21 \pm 1,32$.

Закључак Индекс *OHIP-14* је поуздан и валидан и препоручује се за употребу на црногорском говорном подручју, код старих особа. Постоји значајан утицај оралног здравља на квалитет живота старих особа у средишњем делу Црне Горе.

Кључне речи: квалитет живота; старе особе; Црна Гора

INTRODUCTION

The development of medicine and science in general has led to a prolonged life span. Demographers predict that by 2060 the average age of citizens of the European Union will be 47.2 years. In the next 40 years, people over 65 years of age will make up nearly 30% of the European Union's population [1, 2]. Increased care for the aging, promotion and implementation of the concept of "active aging" aims to contribute to the improvement of health, quality of life and attainment of aging [3]. Knowing the state of the elders oral health

and the impact it has on the quality of life is necessary; in order to be able to plan and organize dental care, as the ever increasing number of individuals will be in need of that service at a future date.

The health of the mouth and teeth is not considered only as the absence of the disease, but also the functional, psychological and social aspect of oral diseases is examined.

This is fully in line with the definition of health of the World Health Organization [4] and the definition of oral health [5]. New definition of oral health: "Oral health is multifaceted and includes the ability to speak, smile, smell, taste, touch, chew, swallow, and convey a range of emotions through facial expressions with confidence and without pain, discomfort, and disease of the craniofacial complex" [5, 6].

The first authors who began to examine the psychosocial aspect of oral health were Cohen and Jago in the 1970s of the 20th century. During the 1980s and 1990s, Reisine, Bailit, Sheiham, Croog, Rosenberg and many others continued the trials. Present day, there is a vast number of clinically verified indices, and each day they are improving postures and creating new indexes [7, 8, 9].

The original version of the OHIP has 49 questions (OHIP-49) [10], but in a large number of researches there are uses of a reduced form of OHIP, with 14 questions. Slade first tested the reduced version in 1997 [11], and then by Locker and Allen in 2002 [12, 13]. Slade had shown that the shorter version has the same reliability and validity as the original version, and due to a smaller number of questions it is particularly suitable for the research of the quality of life in the elderly [11]. To date, OHIP-14 has been translated into a number of languages and has been found in countries of all continents [14–23]. It is a comprehensive and multi-dimensional index that covers all aspects of physical health to the psychological and social sphere of life. It is designed so that it can be applied to people with different general characteristics (education, occupation, culture, social status and other characteristics). Oral Health Impact Profile-14 has experienced linguistic and cultural adaptation, clinical verification and verification in countries in the region of Montenegro and the outlying region [24–27]. The aim of this study is to examine the reliability and validity of OHIP-14 in the Montenegrin population aged 65 and over and to determine the influence of oral health on the quality of their life in the central part of Montenegro.

METHODS

Prior to the realization of the research, the consent was obtained from the Ethical Committee of the Faculty of Medicine of the University of Montenegro in Podgorica. Respondents who participated in the research were previously informed and received their written consent for participation in the research. A research plan had been developed. A dentist who is a specialist in the field of dental prosthetics performed the examination.

The research was conducted in the period from September to December 2016 in the central part of Montenegro, at the Medical University in Podgorica and in the nursing homes for the elderly "Ljubav Spaja" and "Nana" in Spuž and Danilovgrad.

Sample research

The study conducted covered 170 people. The average age of the respondents is 72.32 ± 6.85 (from 65 to 91). The sample accounted for 5% of the total population aged 65 and over, living in the central part of Montenegro. The research includes:

1. Persons aged 65 and over who have appeared for an examination at the Faculty of Medicine in Podgorica - Study program for dentistry, on specific days (Monday and Wednesday) from September to December 2016.

2. The beneficiaries of the services of nursing homes "Nana" and "Ljubav Spaja" in Spuž and Danilovgrad, whose state of general and mental health allowed them to be interviewed.

Research instruments

To determine the impact of oral health on the quality of life, the OHIP-14 index (Oral Health Impact Profile) [11] was used.

The OHIP-14 questionnaire consists of 14 questions. Each response is scored at a value of 0 to 4, using the Likert scale with five responses, depending on the extent to which the patient is affected by the problem. Over the past 12 months, according to one's own

assessment: (0-no problems at all, 1- have problems, 2- often have problems, 3- very often have problems, 4- constantly have problems). The maximum possible number of points is 56. The higher the score, the higher the negative impact of oral health on the quality of life of the respondents.

Since OHIP-14 did not apply in the Montenegrin population, it was first necessary to examine its reliability and validity.

The original OHIP-14 questionnaire was first translated using a two-way (reverse) translation (from English to Montenegrin and back). Two licensed interpreters translated the questionnaire from English into Montenegrin independently of each other. Then a person who was not familiar with the contents of the original text of the questionnaire made a return translation from Montenegrin to English (the person is a good connoisseur of both languages). It was taken into consideration that the essence of the issue was preserved and that the translation was simultaneously adapted to the Montenegrin language, the mentality and culture of the population.

The first 30 respondents of this study who completed the questionnaire (independently or with help of the dentist) understood the significance of all 14 questions and expressed the degree to which certain problems were expressed.

The reliability of the OHIP-14 questionnaire was established using the Cronbach Alpha test, which is standardly used to test the reliability of this and similar questionnaires. The reliability of the questionnaire was assessed by examining the internal consistency (homogeneity) of the answers from the questionnaire. In doing so, testing has been completed in three ways.

1. Omitting individual items (questions), while tracking changes in the Cronbach Alpha value.
2. By following the correlations between items interacting with each other.
3. Calculating total correlations for all items.

For the purposes of further investigation, a regression analysis was performed where the dependent variable was the total OHIP-14 result, and the independent variables were sub-bases OHIP-14.

The OHIP-14 instrument has been tested according to various types of validity. First, the validity of the form and content of the questions was verified in the pilot study. The validity of the OHIP-14 as an instrument was tested by correlating (and examining the existence of a statistically significant difference) of OHIP-14 values depending on the specific characteristics of the respondents (age of the respondent, gender, presence and type of dental remuneration, education and occupation of the respondents).

In statistical data processing, structural validity was tested using Student's T test for two independent samples, one-way ANOVA with Bonferroni or Tamhane T2 after hoc tests. The homogeneity dispersion was checked by Leven test. The level of statistical significance was 0.05.

RESULTS

The study conducted involved 170 people, the average age was 72.32 ± 6.85 , of which: 104 (61.17%) subjects aged 65-74, 53 (31.17%) respondents aged 75-84 and 13 (7.6%) of subjects aged 85 and older. Of the 170 respondents, female examinees were 89 (52.35%) and male examinees were 81 (47.64%). The structure of the respondents according to the level of education: the highest number of respondents 64 (37.64%) have secondary education, 39 (22.94%) have higher education (university), 27 (15.88%) have post secondary education (college), 32 (18.82%) with elementary education, while 8 (4.70%) are without education. The structure of respondents by occupation (prior to retirement) demonstrated that the highest number of respondents were in the field of service activities, 46 (27.05%), 34 (20.00%) from production occupations and 33 in the field of law and economics (19.41%). There is a smaller number of respondents in the field of education 18 (10.58%) and lastly, health workers made up of 14 (8.23%) respondents. The condition of oral health of the respondents who participated in the research was not satisfactory. Out of the 170 people examined, 79(46.47%) are without teeth. It was found that in 83(48.82%) of people there are mobile dental prosthesis in both jaws. Of the respondents examined, 34(20.00%) have mobile dental prosthesis in one jaw and teeth in the opposite jaw are not reimbursed. Denture strings replaced with fixed dental prostheses are present in 16 (9.41%) persons. The respondents who had one jaw mobile, and the other fixed denture was 11(5.88%). In 115(67.65%) of subjects, it was estimated that there was a need for rapid treatment, which mainly relates to the

necessity of making new dental prosthesis. There was no need for emergency dental intervention in any of the respondents.

OHIP-14 reliability analysis

The value of the Cronbach Alpha index, derived from the correlation matrix, is 0.892. The internal consistency of OHIP-14 was first assessed by the analysis of the correlation between the items (questions) (Table 1). Differences in the value of coefficients have shown that no item is superfluous and it is necessary that all questions remain in the questionnaire.

Analyzing results in case of removal of individual items(questions) supports the inclusion of all questions that are in the original questionnaire. The relationship between correlations for individual items(questions) and total, correlations ranges from 0.21 to 0.69. The total correlation analysis (ie correlation between one item and all others) showed that all coefficients are above the minimum recommended value (0.20), which is necessary to include the question in the questionnaire. (Table 2).

All of the above points to the high reliability of the OHIP-14 questionnaires and recommends it to be used in the Montenegrin-speaking area, in older persons.

OHIP-14 Structural Design Analysis

The validity of the OHIP-14 as an instrument was tested by correlating and examining the existence of a statistically significant difference in the value of OHIP-14 according to certain characteristics of the respondents.

Monitoring of the OHIP-14 value in relation to gender, age, education and occupation of respondents (prior to retirement)

Respondents aged 75 and over have a statistically significantly higher OHIP-14 value (20.76 ± 7.39) compared to patients aged 65-74 (18.27 ± 7.43) (Student's T test for two independent samples, $t = -2.132$; $p = 0.034$) (Table 3).

In the male gender ratio, the OHIP-14 value is (19.81 ± 8.53) higher than for female respondents (18.54 ± 6.15), but there is no statistically significant difference between these values (Student's T test for two independent samples, $t = 1.252$; $p = 0.213$) (Table 3).

Respondents who do not have education or have primary education have higher OHIP-14 values (20.50 ± 6.87) compared to those with secondary education (18.78 ± 6.93), post-secondary and higher education ($18.73 \pm 7, 64$). There is no statistically significant difference between these values (ANOVA, $F = 0.391$; $p = 0.815$) (Table 3).

There is no statistically significant difference in the value of OHIP-14 according to the respondents' occupations (ANOVA, $F = 1.072$; $p = 0.384$). The highest value of OHIP-14 is for the respondents who worked in production (20.29 ± 7.50). The lowest value of OHIP-14 is for educators and health workers (18.38 ± 8.35) (Table 3).

Monitoring the OHIP-14 value in relation to the type of dental prosthesis

There is a statistically significant difference in OHIP-14 values relative to the type of dental prosthesis (ANOVA; $F = 111.892$; $p < 0.001$). The Leven test indicates that dispersions may be considered homogeneous ($p = 0.267$) (Table 4).

1. Persons who are partially or completely free of natural teeth and do not have remuneration have a statistically significantly higher OHIP-14 value than:

- a person with a fixed dental prosthesis (Bonferroni test, $p < 0.0001$),
- a person with mobile in both jaws (Bonferroni test, $p < 0.0001$),
- a person with incompletely replaced teeth (there is a mobile denture in one jaw, and in the other lost teeth are not denture) (Bonferroni test, $p < 0.0001$),
- persons with one jaw fixed denture, and in the other mobile (Bonferroni test, $p < 0.0001$).

2. Persons with fixed dental prosthesis have a statistically significantly lower OHIP value of:

- a person with mobile dental prosthesis in both jaws (Bonferroni test, $p < 0.0001$),

- person with incomplete restored tooth (Bonferroni test, $p < 0.0001$),
- persons with one jaw fixed denture, and mobile in the other (Bonferroni test, $p < 0.0001$).

3. Persons with mobile denture in both jaws have a statistically significantly lower OHIP value of:

- a person with incompletely replaced teeth (in one jaw they have mobile denture and in the other they have no natural teeth or dental prosthesis) (Bonferroni test, $p = 0.001$).

The impact of oral health on the quality of life of the elderly

The value of OHIP-14 in this study is 19.24 ± 7.49 (min 0, max 37). The OHIP-14 values expressed in terms of domains are as follows: functional limits 3.31 ± 1.75 , physical pain 4.19 ± 1.31 , psychological discomfort 2.52 ± 1.46 , physical fitness 4.38 ± 1.40 , psychological incompetence 1.42 ± 1.23 , social incapacity 1.18 ± 1.27 and handicap 2.21 ± 1.32 .

The majority of people (91%) expressed difficulties in the field of physical incapacity, ie, Eating disorders (82% answered that "very often" or "constantly" changes the diet regime due to mouth, teeth or dental problems). Physical pain has been present in 90% of people in the last 12 months (74% of respondents "very often" or "constantly" avoids certain foods). Functional constraints were 78% (40% of respondents "often" and 36% "very often" or "constantly" have a feeling of taste change due to the condition of the mouth, teeth and compensation). The impact is least pronounced in domains: psychological incompetence (56%) and social incompetence (51%) (Table 5).

DISCUSSION

The study of the impact of oral health on the quality of life was first carried out in Montenegro. The existence of the OHIP-14 translation into the languages of the states in the region of Montenegro has greatly facilitated the process of linguistic and cultural adjustment of the original text of the OHIP-14.

The Serbian version [24] of this index has 13 questions, because the 5th question was left out (Have you been self-conscious because of your mouth or dentures). In the pilot study, the authors estimated that the translation of this question was such that the issue was not sufficiently understandable for a significant number of respondents and that the questionnaire had a sufficient number of other questions from the psychosocial sphere. Croatian authors translated the fifth question from OHIP-14 questionnaires differently and put it in a questionnaire [24]. In order to adjust the spirit of the language in the Japanese version, a further 14 questions have been added to the 5th. It is considered that the structure of the questionnaire allows such changes, since the overall score is not crucial for the validity of the index [24].

The value of Cronbach alpha coefficient, derived from the correlation matrix, is 0.892 in this study. This is significantly more than the minimum recommended value of 0.70 [13,14]. This recommends OHIP-14 for use among the Montenegrin population of the elderly.

The values of Cronbach alpha coefficient in studies that also examined the applicability of the OHIP-14 index among the elderly (with similar characteristics of the sample) are listed below. In a study conducted in Jordan [17], as in the Montenegrin study, the Cronbach alpha coefficient value is 0.89. In studies conducted in Greece [15] and in Italy [18], the value of Cronbach's alpha coefficient was 0.90. In a study done in Chile [19], the value of Cronbach's alpha coefficient is 0.91. In Poland a study [16] dealing with two GOHAI and OHIP-14 indexes showed that the Cronbach alpha coefficient value was 0.89 for GOHAI and 0.97 for OHIP-14. The value of the Cronbach Alpha coefficient is 0.78, in one of numerous studies in Brazil [20].

The validity of the OHIP-14 index was estimated by correlating the OHIP-14 index values and individual characteristics of the respondents. In the Montenegrin study, patients aged 75+ had a statistically significantly higher OHIP-14 value compared to patients aged 65-74 (20.76 ± 7.39 versus 18.27 ± 7.43). In the elderly, the condition is where the mouth and teeth are worse (more missing teeth) and to a large extent there are mobile dentures older than 5 years. Such remuneration no longer meets aesthetic, functional and prophylactic requirements and does not follow changes in the dental system resulting from the physiological aging process.

The study showed that there is a statistically significant difference in the value of OHIP-14, depending on the type of denture. The highest value was observed in persons whose missing teeth were not replaced by dental prostheses. The lower value is in the holder of fixed prostheses in relation to persons with mobile prostheses. In a study done in Iran [28] where the sample was similar (the average age was 67.5 ± 11 , a high degree of no teeth, 87.5% without front teeth, 85.6% without side teeth, 31.3% had total dentures and 28.8% of the partial), as well as in the research of Montenegrin authors, it was estimated that OHIP-14 was statistically significantly higher in those who were not prosthetically treated compared to persons who did not require prosthetic treatment (25.75 ± 14.5 according to 21.18 ± 9.8 ; $P=0.02$). The Polish authors [17] conducted a two-index study (GOHAI and OHIP-14) in 2014. They concluded that the condition of the teeth, the presence of partial dentures, chewing problems and other problems with the mouth and teeth are significantly related to the values of both indexes. OHIP-14 was significantly higher in patients without teeth (26.0 ± 15.2) compared to those who had their own teeth (12.5 ± 13.0). Prosthetically rehabilitated patients were significantly lower (12.0 ± 12.9) compared to persons without natural teeth and without dental prostheses. (22.5 ± 12.9).

In this study, the researchers estimated that the greatest influence of oral health on their quality of life is present in the domains: physical incapacity, physical pain and functional constraints. The smallest influence is in the domains: psychological disability and social incapacity. If one compares the results of the Montenegrin study with the results of the studies within the region and surrounding countries (Serbia [24], Macedonia [26], Croatia [27], Republic of Serbia [25], Greece [15]), the results are similar. Namely, in the areas of physical incapacity and functional limitations, the respondents expressed the greatest influence. The smallest influence was expressed in the domains of psychological and social incompetence.

Among the questionnaires that are used to examine the effect of oral health on quality of life in the elderly, we decided to use the OHIP-14 in this research because it was used in a large number of countries, on almost all continents. There are large number of scientific papers, references, where OHIP -14 were used. We wanted to do research in Montenegro and compare our results with results in the world. We plan a research that will examine the quality of life before and after prosthodontic rehabilitation in the elderly and we will use the GOHAI questionnaire.

There is a limitation of the study in the use of OHIP-14 in people middle age in Montenegro. In the future, should examine the impact of oral health on quality of life among this and other age groups.

CONCLUSION

The OHIP-14 index is reliable, valid and recommended for use in Montenegro among the elderly. There is a significant influence of the condition in which the mouth, teeth and dental compensations are placed on the quality of life of the elderly in the central region of Montenegro. The influence of these conditions is mostly in the areas of physical incapacity, physical pain and functional limitations, and the smallest in the domains of psychological and social incapacity. In order to improve the condition of oral health and the quality of life of the elderly, it is necessary to plan and continuously work on the development of the dental service and on health literacy and education. It would be useful to form gerontostomatological teams within health institutions and promote the concept of active aging.

Conflict of interest: None declared.

REFERENCES

1. UN report Ageing in the 21st Century: a celebration and a challenge. 2012; 111-18; ISBN 978-0-89714-981-5
2. Sidorenko A. Adjusting to population ageing: evidence for policy action. *Population Horizons*. 2015; 12(2):50–61. [DOI: 10.1515/pophzn-2015-0008]
3. Paúl C, Teixeira L, Ribeiro O. Active Aging in Very Old Age and the Relevance of Psychological Aspects. Original Research Article *Front Med*. October 2017. [DOI: <https://doi.org/10.3389/fmed.2017.00181>] [PMCID: PMC5670329]
4. Huber M, Knottnerus JA, Green L, Horst H, Jadad AR, Kromhout D, et al. How should we define health? *BMJ*. 343. [DOI:<https://doi.org/10.1136/bmj.d4163>] [PMID 21791490. doi:10.1136/bmj.d4163.]
5. Marc L. The New Definition of Oral Health. *Restorative Dentistru*. 2017; 37(1):7. [DOI: 10.11607/prd.2017.1.e]
6. Glick M, Williams DM, Kleinman DV, Vujcic M, Watt RG, Weyant RJ. A new definition for oral health developed by the FDI World Dental Federation opens the door to a universal definition of oral health. *J Am Dent Assoc*. 2016; 147(12):915-917. [DOI: 10.1016/j.adaj.2016.10.001] [PMID: 27886668]
7. Bennadi D, Reddy CVK. Oral health related quality of life. *J Int Soc Prev Community Dent*. 2013; 3(1):1–6. [DOI: 10.4103/2231-0762.115700]
8. Naik A, Kohli N, Flynn P. Validation of the English-language version of 5-item Oral Health Impact Profile. *Journal of Prosthodont. Research*. 2016; 60(2):85-91. [DOI:<http://dx.doi.org/10.1016/j.jpor.2015.12.003>]
9. Hulme C, Robinson P, Douglas G, Baxter P, Gibson B, Godson J, et al. The intensive study: a mixed-methods evaluation of an innovation in commissioning and delivery of primary dental care compared with traditional dental contracting. *Health Services and Delivery Research*. 2016; 4(18):70-99. [DOI: 10.3310/hsdr04180]
10. Adulyanon S, Sheiham A, Slade G. Measuring oral health and quality of life. Slade GD, editor. Chapel Hill: University of North Carolina, Dental Ecology. 1997;1–160.
11. Slade GD. Derivation and validation of a short-form oral health impact profile. *Community Dent Oral Epidemiol*. 1997; 25(4):284-90. [DOI: 10.1111/j.1600-0528.1997.tb00941.x] [PMID: 9332805]
12. Allen F, Locker D. A modified short version of the oral health impact profile for assessing health-related quality of life in edentulous adults. *Int J Prosthodont*. 2002; 15(5):446-50. [PMID:12375458]
13. Locker D, Allen F. What do measures of „oral health-related quality of life“ measure? *Community Dent Oral Epidemiol*. 2007; 35(6):401–11. [DOI:<http://dx.doi.org/10.1111/j.1600-0528.2007.00418.x>] [PMID: 18039281]
14. John MJ, Reißmann DR, Feuerstahler L, Baba K, Larsson P, Čelebić A, et al. Factor Analyses of the Oral Health Impact Profile - Overview and studied population. *J Prosthodont Res*. 2014; 58(1):26–34. [DOI:10.1016/j.jpor.2013.11.002] [PMID: 24440578]
15. Papagiannopoulou V, Papaioannou W, Antonogeorgos G, Yfantopoulos J. Validation of a Greek version of the oral health impact profile (OHIP-14) for use among adults. *Health and Quality of Life Outcomes*. 2012; 10:7. [DOI: 10.1186/1477-7525-10-7]
16. Rodakowska E, Mierzyńska K, Bagińska J, Jamiołkowski J. Quality of life measured by OHIP-14 and GOHAI in elderly people from Białystok, north-east Poland. *BMC Oral Health*. 2014;14:106. [DOI:10.1186/1472-6831-14-106] [PMID:25141902]
17. Al-Habashneh R, Khader YS. Use of the Arabic version of Oral Health Impact Profile-14 to evaluate the impact of periodontal disease on oral health-related quality of life among Jordanian adults. *Journal of Oral Science*. March 2012; 54(1):113-20. [DOI: 10.2334/josnusd.54.113]
18. Corridore D, Campus G, Guerra F, Ripari F, Silvana Sale S, Ottolenghi L. Validation of the Italian version of the Oral Health Impact Profile-14 (IOHIP-14). *Ann Stomatol (Roma)*. 2013; 4(3-4):239–243. [PMID: 24611088]
19. Hoffmann MY, Hoyos JV, Aravena PA. Oral Health Impact Profile in elderly Chileans in southern Chile. *J Oral Res*. 2015; 4(6):365-370. [DOI:10.17126/joralres.2015.071]
20. Murta DP, Lobato WS, Guimaraes ALS, Fraga CA. Impact of Oral Health in the Elderly Patients Quality of Life. *Dent Open J*. 2016; 3(1):4-9. [DOI:<http://dx.doi.org/10.17140/DOJ-3-128>]
21. VikramM, Singh VP. Translation and validation of the Nepalese version of oral health impact profile(OHIP-14) questionnaire. *Oral Biology and Dentistry*. 2014; 2:3. [DOI: 10.7243/2053-5775-2-3]
22. Guarnizo-Herreño C, Watt RG, Fuller E, Steele GJ, Shen J, Morris S, Wildman J, Tsakos G. Socioeconomic position and subjective oral health: findings for the adult population in England, Wales and Northern Ireland. *BMC Public Health*. 2014;14:827. [DOI: 10.1186/1471-2458-14-827] [PMID: 25107286]
23. Lawal FB, Juliana O, Taiwo LO, Arowojolu MO. How Valid are the Psychometric Properties of the Oral Health Impact Profile-14 Measure in Adult Dental Patients in Ibadan Nigeria? *Journal ListEthiop J Health Sciv*. 2014; (3);235-43. [DOI: <http://dx.doi.org/10.4314/ejhs.v24i3.7>] [PMID: 25183930]

24. Ivica S, Tihacek-Šojić Lj, Jelenković A. Adaptation of Oral Health Profile (OHIP-14) index for measuring of health on quality of life elderly to Serbian language. *Vojnosanitetski preglod.* 2009; 66(7):511-515. [DOI:10.2298/VSP0907511S]
25. Eric J, Stančić I, Tihacek Šojić L, Jelenković Popovac A, Tsakos G. Validity and reliability of the Oral Impacts on Daily Performance (OIDP) scale in the elderly population of Bosnia and Herzegovina. *Gerodontology.* 2012; 29(2):1-8. [DOI: 10.1111/j.1741-2358.2011.00584.x] [PMID: 22103883]
26. Šapurik M, Tozija F. OHIP-14 Quesonnaire among elderly over 65 years and its application in the Republic of Macedonia. *Arhiv za javno zdravlje.* 2014; 6(2):36-42. [DOI: <http://dx.doi.org/10.3889/oamjms.2014.064>]
27. Kranjčić J, Mikuš A, Peršić S, Vojvodić D. Factors Affecting Oral Health-Related Quality of Life Among Elderly Croatian Patients. *Acta Stomatol Croat.* 2014 Sep; 48(3):174-182. [DOI: 10.15644/asc48/3/1]
28. Motalebnejad M, Hadian H, Mehdizadeh S, Hajiahmadi M. Validity and reliability of the Persian version of the oral health impact profile (OHIP)-14. *Caspian J Intern Med.* 2011; 2(4):314-320. [PMID:24551438]

Paper accepted

Table 1. Correlation between items (questions) of OHIP-14

	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	1.00													
2	-0.09	1.00												
3	0.41 [#]	0.10	1.00											
4	0.45 [#]	0.15 [*]	0.10	1.00										
5	0.43 [#]	-0.06	0.21 [#]	0.33 [#]	1.00									
6	0.36 [#]	-0.01	0.18 [*]	0.27 [#]	0.42 [#]	1.00								
7	0.37 [#]	0.18 [*]	0.11	0.77 [#]	0.41 [#]	0.22 [#]	1.00							
8	0.18 [*]	0.06	0.15 [*]	0.38 [#]	-0.06	0.28 [#]	0.21 [#]	1.00						
9	0.38 [#]	0.04	0.05	0.47 [#]	0.39 [#]	0.34 [#]	0.54 [#]	0.11	1.00					
10	0.47 [#]	-0.08	0.12	0.44 [#]	0.41 [#]	0.27 [#]	0.42 [#]	0.14	0.57 [#]	1.00				
11	0.47 [#]	-0.05	0.13	0.45 [#]	0.51 [#]	0.29 [#]	0.44 [#]	0.08	0.52 [#]	0.75 [#]	1.00			
12	0.46 [#]	-0.11	0.12	0.43 [#]	0.45 [#]	0.30 [#]	0.48 [#]	0.08	0.57 [#]	0.61 [#]	0.75 [#]	1.00		
13	0.54 [#]	-0.05	0.25 [#]	0.49 [#]	0.46 [#]	0.37 [#]	0.42 [#]	0.26 [#]	0.46 [#]	0.48 [#]	0.48 [#]	0.51 [#]	1.00	
14	0.43 [#]	0.08	0.12	0.47 [#]	0.44 [#]	0.24 [#]	0.57 [#]	0.06	0.52 [#]	0.44 [#]	0.48 [#]	0.55 [#]	0.56 [#]	1.00

1–14 –questions from the OHIP-14 questionnaire;

*p < 0.05;

#p < 0.01

Table 2. Values of Cronbach Alpha coefficients and degree of correlation expressed on issues from Question No. 1 to Question No. 14

Questions	Correlation of questions/total correlation	Value of Cronbach's alpha index when a question is omitted
Q1	0.611	0.835
Q2	0.211	0.873
Q3	0.285	0.850
Q4	0.690	0.826
Q5	0.511	0.83
Q6	0.425	0.843
Q7	0.666	0.828
Q8	0.250	0.855
Q9	0.627	0.833
Q10	0.628	0.835
Q11	0.665	0.833
Q12	0.646	0.834
Q13	0.678	0.830
Q14	0.635	0.834

Q1-Q14 – questions from the OHIP-14 questionnaire

Table 3. Values of the OHIP according to the gender, age, education, and occupation of the respondents (before retirement)

Structure of respondents	Number n	Percentage %	Mean value OHIP-14	Standard deviation \pm SD	Statistical test and statistical significance
Gender					
Male	81	47.64%	19.81	\pm 8.53	Student t-test for two independent samples; t=1.252; p=0.213
Female	89	52.35%	18.54	\pm 6.15	
Age					
65-74 years	104	61.17%	18.27	\pm 7.43	Student t-test for two independent samples; t=-2.132; p=0.034**
75 years and older	66	38.77%	20.76	\pm 7.39	
Education					
Uneducated	8	4.70%	20.50	\pm 6.87	ANOVA test F=0.391; p=0.815
Elementary education		18.82%			
Secondary education	64	37.64%	18.78	\pm 6.93	
Post-secondary (college)	27	15.88%	18.73	\pm 7.64	
Higher education (university)	39	22.94%			
Occupation					
Service activities	46	27.05%	19.09	\pm 7.29	ANOVA test F=1.072; p=0.384
Production	34	20.00%	20.29	\pm 7.50	
Field of law & economic	33	19.41%	19.96	\pm 6.96	
Field of education	18	10.58%	18.38	\pm 8.35	
Health workers	14	8.23%			

**Old significant difference

Table 4. Values of OHIP-14 according to the type of denture

Type of (I) denture	n	Value of OHIP-14	Type of (J) denture	Difference of Mean Value (I and J)	p-value
NN	27	31.66±3.00	FN	24.479*	<0.001
			2MN	14.438*	<0.001
			NZ	11.108*	<0.001
			MN+FN	14.567*	<0.001
FN	16	7.19±4.25	NN	-24.479*	<0.001
			2MN	-10.041*	<0.001
			NZ	-13.371*	<0.001
			MN+FN	-9.913*	<0.001
2MN	82	17.17±5.01	NN	-14.438*	<0.001
			FN	10.041*	<0.001
			NZ	-3.330*	<0.001
			MN+FN	0.129	1.000
NZ	34	20.55±3.38	NN	-11.108*	<0.001
			FN	13.371*	<0.001
			2MN	3.330*	0.001
			MN+FN	3.459	0.156
MN+FN	11	17.54±4.11	NN	-14.567*	<0.001
			FN	9.913*	<0.001
			2MN	-0.129	1.000
			NZ	-3.459	0.156

NN – no denture; 2MN – mobile denture in both jaws; MN + FN – mobile denture in one jaw, fixed denture in the other; NZ – incompletely replaced tooth (due to mobile denture in one jaw, no denture in the other); FN – fixed denture

Table 5. Expression of the influence of oral health on the quality of life

Domains OHIP-14	Quest. OHIP-14	“I had no problems“ n (%)	“Rarely” n (%)	“Often” n (%)	“Very often” “Constantly” n (%)	Domains in (%)
Functional constraints	Q1	76 (46%)	43 (25%)	18 (10%)	33 (19%)	78%
	Q2	18 (10%)	25 (14%)	65 (40%)	72 (36%)	
Physical pain	Q3	27 (15%)	91 (53%)	45 (28%)	7 (4%)	90%
	Q4	5 (3%)	6 (3%)	33 (20%)	126 (74%)	
Psychological discomfort	Q5	34 (20%)	57 (33%)	58 (35%)	21 (12%)	77%
	Q6	41 (24%)	78 (45%)	40 (29%)	11 (6%)	
Physical incapacity	Q7	5 (3%)	11 (6%)	13 (7%)	141 (82%)	91%
	Q8	25 (15%)	58 (34%)	63 (37%)	24 (14%)	
Psychological incompetence	Q9	77 (45%)	67 (39%)	24 (15%)	2 (1%)	56%
	Q10	70 (41%)	81 (47%)	18 (11%)	1 (1%)	
Social incapacity	Q11	70 (41%)	75 (44%)	23 (14%)	2 (1%)	51%
	Q12	95 (55%)	59 (34%)	15 (10%)	1 (1%)	
Handicap	Q13	11 (6%)	95 (56%)	33 (19%)	31 (18%)	78%
	Q14	62 (36%)	86 (50%)	19 (12%)	3 (2%)	

Q1–Q14 – questions from the OHIP-14 questionnaire