



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Case Report / Приказ болесника

Ružica Kravljjanac^{1,2,*}, Slobodan Gazikalović¹, Biljana Vučetić-Tadić^{1,2}

**Infantile hemiconvulsion-hemiplegia epilepsy syndrome
associated with *GRIN2A* gene mutation**

Синдром инфантилне хемиконвулзије – хемиплегије и епилепсије
удружен са мутацијом гена *GRIN2A*

¹Dr. Vukan Čupić Institute for Mother and Child Health Care of Serbia, Belgrade, Serbia;

²University of Belgrade, Faculty of Medicine, Belgrade, Serbia

Received: November 25, 2020

Revised: May 10, 2021

Accepted: May 20, 2021

Online First: June 1, 2021

DOI: <https://doi.org/10.2298/SARH201125049K>

*Accepted papers are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal.

The date the article was made available online first will be carried over.

***Correspondence to:**

Ružica KRAVLJANAC

Faculty of Medicine, University of Belgrade, Institute for Mother and Child Healthcare of Serbia, Radoje Dakić 6–8, 11070 Belgrade, Serbia

E-mail: ruzica.kravljjanac@gmail.com

Infantile hemiconvulsion-hemiplegia epilepsy syndrome associated with *GRIN2A* gene mutation

Синдром инфантилне хемиконвулзије – хемиплегије и епилепсије удружен са мутацијом гена *GRIN2A*

SUMMARY

Introduction Infantile hemiconvulsion-hemiplegia and epilepsy (IHHE) syndrome is defined as a specific syndrome in a patient < 2 years of age, presenting as a new onset refractory status epilepticus with unilateral motor seizures and acute imaging abnormalities, fever, hemiparesis >24 hours, and excluding infectious encephalitis.

Case outline We present results of follow-up in 11 year-old girl with IHHE, associated with *GRIN2A* mutation. The girl had normal development until the first febrile hemiconvulsive status epilepticus at the age of seven months. Neuroimaging initially showed right hemisphere edema, followed by progressive right side hemiatrophy. The patient has resistant epilepsy, left side hemiparesis, and good language and cognitive development.

Conclusion Despite IHHE described many years ago, some syndrome's features, including etiology have remained unexplained. The association of IHHE and *GRIN2A* mutation is described in current manuscript for the first time in scientific literature.

Keywords: Infantile hemiconvulsion; hemiplegia epilepsy; *GRIN2A*; *status epilepticus*

САЖЕТАК

Увод Синдром инфантилне хемиконвулзије – хемиплегије и епилепсије (ИХХЕ) је специфичан ентитет код болесника < 2 године. ИХХЕ представља новонастали рефрактарни *status epilepticus* кога карактеришу: унилатерални моторички напади и акутне неурорадиолошке промене на мозгу, фебрилност, хемипареза у трајању > 24 сата, у одсуству инфективног енцефалитиса.

Приказ болесника Приказујемо резултате праћења једанаестогодишње девојчице са ИХХЕ удруженим са мутацијом *GRIN2A*. Девојчица је имала нормалан развој до првог фебрилног хемиконвулзивног епилептичког статуса у седмом месецу живота. Неурорадиолошки налаз је иницијално показао едем десне хемисфере мозга, а касније прогресивну деснострану хемиатрофију мозга. Болесница је имала резистентну епилепсију, левострану хемипарезу и задовољавајући когнитивни развој и говор.

Закључак Иако је ИХХЕ давно описан, бројне карактеристике синдрома, укључујући етиологију, још су увек необјашњене. Удруженост ИХХЕ и мутације *GRIN2A* у стручној је литератури сада први пут описана.

Кључне речи: инфантилне хемиконвулзије; хемиплегија; епилепсија; *GRIN2A*; *status epilepticus*

INTRODUCTION

Infantile hemiconvulsion – hemiplegia and epilepsy (IHHE) syndrome is defined as a specific syndrome in a patient younger than two years of age, presenting as a new onset refractory status epilepticus (NORSE). IHHE characteristics are: unilateral motor seizures, high grade fever at the time of onset of refractory status epilepticus (SE), and unilaterally abnormal acute imaging, followed by hemiparesis lasting at least 24 hours, and excluding infectious encephalitis [1]. In variable period, from a few months to years later, intractable epilepsy emerges in two thirds to three fourths of children with IHHE. The mechanisms underlying the IHHE syndrome remain unclear. A neuronal injury induced by venous thrombosis and/or hypoxia, and previous abnormalities of the brain were suggested as

underlying mechanism [2]. Gene mutations associated with IHHE syndrome were described in the literature, such as *CACNA1A*, *SCN1A*, *HNRNPU*, *ATPIA3* genes [3, 4, 5]. Initial acute cytotoxic edema of the affected hemisphere at the time of SE is followed by chronic atrophic changes of the same hemisphere [6]. Clinical presentation is characterized by prolonged hemiconvulsions followed by hemiplegia and medication-resistant epilepsy [6–9]. The motor deficit has a variable course, from definitive hemiplegia to complete resolution [6]. The long-term cognitive outcome following IHHE syndrome has been poorly studied [9].

CASE REPORT

We present 11-year-old girl with resistant focal epilepsy and left side hemiparesis. The girl was the second child from an uneventful pregnancy and perinatal history. She had normal global development until the first epileptic attack at the age of seven months. The child was febrile (39°C) for three days before the first seizure and had facial exanthema. NORSE started with the head and eyes deviation toward the left side, clonic jerks of left limbs and impaired consciousness lasting for hours, and repeating in clusters within three days. All hematological, coagulation factors, biochemical and metabolic blood analyses were normal. Cytological, biochemical and microbiological analyses of cerebrospinal fluid were normal. Fundoscopy was normal. Electroencephalographic (EEG) record showed continuous slow epileptic discharges above right hemisphere (figure 1). Urgent brain CT showed edema of right hemisphere (figure 2). The girl was treated according to the protocol for SE: benzodiazepine as the first line (intravenously midazolam in dosage of 0.2 mg/kg), Phenobarbital as a second line (20 mg/kg), and since it was ineffective, midazolam was given in continuous intravenous infusion (0.2 mg/kg/h). Anti-edematous therapy was given (Manitol, Dexamethasone). From the fourth day of hospitalization, the infant was seizure free. Phenobarbital was continuing in maintaining dosage of 5 mg/kg. After two months, focal seizures started in afebrile, awake state: the infant interrupted activities, had staring, cyanosis, deviation of the head lasting up to two minutes. Left side hemiparesis persisted from NORSE. At the age of ten months, the child was referred to our clinic. EEG showed asymmetric background activity, slower above the right hemisphere with almost continuous discharges above parietotemporal right region. Brain MR showed right cerebral white matter hypotrophy. Diagnosis of IHHE was based on the data of febrile refractory hemiconvulsive

SE at the early age followed by recurrent focal-onset seizures, persistent hemiparesis, and MR evidence of brain hemiatrophy. Antiepileptic therapy was corrected, phenobarbital was withdrawn, and carbamazepine was introduced leading to seizure control for one year (25 mg/kg). At the age of 18 months, hemiconvulsive seizures reappeared, and clobasam (1.2 mg/kg) was added. At the age of 28 months, brain MR showed progressive right side hemiatrophy (figure 3). Despite this progression in brain atrophy, the girl had improvement in global development. Psychological testing showed mild language delay, and developmental quotient 80-90. Levetiracetam was introduced due to worsening of seizure control. The girl was continuing to have sporadic focal seizures in frequency up to five seizures per month. At the age of six, the girl started to complain on feeling of fast heart beating, chest pain and tremor of the hands. If it happened during the sleep, the girl awoke and complained of fast heart beating. These episodes were short repeating up to ten per day and night. Cardiologist found normal clinical and heart ultrasound findings, while second degree of A-V block was registered on 24-hour ECG holter. Since carbamazepine might contribute to ECG changes, the dosage was reduced and substituted by oxcarbazepine. Two repeated 24-hour ECG holter in period of two months, showed normal heart rhythm. Since the symptoms were continuing at the age of eight, long term video EEG was performed. No clinical seizures were observed during recording, while EEG showed clear asymmetric background activity, slower above right hemisphere with frequent epileptic discharges. Sulthium was added, but since the girl became drowsy, the drug was excluded. At the age of nine, topiramate was introduced with good effect to both, seizure control and episodes of paroxysmal tachycardia. With current medications (Levetiracetam 40 mg/kg/day, Clobazam 0.9 mg/kg/day, Topiramate 5mg/kg/day) seizure control was achieved. At the end of follow up, the speech and intelligence are normal, as well as physical finding, and only discrete spastic left side hemiparesis was observed. She goes to regular school and she is successful in achieving academic knowledge.

Whole gene sequencing showed that the patient has an extremely rare heterozygous nonsense likely pathogenic variant in the mutation in GRIN2A gene (NM_001134407:c.2776C>T).

All procedures performed were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Written consent to publish all shown material was obtained from the patient's caregiver.

DISCUSSION

IHHE was described more than half a century ago, but still many features of the syndrome have remained unexplained [6]. There is a diagnostic challenge since clinical presentation of IHHE has similarities with other neurological disorders which has to be considered in differential diagnosis such as Rasmussen's encephalitis, cerebrovascular insult and mitochondrial disorders. Infective, genetic, and structural underlying causes are proposed in IHHE etiology [10]. There are case reports of acute infantile hemiplegia associated with exanthema subitum caused by HHV-7 [11]. In our case, at the time of NORSE, exanthema was observed for two days, but PCR for HHV-6 and HHV-7 was not done in regional hospital. Various gene mutations are associated with IHHE [3, 4, 5, 12], and this is the first case with IHHE associated with mutation of *GRIN1A* gene. *GRIN1A* mutation is associated with different epileptic disorders, such as Rolandic epilepsy, Landau-Kleffner syndrome, and continuous spike waves during slow wave sleep [13, 14]. A great number of variants in the *GRIN* genes that encode N-methyl-D-aspartate (NMDA) glutamatergic receptor subunits have been found in patients with neuropsychiatric disorders, including epilepsy. There is evidence that NMDA receptor activation might play an important role in epilepsy, and NMDA receptor antagonists have been considered in treating epilepsy, including SE (MgSO₄, felbamate, ketamine). The results of preclinical studies showed that excessive activation of NMDA receptors has been implicated in the pathogenesis of neuronal injury in acute neurologic disorders, and that magnesium reduced NMDA-mediated brain injury [15, 16]. Magnesium normally does not readily penetrate the fully developed blood-brain barrier (BBB), but BBB permeability is greater in infants especially during seizures, thus, systemically administered magnesium may provide effective therapy in these conditions [15]. NMDA receptors play an important role in the induction of programmed cell death during SE [17, 18]. If further genetic investigation shows more frequent association of IHHE with mutation of the genes responsible for NMDA receptors, the strategy for the treatment of hemiconvulsive SE in IHHE syndrome, should include NMDA antagonists, such as MgSO₄ in high dosage [19], or ketamine [20].

Auvin et al. [2] suggested that anti-edema therapy should be discussed to prevent the cell injury, since the pathological studies showed that cytotoxic edema in HHE is responsible for neuronal damage playing a role in the development of a later epilepsy. Very aggressive anti-edema treatment by mannitol and corticosteroids, and antiseizure therapy in acute phase in our patient, did not prevent further damage and epilepsy.

In patients with IHHE, magnetic resonance imaging (MRI) of the brain reveals evidence of abnormal diffusion within the white matter in the acute phase, but hippocampal sclerosis (HS) and/or hemiatrophy later in life [2, 21]. In our case, three months after initial brain edema, increased T2 hyperintensity, severe gliosis and unilateral brain atrophy were evident by MRI of the brain. The similar MRI unilateral changes were described in literature, only one month after initial hemiconvulsive febrile SE [21]. Recently, some unusual IHHE presentation and etiology were published, such as IHHE in adult cases [22], or associated with cortical dysplasia type III [23], or cobalamin C deficiency [24].

The long-term cognitive outcome following HHE syndrome has been poorly studied, although mental retardation has been reported in cases with IHHE. Recent study reported that mental retardation is not a uniform feature of the IHHE syndrome and demonstrated that reorganization of language to the right cerebral hemisphere or its bilateral representation is common in patients with IHHE syndrome affecting the left cerebral hemisphere [9]. In our patient, right cerebral hemisphere was affected, and language delay was present initially, but during follow up, the girl developed normal speech, and had normal intellectual and school performance. Study which included ten patients with IHHE showed that peri-insular hemispherotomy provides excellent long-term seizure control in patients with drug-resistant hemispheric epilepsy [25]. Recent reports pointed the rule of inflammation in epilepsy associated with *GRIN2A* mutation and good response to intravenous immunoglobulin [26], and this therapy might be considered also in our patient if other treatment failed.

In conclusion, we suggest to explore genetic underlying in all cases with IHHE. Our case with IHHE and *GRIN1A* mutation has favorable neurological and cognitive development despite resistant epilepsy from early age. Since the disorder is very rare, multicenter investigation is recommended, with particular attention to genetic underlying of IHHE. It seems that genetic factors contribute to IHHE pathogenesis in addition to all other proposed factors, such as excessive neuronal excitation, hyperthermia, inflammation and blood-brain

barrier damage [27, 28]. Further investigations are necessary in order to find the answers to present questions about pathogenesis, treatment and prognosis of IHHE.

Conflict of interest: None declared.

Paper accepted

REFERENCES

1. Hirsch LJ, Gaspard N, van Baalen A, Nabbout R, Demeret S, Loddenkemper T. et al. Proposed consensus definitions for new-onset refractory status epilepticus (NORSE), febrile infection-related epilepsy syndrome (FIRES), and related conditions *Epilepsia* 2018; 59:739–744. doi: 10.1111/epi.14016. Epub 2018 Feb 5.
2. Auvin S, Devisme L, Maurage CA, Soto-Ares G, Cuisset JM, Leclerc F, Vallée L. Neuropathological and MRI Findings in an Acute Presentation of Hemiconvulsion-Hemiplegia: A Report With Pathophysiological Implications. *Seizure* 2007; 16(4):371–376. doi: 10.1016/j.seizure.2007.01.009. Epub 2007 Mar 12.
3. Yamazaki S, Ikeno K, Abe T, Tohyama J, Adachi Y. Hemiconvulsion-hemiplegia-epilepsy syndrome associated with CACNA1A S218L mutation. *Pediatr Neurol* 2011; 45:193e6. doi: 10.1016/j.pediatrneurol.2011.04.010.
4. Shimada S, Oguni H, Otani Y, Nishikawa A, Ito S, Eto K, et al. An episode of acute encephalopathy with biphasic seizures and late reduced diffusion followed by hemiplegia and intractable epilepsy observed in a patient with a novel frameshift mutation in HNRNPU. *Brain Dev* 2018; 40(9):813–818. doi: 10.1016/j.braindev.2018.05.010. Epub 2018 May 29. PMID: 29858110.
5. Boonsimma P, Michael Gasser M, Netbaramee W, Wechapinan T, Srichomthong C, Ittiwut C, et al. Mutational and phenotypic expansion of ATP1A3-related disorders: Report of nine cases. *Gene*. 2020 Jul 30; 749:144709. doi: 10.1016/j.gene.2020.144709. Epub 2020 Apr 25. PMID: 32339621.)
6. Gastaut H, Poirier F, Payan H, et al. H.H.E. syndrome: hemiconvulsion, hemiplegia, epilepsy. *Epilepsia* 1959/1960; 1:418–447. doi: 10.1111/j.1528311157.1959.tb04278.
7. Myers KA, Scheffer IE, Archer JS. Hemiconvulsion-hemiplegia-epilepsy evolving to contralateral hemi-Lennox-Gastaut-like phenotype. *Brain Dev*. 2018 May;40(5):425–428. doi: 10.1016/j.braindev.2018.01.005. Epub 2018 Feb 16. PMID: 29459061
8. Kim DW, Kim KK, Chu K, Chung CK, Lee SK. Surgical treatment of delayed epilepsy in hemiconvulsion–hemiplegia–epilepsy syndrome. *Neurology* 2008; 70:2116–2122. doi: 10.1212/01.wnl.0000289192.50924.93.
9. Mirsattari SM, Wilde NJ, Pigott SE. Long-term cognitive outcome of hemiconvulsion–hemiplegia–epilepsy syndrome affecting the left cerebral hemisphere *Epilepsy & Behavior* 2008; 13: 678–680. doi: 10.1016/j.yebeh.2008.07.008. Epub 2008 Aug 16.
10. Hirayama Y, Saito Y, Maegaki Y; Status Epilepticus Study Group. "Symptomatic" infection-associated acute encephalopathy in children with underlying neurological disorders. *Brain Dev*. 2017 Mar;39(3):243–247. doi: 10.1016/j.braindev.2016.09.014. Epub 2016 Oct 22. PMID: 27780632
11. Kawada JI, Kimurab H, Yoshikawac T, Ihirad M, Okumurab A, Morishimae T, Hayakawaa F et al. Hemiconvulsion–hemiplegia syndrome and primary human herpesvirus 7 infection. *Brain & Dev* 2004; 26:412–414. doi:10.1016/j.braindev.2003.12.003 48
12. Robinson JE, Wolfe SM, Kaiser-Rogers K, Greenwood RS. Stroke-Like Presentation Following Febrile Seizure in a Patient with 1q43q44 Deletion Syndrome. *Front Neurol*. 2016 May 4;7:67. doi: 10.3389/fneur.2016.00067. PMID: 27199890; PMCID: PMC4854866
13. Strehlow V, Heyne HO, Vlaskamp DRM, Marwick KFM, Rudolf G, de Bellescize J, et al. GRIN2A-related disorders: genotype and functional consequence predict phenotype. *Brain*. 2019 Jan 1; 142(1):80–92. doi: 10.1093/brain/awy304. PMID: 30544257; PMCID: PMC6308310.
14. Myers SJ, Yuan H, Kang JQ, Tan FCK, Traynelis SF, Low CM. Distinct roles of GRIN2A and GRIN2B variants in neurological conditions. *F1000Res*. 2019 Nov 20; 8:F1000 Faculty Rev-1940. doi: 10.12688/f1000research.18949.1. PMID: 31807283; PMCID: PMC6871362.
15. McDonald JW, Silverstein FS, Johnston MV. Magnesium reduces N-methyl-D-aspartate (NMDA)-mediated brain injury in perinatal rats. *Neuroscience Letters* 1990; 109:234–8. [https://doi.org/10.1016/0304-3940\(90\)90569-U](https://doi.org/10.1016/0304-3940(90)90569-U)
16. Amador A, Bostick CD, Olson H, Peters J, Camp CR, Krizay D, Chen W et al. Modelling and treating GRIN2A developmental and epileptic encephalopathy in mice. *Brain*. 2020 Jul 1;143(7):2039–2057. doi: 10.1093/brain/awaa147. PMID: 32577763; PMCID: PMC7363493.
17. Kapur J. Role of NMDA receptors in the pathophysiology and treatment of status epilepticus. *Epilepsia Open*, 2018; 3(s2):165–168, doi: 10.1002/epi4.1227019
18. Fujikawa DG. Prolonged seizures and cellular injury: understanding the connection. *Epilepsy Behav* 2005; 7(Suppl 3):S3–S11. <https://doi.org/10.1016/j.yebeh.2005.08.003 21>
19. Zeiler FA, Matuszczak M, Teitelbaum J, Gillman LM, Kazina CJ. Magnesium sulfate for non-eclamptic status epilepticus. *Seizure* 2015; 32:100–108. DOI:<https://doi.org/10.1016/j.seizure.2015.09.017>
20. Gaspard N, Foreman B, Judd LM, Nathan BR, McCoy BM, Al Otaibi A, et al. Intravenous ketamine for the treatment of refractory status epilepticus: a retrospective multicenter study. *Epilepsia* 2013; 54:1498–1503. <https://doi.org/10.1111/epi.12247>.
21. Toldo I, Calderone M, Boniver C, Dravet C, Guerrini R, Laverda AM. Hemiconvulsion–hemiplegia–epilepsy syndrome: early magnetic resonance imaging findings and neuroradiological follow-up. *Brain Dev* 2007; 29:109–111. <https://doi.org/10.1016/j.braindev.2006.06.005>.

22. Athwal PSS, Aggarwal S, Eubanks JE, Kahlon S, Singh P. Hemiconvulsion-Hemiplegia-Epilepsy Syndrome in Adult with Uncontrolled Seizures and Phenytoin Toxicity. *Cureus* 2020; 2; 12(5):e7924. doi: 10.7759/cureus.7924
23. Itamura S, Okanishi T, Arai Y, Nishimura M, Baba S, Ichikawa N, Hirayama Y, Ishihara N, Hiraide T, Ishigaki H, Fukuda T, Otsuki Y, Enoki H, Fujimoto A. Three Cases of Hemiconvulsion-Hemiplegia-Epilepsy Syndrome with Focal Cortical Dysplasia Type III. *Front Neurol.* 2019; 20;10:1233. doi: 10.3389/fneur.2019.01233. eCollection 2019
24. Myers KA, Dudley RW, Srour M. Hemiconvulsion-Hemiplegia-Epilepsy in a girl with cobalamin C deficiency. *Epileptic Disord.* 2018; 20(6):545-550. doi: 10.1684/epd.2018.1017
25. Abraham AP, Thomas MM, Mathew V, Muthusamy K, Yoganathan S, Jonathan GE, Prabhu K, Daniel RT, Chacko AG. EEG lateralization and seizure outcome following peri-insular hemispherotomy for pediatric hemispheric epilepsy. *Childs Nerv Syst.* 2019 Jul; 35(7):1189-1195. doi: 10.1007/s00381-019-04067-6. Epub 2019 Jan 30. PMID: 30701299.
26. Hausman-Kedem M, Menascu S, Greenstein Y, Fattal-Valevski A. Immunotherapy for GRIN2A and GRIN2D-related epileptic encephalopathy. *Epilepsy Res.* 2020 Jul;163:106325. doi: 10.1016/j.eplepsyres.2020.106325. Epub 2020 Apr 2. PMID: 32289570.
27. Vega-García A, Ferial-Romero I, García-Juárez A, Munguia-Madera AC, Montes-Aparicio AV, Zequeida-Muñoz E, Garcia-Albavera E, Orozco-Suárez S. Cannabinoids: A New Perspective on Epileptogenesis and Seizure Treatment in Early Life in Basic and Clinical Studies. *Front Behav Neurosci.* 2021; 14:610484. doi: 10.3389/fnbeh.2020.610484. PMID: 33510627; PMCID: PMC7835327.
28. Tan TH, Perucca P, O'Brien TJ, Kwan P, Monif M. Inflammation, ictogenesis, and epileptogenesis: An exploration through human disease. *Epilepsia.* 2021 Feb;62(2):303-324. doi: 10.1111/epi.16788. Epub 2020 Dec 14. PMID: 33316111.




Figure 1. Interictal electroencephalogram during the first day of seizure onset showed high-amplitude, slow, 2 Hz spikes and waves, above the right centro-parietal region with spreading to the whole right

Paper accepted




Figure 2. Brain computed tomography scan during hemiconvulsive *status epilepticus*, showing right cerebral edema

Paper accepted




Figure 3. Brain magnetic resonance scan at the age of 28 months, showing progressive right hemisphere atrophy

Paper accepted