



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original article / Оригинални рад

Ivana Baralić^{1,2,†}, Tamara Stojmenović², Marija Anđelković², Brižita Đorđević³,
Nenad Dikić², Sanja Radojević Škodrić⁴, Snežana Pejić⁵

Effect of combined antioxidant treatment on oxidative stress, muscle damage and sport performance in female basketball players

Утицај комбиноване суплементације антиоксидансима на оксидативни стрес, оштећење мишића и спортску способност код кошаркашица

¹Zvezdara University Medical Center, Belgrade, Serbia;

²Sports Medicine Association of Serbia, Belgrade, Serbia;

³University of Belgrade, Faculty of Pharmacy, Institute for bromatology, Belgrade, Serbia;

⁴University of Belgrade, Faculty of Medicine, Institute for pathology, Belgrade, Serbia;

⁵University of Belgrade, "Vinča" Institute of Nuclear Sciences, Department of Molecular Biology and Endocrinology, Belgrade, Serbia

Received: January 18, 2019

Revised: January 29, 2019

Accepted: May 21, 2019

Online First: June 18, 2019

DOI: <https://doi.org/10.2298/SARH190118063B>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. Srp Arh Celok Lek. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

† **Correspondence to:**

Ivana BARALIĆ

Vatroslava Lisinskog 19/17, 11000 Belgrade

E-mail: ivanabaralic111@gmail.com

Effect of combined antioxidant treatment on oxidative stress, muscle damage and sport performance in female basketball players

Утицај комбиноване суплементације антиоксидансима на оксидативни стрес, оштећење мишића и спортску способност код кошаркашица

SUMMARY

Introduction We determined the impact of antioxidant supplementation by GE132[®] on sports performance, oxidative stress markers, and muscle enzymes activities in professional female basketball players.

Methods Repetitive strength, explosive power, anaerobic endurance, and agility performance were measured before/after the 45-day supplementation period. The FORT (Free Oxygen Radicals Test) and FORD analysis (Free Oxygen Radical Defense) were assessed before/after basketball specific exercise bout, at the beginning/end of observational period. The grade of muscle damage was evaluated by aspartate aminotransferase (AST), creatine kinase (CK) and lactate dehydrogenase (LDH).

Results After supplementation period, significant difference was not recorded regarding the basic motor skills tests. Basketball specific exercise bout induced significant increase in FORT ($p < 0.05$) only at the beginning of supplementation period. Both FORT and FORD significantly decreased over the observational period ($p < 0.001$, $p < 0.01$, respectively). CK and LDH were remarkably lower at the end of observational period ($p < 0.05$), compared to the baseline.

Conclusion Exogenous supplementation with protective nutraceuticals such as those found in GE132[®], could reduce acute/chronic oxidative stress and muscle damage, but had no effect on sport performance in basketball players.

Keywords: oxidative stress; FORT; FORD; basketball; muscle damage

САЖЕТАК

Увод Утврдили смо утицај комбинације антиоксиданата GE132[®] на спортску способност, маркере оксидативног стреса и активности мишићних ензима код професионалних кошаркашица.

Методе Поновљена снага, експлозивна снага, анаеробна издржљивост и агилност су мерени пре/након 45-дневног периода суплементације. ФОРТ (Тест слободних кисеоничних радикала) и ФОРД (Тест антиоксидативне заштите) су процењени пре/ после специфичног кошаркашког тренинга на почетку / крају периода суплементације. Степен оштећења мишића је процењен мерењем активности аспарат аминотрансферазе (АСТ), креатин киназе (ЦК) и лактат дехидрогеназе (ЛДХ) у серуму.

Резултати Након суплементације, није забележена значајна разлика у резултатима тестова моторичких способности у односу на период пре суплементације. Кошаркашки специфични тренинг је изазвао значајно повећање ФОРТ-а ($p < 0.05$) само на почетку периода суплементације. И ФОРТ и ФОРД су значајно опали током посматраног периода ($p < 0.001$, $p < 0.01$, редом). ЦК и ЛДХ су били значајно нижи на крају периода посматрања ($p < 0.05$), у поређењу са вредностима пре суплементације.

Закључак Суплементација са заштитним нутритивним препаратима, као што су они у GE132[®], може смањити акутни/хронични оксидативни стрес и оштећење мишића, али није показан утицаја на спортске перформансе кошаркашица.

Кључне речи: оксидативни стрес; ФОРТ; ФОРД; кошарка; оштећење мишића

INTRODUCTION

Moderate physical activity, associated with a balanced diet, provides many health benefits. However, exhaustive and/or intense training sessions are associated with increased production of reactive oxygen species (ROS) and might lead to oxidative stress in skeletal muscles, blood, and perhaps other tissues [1]. An increasing body of evidence implicates oxidative stress in the pathogenesis of numerous diseases, including diabetes, certain cancers,

and cardiovascular disease. Importantly, exercise-induced oxidative stress might be associated with fatigue, muscle damage, and increased recovery time, which can all affect exercise performance [2, 3, 4].

The body contains an antioxidant defense system that depends on dietary intake of antioxidant vitamins and minerals and the endogenous production of antioxidant compounds such as antioxidant enzymes and numerous non-enzymatic antioxidants, involved in the quenching or removal of free radicals [5]. Physical training may enhance the antioxidant defense system to offset the barrage of ROS generated during exercise [6]. However, body's natural antioxidant defense system might not be sufficient to counteract the increase in ROS production during high-intensity or prolonged intermittent aerobic or anaerobic exercise [7]. Additionally, a large number of the athletes failed to ingest sufficient quantities of fruits and vegetables, which also suggest suboptimal intake of various antioxidants.

Previous research has demonstrated that antioxidants obtained through antioxidant rich diet or supplementation can reduce lipid peroxidation and muscle damage, indicating reduced oxidative stress [8–11]. Although isolated studies reported potential ergogenic properties of antioxidants [12, 13], overall conclusion is that there is limited evidence that dietary supplementation with antioxidants improves human performance.

We investigated the effect of proprietary nutraceutical blend GE132[®] that contains several components with various biological effects including antioxidant properties. The efficacy of this blend has never been tested before. Because of the great interest in using antioxidant nutrients as a preventive and therapeutic tool in clinical medicine and in physical activity, the aim of the present study was to determine the effects of this dietary supplement on oxidative stress, muscle damage and sport performance in female basketball players during competitive half season. It was hypothesized that athletes would demonstrate lower oxidative stress and muscle damage in response to exercise and training after supplementation period.

METHODS

Subjects

Fourteen senior female basketball players, who play in the first league club “Red Star”, Belgrade, Serbia, participated in this study. Athletes gave written consent after explanation of the purpose, demands, and possible risks associated with the study. The protocol was in accordance with the Declaration of Helsinki for Research on Human Subjects and it was approved by the Ethical Committee of Sport Medicine Association of Serbia. All participants passed sports medical examination and were eligible for participation in competitive sport. None of the subjects reported any serious injury or disease 6 months prior to or during the study. Subjects were non-smokers and did not use oral contraceptives, anti-inflammatory drugs or dietary supplements (i.e. antioxidants) one month before and during the study. Subjects were instructed to restrain themselves from making any drastic changes in the diet. All of them had regular menstrual cycles and none of them were in the menstrual phase at the time of blood sampling.

Study design and supplementation

The study was conducted during a competitive half season, over the 45-day period. During this period, athletes were engaged in a controlled training program and participation in the study did not have any effect on previously determined training and competition schedule. Subjects completed two basketball specific exercise bouts, at the beginning and at the end of the observational period. Supplementation started after the first exercise bout and continued for 45 days. Before and after each exercise bout, capillary blood samples were collected for oxidative stress measurement. Venous blood samples were collected at the beginning and at the end of observational period for muscle enzyme analysis. In addition, all players performed basic motor skills tests at the beginning and at the end of the study, in

order to evaluate strength, endurance, and agility as the most commonly used motor skills in basketball.

All subjects received antioxidant complex supplement, GE132[®], during 45 days. Each capsule contained 100 mg of *Ganoderma lucidum* extract (20%), 130 mg of royal jelly (26%), 80 mg of resveratrol (16%), 30 mg of shark protein complex (6%), 80 mg of green tea extract (16%) and 80 mg of rosehip extract (16%). Athletes were told to comply with supplementation protocol and to take two capsules daily, one before lunch and the other before dinner. Capsules were counted upon return of the capsule bottles to assess compliance.

Procedures and measurements

Baseline measurements: Prior to enrollment into the study, all subjects completed a body composition assessment, standard blood chemistry screening, medical history, and physical activity questionnaire. Anthropometric and body composition characteristics were determined by using *Seca* height measuring instrument and *Tanita scale BC-418MA*.

Basketball specific exercise bout: Each exercise bout consisted of a general warm up and stretching (approx. 10 min), technical-tactical training (approx. 30 min), heavy training, including training of counterattacks and simulated full- or half-court basketball games (approx. 40 min), and finally a cool-down phase (approx. 10 min). Each subject served as self-control to eliminate any biological variability in the response to antioxidant supplementation. The exercises were carried out under the same conditions, in the same place and time of the day to avoid circadian variations.

Basic motor skills tests

Strength: In order to evaluate repetitive strength, players performed push-ups and sit-ups to failure. Push-ups were done by placing hands just wider than shoulders. Subjects were

told to keep their elbows fairly close to body and point them back and not to flare them out to the sides. They lowered until their chests were just above the floor, paused for a split second, and then pressed themselves back up. In order to do sit-ups, subjects were told to raise the torso from a supine to a sitting position and then lie back down again without moving the legs. Knees were bent at an angle of 90 degrees and arms were held crossed behind the neck during the test. Only correct repetitions were taken into account.

Explosive strength was assessed by Globus Ergo Tester Platform. Squat jump (SJ), countermovement jump (CMJ), vertical jump (VJ) and left (LL) and right leg (RL) jumps values were obtained. The subjects stood on the contact platform connected to a digital timer that recorded the flight time and height of all jumps. The timer was triggered by the release of the player's feet from the platform, and stopped at the moment of touchdown. Squat jump was performed from a starting position of 90 degrees knee angle without allowing any counter movement. The subjects were told to jump as high as they can without performing a countermovement. The hands were held on the hips during the jump, thus avoiding any arm swing. During the countermovement jump, subjects were in the position with knees slightly bent and moved into a semi-squat position before jumping. Subject's hands also remained on hips throughout countermovement jump. One leg jumps (right and left leg) were performed in the same way as the countermovement jump. The only difference was that subjects were jumping from and landing on the same leg. Considering the vertical jump, the similar technique was used as in the countermovement jump, but subjects were able to perform the arm swing during the jump. Each player performed three jumps and the highest values achieved were recorded.

Endurance: Anaerobic endurance, as an important aspect of basketball, was evaluated by 300-yard shuttle test. Marker cones and lines were placed 25 yards apart to indicate the sprint distance. Stopwatch was used to record results of the test. Athletes were told to run as

fast as they can to the opposite 25-yard line, touch it with their foot, turn and run back to the start. This was repeated six times without stopping (covering 300 yards total). After a rest of five minutes, the test was repeated. The average values of the two 300-yard shuttles were recorded.

Agility: Agility performance of basketball players was assessed by agility t-test. Four cones were set in the court (5 yards = 4.57 m, 10 yards = 9.14 m), and the subjects started at cone A. On the command of the timer, the subjects sprinted to cone B and touched the base of the cone with their right hand. Then, they turned left and shuffled sideways to cone C, and touched its base, this time with their left hand. The subjects then were shuffling sideways to the right to cone D, touching the base with the right hand. At last they shuffled back to cone B, touching it with the left hand, and ran backwards to cone A. The stopwatch was stopped as they passed the cone A. Test was performed three times and average value of all three attempts was taken into account.

Oxidative stress and biochemical measurements: In order to evaluate oxidative stress status, approximately 15 minutes before and 15 minutes after each basketball specific exercise bout, capillary blood samples were collected for FORT (free oxygen radicals test) and FORD (free oxygen radical defense) measurements. The free radical analysis system (free oxygen radicals monitor [FORM] Plus; Callegari S.P.A., Parma, Italy), incorporating a spectrophotometric device reader, was used to measure these parameters.

FORT assay provides an indirect measurement of hydroperoxide, which are intermediate oxidative products of lipids, amino acids and peptides and therefore useful measure of oxidative stress. It is a colorimetric test based on the ability of transition metals, such as iron, to catalyze the breakdown of hydroperoxide into derivative radicals. These derivative radicals are then preferentially trapped by a suitably buffered chromogen - 4-amino-N-ethyl-N-isopropylaniline hydrochloride and develop, in a linear kinetic based

reaction at 37°C, a colored fairly long-lived radical cation spectrophotometrically detectable at 505 nm. The intensity of the color correlates directly with the quantity of radical compounds, which is related to the oxidative status of the sample [14].

FORD test provides an estimation of the overall antioxidant capacity of blood plasma. This test is based on the ability of antioxidants present in plasma to reduce a preformed radical cation. A stable colored cation (photometrically detectable at 505 nm) is formed in the presence of an acidic buffer (pH = 5.2) and an oxidant (FeCl₃). Antioxidant compounds present in the analyzed sample, reduce the radical cation of the chromogen, quenching the color and producing a discoloration of the solution, which is proportional to their amount in the sample [14].

To examine the extent of muscle damage, venous blood samples were collected from the antecubital vein of athletes in serum separator tube using vacutainer system (Greiner Bio-one, Kremsmünster, Austria). The serum separator tubes were placed on ice and left to stand for 30 minutes to facilitate clotting before being centrifuged at 3500 g for 15 minutes to obtain serum. Creatine kinase (CK), lactate dehydrogenase (LDH) and aspartate aminotransferase (AST) were determined in a clinical laboratory using current bioassays.

Statistical analysis

Statistical analyses were performed with the software IBM SPSS Statistic version 20.0. All data were assessed for normality (one-sample Kolmogorov-Smirnov test). FORT and FORD were analyzed using two-way analysis of variance (ANOVA) with repeated measures. Significant changes in muscle enzyme activities at rest, as well as values of basic motor skills test obtained at the beginning and at the end of the study were analyzed using paired sample t-tests. Data were expressed as mean ± SD. A p-value < 0.05 was considered statistically significant.

RESULTS

Descriptive characteristics of the basketball players are shown in Table 1. All subjects consumed the appropriate amount of product throughout the study period. None of the subjects reported adverse effects related to the dietary supplement.

Statistically significant difference was not recorded regarding the results of basic motor skills tests after 45-day supplementation period. The obtained results are shown in Table 2.

Basketball specific exercise bout induced significant increase in FORT at the beginning of observational period ($p < 0.05$), (Figure 1). However, these changes were not recorded after 45 days of supplementation. In addition, the FORT significantly decreased after 45-day supplementation period ($p < 0.001$). ANOVA repeated measures revealed significant decrease in FORD over the observational period ($p < 0.01$).

We established the overall oxidative stress (OS) status of the athletes based on both the FORT and FORD results, according to the manufacturers' direction. Five major profiles have been depicted and the results are presented in Table 3. The number of athletes with OS in progress was reduced and the number of athletes with normal status was increased because of antioxidant supplementation.

The CK and LDH levels at rest, as indicators of muscle damage, significantly decreased after 45 days of supplementation ($p < 0.05$), while no changes were detected regarding the AST levels (Table 4).

DISCUSSION

In the present study, female basketball players were supplemented with complex antioxidant supplement GE132[®] during 45 days. The study was performed just before the start of regular basketball season and after completion of basic conditioning training period.

This period was chosen since preseason training is highly demanding for athletes because they are engaged in both frequent and high intensity workouts with little or no time to recover. This training program allows neuromuscular and endocrine systems to adapt after the loads placed to them and potential redox status adaptations occur [5, 15].

The major findings of this study indicate that:

- 1) Single basketball training session can increase oxidative stress in trained females;
- 2) the antioxidant combination treatment with GE132[®] used in this study can significantly attenuate the rise of blood oxidative stress markers and muscle damage after basketball exercise and training;
- 3) GE132[®] supplementation does not provide benefit for enhancing motor skills of female basketball players.

Based on FORT assay measurement, we found that oxidative stress was significantly increased in response to basketball specific exercise bout before supplementation. Basketball is one of the mixed sports which include aerobic phases (intermittent running at different intensity) and anaerobic phases (jumps, sprints). Therefore, the increased free radical generation in basketball can occur via several pathways: mitochondrial respiration, oxidase enzymatic activity (NAD(P)H oxidase, xanthine oxidase), via phagocytic respiratory burst, a loss of calcium homeostasis and/or the destruction of iron containing proteins [16]. The increase in ROS production resulting from any of the above sources, could lead to oxidative changes of different biomolecules, and increased levels of oxidative stress. The antioxidant supplementation attenuated this increase after exercise bout at the end of study period, as evidenced by the non-significant changes in FORT levels. Additionally, the overall decreased FORT levels indicate less oxidative damage after 45 days of supplementation. The attenuated oxidative stress response is consistent with other studies using antioxidant supplementation [8, 17, 18, 19]. On the other hand, similar changes might occur as a result of adaptive

response to chronic exercise [5]. However, since present study was conducted after the conditioning preseason training, which allowed redox status adaptations, the reduced oxidative stress observed in female basketball players might be the result of antioxidant supplementation alone.

The antioxidant system capacity of plasma, measured by FORD test, depends on individual and synergic effects of different molecules, such as proteins, glutathione, vitamin E, ascorbate, carotenoids, and phenolic compounds. We detected no changes of FORD neither in response to exercise at the beginning of the observational period nor at the end. Mobilization of tissue antioxidant stores into the plasma is an accepted phenomenon that would help maintain antioxidant status in plasma at certain level and protect body against ROS [20]. In addition, soluble plasma antioxidants work synergistically to defend against oxidant production, meaning that when one antioxidant nutrient is lacking at a particular period in time, another could substitute or it may be regenerated by another that is in abundance [21]. These rapid, dynamic responses in order to maintain redox homeostasis could be the reason for non-significant changes observed in FORD after the exercise. However, plasma levels of antioxidants, measured by FORD, decreased over the entire observational period in response to supplementation. This may not increase athlete's susceptibility to oxidative stress, since supplementation decreased oxidative modifications of various biomolecules, as indicated by FORT test. In addition, OS status of the female basketball players was improved after supplementation, judging by the increased number of athletes with normal status and decreased number of athletes with OS in progress. Therefore, this antioxidant supplement may provide protection against the negative health consequences of free radicals produced during training.

Although the mechanisms behind exercise-induced muscle damage are not precisely known, it is believed that along with initial mechanically induced disruption, secondary

damage is caused by the free radical production and subsequent oxidative stress [22]. Some markers, such as AST, CK and LDH, have been used as a way to indicate the grade of muscle cell damage, especially after playing a sport, since microfiber breakdown releases cell content [23]. The supplementation with antioxidants significantly reduced plasma muscle enzyme activities (CK and LDH), suggesting the involvement of oxidant mechanisms on tissue injury induced by the exercise. This finding can be explained by protective effect of antioxidants against lipid peroxidation, resulting in less muscle membrane damage. Our results are in accordance with several studies, which reported beneficial effects of antioxidants in terms of minimizing the rise in muscle enzyme activity in response to exercise [11, 24, 25].

There has been a general inconsistency of outcomes when investigating the role of antioxidant supplementation in exercise performance, with the majority of the studies reporting no benefits. In accordance, in the present study no statistically significant difference was observed regarding repetitive or explosive strength, endurance, or agility performance after supplementation period in comparison to baseline.

CONCLUSIONS

Studies showed that professional athletes are exposed to increased oxidative stress over the periods of intensive training. Exercise-induced oxidative stress might be associated with fatigue, muscle damage, and increased recovery time, which can all affect exercise performance. For that reason, reliable and quick tests for OS status measurements, such as FORD and FORT assays, might be very useful for training and supplementation planning. Exogenous supplementation with protective nutraceuticals such as those found in GE132[®], could reduce acute and chronic oxidative stress during high intensity efforts and provide beneficial effect on muscle function recovery. The results of the present study suggest that GE132[®] supplementation does not enhance performance of female basketball players, but

rather provide protection against detrimental health consequences of ROS produced during training.

ACKNOWLEDGEMENTS

This work was financially supported by grants from the Ministry of Education, Science, and Technological Development, Republic of Serbia (III 46001 and III41027). The authors wish to thank basketball club “Red Star”, Belgrade, Serbia.

Conflict of interest: None declared.

Paper accepted

REFERENCES

1. Finaud J, Lac G, Filaire E. Oxidative stress: relationship with exercise and training. *Sports Med* 2006; 36(4): 327-58. PMID: 16573358. DOI: [10.2165/00007256-200636040-00004](https://doi.org/10.2165/00007256-200636040-00004)
2. Barclay JK, Ansel MH. Free radicals may contribute to oxidative skeletal muscle fatigue. *Can J Physiol Pharmacol* 1991; 69(2): 279-84. PMID: 2054745
3. Pyne DB. Exercise-induced muscle damage and inflammation: a review. *Aust J Sci Med Sport* 1994;26(3-4):49-58. PMID: 8665277
4. Alessio HM. Exercise-induced oxidative stress. *Med Sci Sports Exerc* 1993; 25(2): 218-24. PMID: 8383786
5. Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol Rev.* 2008; 88(4): 1243–76. PMID:18923182. DOI: [10.1152/physrev.00031.2007](https://doi.org/10.1152/physrev.00031.2007).
6. Radak Z, Chung HY, Goto S. Systemic adaptation to oxidative challenge induced by regular exercise. *Free Radical Biol Mol.* 2008; 44(2): 153-9. PMID:18191751.. DOI: [10.1016/j.freeradbiomed.2007.01.029](https://doi.org/10.1016/j.freeradbiomed.2007.01.029).
7. Bloomer RJ, Goldfarb AH, Wideman L, McKenzie MJ, Consitt LA. Effect of acute aerobic and anaerobic exercise on blood markers of oxidative stress. *J Strength Cond Res* 2005; 19(2):276-85. PMID:15903362. DOI: [10.1519/14823.1](https://doi.org/10.1519/14823.1).
8. Goldfarb AH, Bloomer RJ, McKenzie MJ. Combined antioxidant treatment effects on blood oxidative stress after eccentric exercise. *Med Sci Sports Exerc.* 2005; 37 (2): 234-9. PMID: 15692318.
9. Panza VS, Wazlawik E, Ricardo Schütz G, Comin L, Hecht KC, da Silva EL. Consumption of green tea favorably affects oxidative stress markers in weight-trained men. *Nutrition* 2008; 24(5): 433-42. PMID:18337059. DOI:[10.1016/j.nut.2008.01.009](https://doi.org/10.1016/j.nut.2008.01.009).
10. Mastaloudis A, Morrow JD, Hopkins DW, Devaraj S, Traber MG. Antioxidant supplementation prevents exercise-induced lipid peroxidation, but not inflammation, in ultramarathon runners. *Free Radic Biol Med.* 2004; 36(10): 1329-41. PMID:15110397. DOI: [10.1016/j.freeradbiomed.2004.02.069](https://doi.org/10.1016/j.freeradbiomed.2004.02.069).
11. Djordjevic B, Baralic I, Kotur-Stevuljevic J, Stefanovic A, Ivanisevic J, Radivojevic N et al. Effect of astaxanthin supplementation on muscle damage and oxidative stress markers in elite young soccer players. *J Sports Med Phys Fitness* 2012; 52(4): 382-92. PMID:22828460.
12. MacRae HS, Mefferd KM. Dietary antioxidant supplementation combined with quercetin improves cycling time trial performance. *Int J Sport Nutri Exerc Metab* 2006;16(4): 405-19. PMID:17136942.

13. Earnest CP, Lupo M, White KM, Church TS. Effect of astaxanthin on cycling time trial performance. *Int J Sports Med* 2011; 32(11): 882-8. PMID:21984399. DOI: [10.1055/s-0031-1280779](https://doi.org/10.1055/s-0031-1280779).
14. Palmieri B, Sblendorio V. Oxidative stress tests: overview on reliability and use. Part II. *Eur Rev Med Pharmacol Sci* 2007;11(6): 383-99. PMID:18306907.
15. Li JX, Tong CWC, Xu DQ, Chan KM. Changes in membrane fluidity and lipid peroxidation of skeletal muscle mitochondria after exhaustive exercise in rats. *Eur J Appl Physiol* 1999;80(2): 113-7. PMID:10408321. DOI:[10.1007/s004210050566](https://doi.org/10.1007/s004210050566).
16. Fisher-Wellman K, Bloomer RJ. Acute exercise and oxidative stress: a 30 year history. *Dyn Med* 2009; 8: 1. PMID:19144121. DOI:[10.1186/1476-5918-8-1](https://doi.org/10.1186/1476-5918-8-1).
17. Bloomer RJ, Goldfarb AH, McKenzie MJ. Oxidative stress response to aerobic exercise: comparison of antioxidant supplements. *Med Sci Sports Exerc* 2006;38(6): 1098-105. PMID:16775552. DOI:[10.1249/01.mss.0000222839.51144.3e](https://doi.org/10.1249/01.mss.0000222839.51144.3e).
18. Machefer G, Groussard C, Vincent S, Zouhal H, Faure H, Cillard J et al. Multivitamin-mineral supplementation prevents lipid peroxidation during "the Marathon des Sables". *J Am Coll Nutr* 2007; 26(2): 111-20. PMID:17536122.
19. Naziroğlu M, Kiliç F, Uğuz AC, Celik O, Bal R, Butterworth PJ et al. Oral vitamin C and E combination modulates blood lipid peroxidation and antioxidant vitamin levels in maximal exercising basketball players. *Cell Biochem Funct*. 2010;28(4): 300-5. PMID: 20517894. DOI: [10.1002/cbf.1657](https://doi.org/10.1002/cbf.1657).
20. Balakrishnan SD, Anuradha CV. Exercise, depletion of antioxidants and antioxidant manipulation. *Cell Biochem Funct*. 1998;16(4):269–75. PMID:9857489. DOI: [10.1002/\(SICI\)1099-0844\(1998120\)16:4<269::AID-CBF797>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1099-0844(1998120)16:4<269::AID-CBF797>3.0.CO;2-B).
21. Wayner DD, Burtin GW, Ingold KU, Barclay LR, Locke SJ. The relative contributions of vitamin E, urate, ascorbate and proteins to the total peroxy radical-trapping antioxidant activity of human blood plasma. *Biochim Biophys Acta*. 1987; 924(3):408–19. PMID:3593759.
22. Charge SBP, Rudnicki MA. Cellular and molecular regulation of muscle regeneration. *Physiol Rev* 2004; 84(1): 209–38. PMID:14715915. DOI:[10.1152/physrev.00019.2003](https://doi.org/10.1152/physrev.00019.2003).
23. Warren GL, Lowe DA, Armstrong RB. Measurement tools used in the study of eccentric contraction-induced injury. *Sports Med*. 1999; 27(1): 43-59. PMID:10028132. DOI: [10.2165/00007256-199927010-00004](https://doi.org/10.2165/00007256-199927010-00004).

24. Bloomer RJ, Goldfarb AH, McKenzie MJ, You T, Nguyen L. Effects of antioxidant therapy in females exposed to eccentric exercise. *Int J Sport Nutr Exerc Metab* 2004;14(4): 377-88. PMID: 15467097.
25. Miranda-Vilela AL, Akimoto AK, Alves PC, Pereira LC, Gonçalves CA, Klautau-Guimarães MN et al. Dietary carotenoid-rich pequi oil reduces plasma lipid peroxidation and DNA damage in runners and evidence for an association with MnSOD genetic variant -Val9Ala. *Genet Mol Res.* 2009; 8(4):1481-95. PMID:20082261. DOI: [10.4238/vol8-4gmr684](https://doi.org/10.4238/vol8-4gmr684)

Paper accepted

Table 1. Characteristics of female basketball players

Variables	Mean \pm SD
Age (years)	20.6 \pm 2.7
Height (cm)	178.1 \pm 7.8
Weight (kg)	72.9 \pm 10.9
Body Mass Index (kg/m ²)	22.7 \pm 2.2
Percent body fat (%)	20.7 \pm 4.5
Years of training (years)	8.2 \pm 2.7
Hours per week of training (h)	14.2 \pm 4.8

SD – standard deviation.

Paper accepted

Table 2. Performance changes before and after supplementation period in female basketball players

Motor skills	Before supplementation	After supplementation	<i>p</i>
Push-ups	31.5 ± 10.2	38.3 ± 10.9	< 0.05
Sit-ups	95.5 ± 50.1	121.3 ± 50.4	n.s.
Squat jump (cm)	22.4 ± 2.9	23.4 ± 2.8	n.s.
Countermovement jump (cm)	27.5 ± 5.6	28.1 ± 4.3	n.s.
Vertical jump (cm)	36.1 ± 7.91	36.77 ± 6.55	n.s.
Right leg jump (cm)	14.7 ± 3.7	15.8 ± 2.8	n.s.
Left leg jump (cm)	15.2 ± 2.9	15.9 ± 2.0	n.s.
Anaerobic endurance (sec)	75.8 ± 5.7	74.1 ± 4.9	< 0.05
Agility (sec)	11.1 ± 0.5	10.9 ± 0.7	n.s.

Values are presented as mean ± standard deviation.

Table 3. Oxidative stress status interpretation at the beginning and at the end of the study;

Based on both the FORT and FORD results

Profile *	Before supplementation	After supplementation
Normal status FORT < 300 units FORD > 1.00 mmol/L Trolox eq.	2 athletes	12 athletes
Latent OS FORT < 300 units FORD < 1.00 mmol/L Trolox eq.	-	-
Compensated OS FORT – 300–330 units FORD >1.00 mmol/L Trolox eq.	2 athletes	2 athletes
At risk of OS FORT – 300–330 units FORD < 1.00 mmol/L Trolox eq.	-	-
OS in progress FORT > 330 units	10 athletes	-

* Five major OS profiles are defined according to the manufacturers direction.

Table 4. Muscle enzyme activities at rest before and after supplementation period in female basketball players

	Before supplementation	After supplementation	<i>p</i>
AST (IU/L)	21.7 ± 3.6	19.5 ± 2.3	n.s.
CK (IU/L)	143.5 ± 35.6	108.7 ± 32.3	< 0.05
LDH (IU/L)	179.4 ± 14.4	164.0 ± 25.6	< 0.05

Values are presented as mean ± SD.

Paper accepted




Figure 1. a) The levels of reactive oxygen species measured by the FORT test before and after supplementation, in pre-exercise and post-exercise conditions; b) The capacity of plasmatic antioxidants (FORD), measured by FORD test before and after supplementation, in pre-exercise and post-exercise conditions. Values are presented as mean \pm SD, (* $p < 0.05$, ## $p < 0.01$, ### $p < 0.001$).