



Paper Accepted*

ISSN Online 2406-0895

Case Report / Приказ болесника

Ivan Stanković^{1†}, Gojko D. Obradović¹, Radosav Vidaković¹, Ružica Maksimović²,
Ivan Ilić¹, Biljana Putniković¹, Aleksandar N. Nešković¹

Severe short-lasting left ventricular dysfunction associated with a respiratory infection

Тешка, транзиторна дисфункција леве коморе
удружена са респираторном инфекцијом

¹University of Belgrade, University Clinical Hospital Center Zemun, Faculty of Medicine, Department of Cardiology, Belgrade, Serbia;

²University of Belgrade, University Clinical Center of Serbia, Faculty of Medicine, Center for Radiology and Magnetic Resonance Imaging, Belgrade, Serbia

Received: October 13, 2017

Revised: October 8, 2018

Accepted: October 9, 2018

Online First: October 30, 2018

DOI: <https://doi.org/10.2298/SARH171013068S>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

† **Correspondence to:**

Ivan STANKOVIĆ

Department of Cardiology, University Clinical Hospital Centre Zemun, Vukova 9, 11070 Belgrade, Serbia

Email: future.ivan@gmail.com

Severe short-lasting left ventricular dysfunction associated with a respiratory infection

Тешка, транзиторна дисфункција леве коморе
удружена са респираторном инфекцијом

SUMMARY

Introduction Since clinical and electrocardiographic features of various cardiac disorders may overlap, the differential diagnosis of left ventricular (LV) dysfunction may be difficult even for the most experienced physicians. Recent advances in cardiac imaging may help clinicians to establish an accurate diagnosis and initiate adequate treatment. The aim of this case report is to raise awareness of a very short-lasting LV dysfunction during respiratory infections and to underline the importance of multimodality imaging in this clinical setting.

Case outline A previously healthy 37-year-old male presented with atypical chest pain and ST-segment elevation in the inferolateral leads during severe mental stress and acute respiratory infection. Acute myocardial infarction, myocarditis, coronary vasospasm and stress cardiomyopathy were all considered as a differential diagnosis. A rapid onset of severe LV dysfunction and a complete recovery within 4 days was detected by echocardiography and further evaluated by multimodality imaging, including multislice computed tomography and cardiac magnetic resonance imaging.

Conclusion Severe, but very short-lasting LV dysfunction may be triggered by various causes, including upper respiratory tract infections. Since the symptoms of respiratory infections may obscure those of LV dysfunction, myocardial dysfunction in these patients may go undetected with possible serious consequences.

Keywords: chest pain; ST-segment elevation; transient left ventricular dysfunction

САЖЕТАК

Увод С обзиром на то да клиничке и електрокардиографске карактеристике различитих срчаних поремећаја могу да се преклапају, диференцијална дијагноза дисфункције леве коморе може бити тешка и за најiskusније лекаре. Скорашњи напредак на пољу визуализационих метода срца може да помогне клиничарима да поставе тачну дијагнозу и започну адекватно лечење. Циљ овог рада је да скрене пажњу на дисфункцију леве коморе врло кратког трајања током респираторних инфекција и да нагласи значај различитих визуализационих техника у овим клиничким околностима.

Приказ болесника Претходно здрав, 37-годишњи мушкарац јавио се лекару због атипичних болова у грудима и елевације СТ-сегмента у инферолатералним одводима током психичког стреса и акутне респираторне инфекције. Акутни инфаркт миокарда, миокардитис, коронарни вазоспазам и стрес кардиомиопатија разматрани су као диференцијалне дијагнозе. Нагла појава тешке дисфункције леве коморе и потпун опоравак за 4 дана регистрован је ехокардиографијом и даље евалуиран другим визуализационим техникама, укључујући вишеслојну компјутеризовану томографију и магнетну резонанцу срца.

Закључак Бројни узроци, укључујући респираторне инфекције горњих дисајних путева, могу бити окидач тешке, али врло краткотрајне дисфункције леве коморе. С обзиром да симптоми респираторних инфекција могу да маскирају симптоме дисфункције леве коморе, дисфункција миокарда у ових болесника може проћи незапажено са могућим озбиљним последицама.

Кључне речи: бол у грудима, елевација СТ-сегмента, пролазна дисфункција леве коморе

INTRODUCTION

A number of cardiac and noncardiac causes can induce left ventricular (LV) dysfunction, including myocardial ischemia, severe mental stress, endocrine disorders and systemic or myocardial inflammation [1, 2, 3, 4]. Various patient's characteristics and clinical features of the disease are useful for reaching an accurate diagnosis in a typical patient. However, patients having a myocarditis

may share many electrocardiographic (ECG), echocardiographic and clinical features with those with stress cardiomyopathy and acute myocardial infarction (AMI). We present a case of a young adult male who presented with atypical chest pain and ST-segment elevation (STE) during severe mental stress and acute respiratory infection. The role of echocardiography and other imaging modalities in this case scenario is also briefly discussed.

CASE REPORT

A 37-year-old, previously healthy male with multiple risk factors (arterial hypertension, smoking, new-onset diabetes mellitus) for coronary artery disease (CAD) was admitted to hospital due to atypical chest pain associated with STE in the inferolateral ECG leads. A sharp piercing chest pain that partially improves with movement started 5 hours before admission. A week before admission, the patient was treated for an upper respiratory tract infection and also was exposed to severe mental stress. At the time of hospital admission, the patient was afebrile and his physical examination was unremarkable. The initial ECG showed the accelerated junctional rhythm with STE in the leads I, II, aVL, aVF, V3-V6 (Figure 1A). Transthoracic echocardiography (TTE) revealed inferoposterolateral wall hypokinesia with LV ejection fraction (LVEF) of 50% ([Video 1](#)). A mild elevation of cardiac troponin I of 0.08 ng/mL [normal range <0.04 ng/mL] was also noted. The invasive coronary angiography to rule out CAD was considered, but the patient did not consent to any invasive procedure. Despite the treatment with aspirin, benzodiazepines, and morphine, the patient remained anxious and on hospital day 2 complained of recurring severe, sharp, non-radiating chest pain and dyspnea accompanied by S3 gallop. There were no significant changes in blood pressure and heart rate. The troponin I levels rose to 16.6 ng/ml with C-reactive protein (CRP) levels of 76.2 ng/ml (normal range < 5.0 ng/ml) (Figure 2); leukocytes were in the normal range and blood and urine cultures were negative. Beta-blocker, low molecular weight heparin and angiotensin-converting enzyme inhibitor were added to the therapy. The ECG revealed negative T-waves in the leads I, II, aVL, aVF, V3-V6 (Figure 1B) while a repeated TTE showed a worsening of LV function (LVEF drop to 30%) due to global LV hypokinesia with intracavitary spontaneous echo contrast (SEC) (Figure 3A, [Video 2](#)). Two-dimensional speckle-tracking strain analysis showed the global longitudinal peak systolic strain (PSS) of -11.1%. As the patient was extremely anxious and unwilling to undergo invasive coronary angiography, multi-slice computed tomography (MSCT) coronary angiography was performed the same day and revealed normal coronary angiogram (Figure 4). Four days later, a repeated TTE study demonstrated a complete recovery of LV systolic function and disappearance of intracavitary SEC accompanied by the improvement of segmental and global longitudinal PSS (Figure 3B, [Video 3](#)). Troponin I decreased in parallel with CRP and returned to near normal values 5 days following admission. Due to rapid and complete recovery of LV function, myocardial biopsy was not

considered and the patient was discharged 2 days later. Cardiac magnetic resonance imaging (MRI) was not available during the acute phase of the disease; gadolinium-enhanced cardiac MRI was performed 9 months later and was unremarkable (Figure 5). The ECG returned to normal during an uneventful 1-year follow-up (Figure 1C).

DISCUSSION

Transient LV dysfunction may be encountered in a number of cardiac and non-cardiac conditions. As shown in this case, TTE and speckle tracking imaging are excellent tools for detecting rapid changes of LV function – bedside TTE showed that a severe LV dysfunction may last even shorter than usually believed in the setting of acute myocarditis or stress cardiomyopathy. In experienced hands, speckle tracking echocardiography may be useful to detect subtle changes in LV function [5]. However, echocardiography alone is usually not sufficient to make the distinction between ischemic and non-ischemic causes of LV dysfunction, although it has recently been suggested that layer-specific strain may be useful for diagnosing an acute myocarditis [6]. In the present case, the chest pain accompanied by STE and transient LV dysfunction were indicative of an AMI, acute myocarditis and atypical variant of stress cardiomyopathy. The diagnosis of an AMI was ruled out by normal MSCT coronary angiography. However, approximately 3% of patients with AMI may have normal coronary angiograms while unstable coronary lesions may be detected with coronary intravascular ultrasound [7]. Further, approximately 2% of patients initially presenting as AMI are diagnosed with stress cardiomyopathy which commonly affects postmenopausal women and is characterized by transient apical ballooning in the absence of obstructive CAD. Our patient did not fit this paradigm, but it should be noted that patients of both sexes and all ages can be affected and various atypical forms of transient LV dysfunction have been reported [8]. Vasospastic (Prinzmetal) angina might also be considered in this patient since transient chest pain and LV dysfunction can also be due to coronary vasospasm that may occur in angiographically normal epicardial arteries and be triggered by stress conditions. However, the chest pain was atypical and not responsive to nitrates while provocative pharmacologic tests were not carried out. If had been performed early after the onset of disease, cardiac MRI could have contributed to differential diagnosis. The most characteristic MRI feature of stress cardiomyopathy is myocardial edema that appears as high T2 signal intensity with a diffuse or transmural distribution. However, a few weeks after the onset of symptoms, the signal intensity decreases and in many cases it may be impossible to differentiate it from the normal ventricular wall [9]. In case of AMI, myocardial edema is located transmurally or subendocardially, follows vascular distribution and high signal intensity may persist for several months after the event [9]. Finally, in acute myocarditis, the distribution of myocardial edema is more heterogeneous, usually has a mid or subepicardial location and it has been reported to persist on MRI for an average of 111

days (ranging from 56 to 313 days) after the symptom onset [10]. Typical clinical presentations and diagnostic findings in patients presenting with chest pain and left ventricular dysfunction are summarized in Table 1. Mechanisms underlying transient LV dysfunction in various clinical scenarios, including stress cardiomyopathy, neurogenic stunning or myocarditis, are not fully understood. Since the area of edema corresponds to the area of wall motion abnormalities, it could be speculated that swift onset and recovery in some cases of myocarditis can be explained by the presence of myocardial edema and the absence of necrosis and fibrosis [11].

Given the main features of the disease (the ongoing respiratory infection, synchronous rise and fall of cardiac troponin and CRP levels and normal MSCT coronary angiogram), the patient was suspected of having a myocarditis and responded well to supportive treatment including aspirin until CAD was ruled out. Importantly, although data from animal studies found an association of aspirin and increased mortality in myocarditis, this association was not confirmed in a recent prospective, multicenter study [12]. Prompt worsening and recovery of LV function in myocarditis is rare, but possible. In a study by Martin SS et al [13], 6 out of 24 patients who underwent echocardiography during pandemic influenza A (H1N1) in 2009, had a new-onset or worsening LV dysfunction. Improved or normalized ejection fraction was observed in 4 patients, within 4 to 22 days following hospitalization. We present this case to underline the value of multimodality cardiac imaging in patients with chest pain and ambiguous clinical presentations but also to raise awareness of a very short-lasting LV dysfunction that may complicate upper respiratory tract infections. Since the symptoms of respiratory infections may obscure those of LV dysfunction, myocardial dysfunction in these patients may go undetected. Its prevalence and clinical importance is yet to be determined.

REFERENCES

1. Aurigemma GP, Tighe DA. Echocardiography and reversible left ventricular dysfunction. *Am J Med.* 2006; 119(1):18–21. PMID: 16431177 DOI: 10.1016/j.amjmed.2005.10.031.
2. Sosa S, Banchs J. Early recognition of apical ballooning syndrome by global longitudinal strain using speckle tracking imaging--the evil eye pattern, a case series. *Echocardiography.* 2015;32(7):1184–92. PMID: 25556622 DOI: 10.1111/echo.12875.
3. Pešut DP, Petrović ŽM, Vasić NR, Stević RS. Synchronous advanced pulmonary tuberculosis and acute virus myocarditis mimicked Wegener granulomatosis in a 26-year-old man – Case report. *Srp Arh Celok Lek.* 2016; 144(11-12):645–9. DOI: 10.2298/SARH1612645P.
4. Simić D, Milojević I, Budić I, Strajina V. Myocarditis Exacerbation in a Child Undergoing Inguinal Hernioplasty after Viral Infection. *Srp Arh Celok Lek.* 2009; 137(9-10):537–9. PMID: 19950763 DOI: 10.2298/SARH0910537S.
5. Stankovic I, Cvorovic V, Putnikovic B, Vuksanovic I, Panic M, Neskovic AN. Two-dimensional speckle tracking-derived strain to distinguish acute coronary syndrome from a marked early repolarization in a patient with chest pain: a fancy gadget or a useful tool? *Echocardiography.* 2014; 31(2):E48–51. PMID: 24134415 DOI: 10.1111/echo.12411.
6. Caspar T, Germain P, El Ghannudi S, Morel O, Samet H, Trinh A et al. Acute myocarditis diagnosed by layer-specific 2D longitudinal speckle tracking analysis. *Echocardiography.* 2016; 33(1):157–8. PMID: 26343232 DOI: 10.1111/echo.13045.
7. Reynolds HR, Srichai MB, Iqbal SN, Slater JN, Mancini GB, Feit F, et al. Mechanisms of myocardial infarction in women without angiographically obstructive coronary artery disease. *Circulation.* 2011; 124(13):1414–25. PMID: 21636193 DOI: 10.1161/CIRCULATIONAHA.111.026542.
8. Cvorovic V, Stankovic I, Panic M, Stipac AV, Zivkovic A, Neskovic AN, et al. Establishing the diagnosis of inverted stress cardiomyopathy in a patient with cardiac arrest during general anesthesia: a potential role of myocardial strain? *Echocardiography.* 2013; 30(6):E161–3. PMID: 23489387 DOI: 10.1111/echo.12175.
9. Fernández-Pérez GC, Aguilar-Arjona JA, de la Fuente GT, Samartín M, Ghioldi A, Arias JC, et al. Takotsubo cardiomyopathy: assessment with cardiac MRI. *Am J Roentgenol.* 2010 ; 195(2):W139–45. PMID: 20651173 DOI: 10.2214/AJR.09.3369.
10. Assomull RG, Lyne JC, Keenan N, Gulati A, Bunce NH, Davies SW, et al. The role of cardiovascular magnetic resonance in patients presenting with chest pain, raised troponin, and unobstructed coronary arteries. *Eur Heart J.* 2007; 28(10):1242–9. PMID: 17478458 DOI: 10.1093/eurheartj/ehm113.
11. Friedrich MG. Myocardial edema – a new clinical entity? *Nat Rev Cardiol.* 2010;7(5):292–6. doi: 10.1038/nrcardio.2010.28.
12. Imazio M, Brucato A, Barbieri A, Ferroni F, Maestroni S, Ligabue G et al. Good prognosis for pericarditis with and without myocardial involvement: results from a multicenter, prospective cohort study. *Circulation.* 2013;128(1):42–9. doi: 10.1161/CIRCULATIONAHA.113.001531.
13. Martin SS, Hollingsworth CL, Norfolk SG, Wolfe CR, Hollingsworth JW. Reversible cardiac dysfunction associated with pandemic 2009 influenza A(H1N1). *Chest.* 2010;137(5):1195–7. doi: 10.1378/chest.10-0032.

Table 1. Typical clinical presentations and diagnostic findings in patients presenting with chest pain and left ventricular dysfunction*

Parameter	Severe mental stress	Recent infection	ST-segment elevation	Transient LV dysfunction	Elevated troponin	Abnormal coronary angiogram	Late gadolinium enhancement
Acute MI	++	–	+++	+	+++	+++	++ (subendocardial)
Stress CMP	+++	–	++	+++	+	+/- §	–
Coronary vasospasm	+	–	+++	++	+	+/- §	–
Acute myocarditis	–	++	++	++	++	+/- §	++ (subepicardial)

MI – myocardial infarction; CMP – cardiomyopathy; LV – left ventricle;

*number of + in table corresponds to the likelihood of the occurrence of diagnostic findings;

§may be present in the presence of concomitant coronary artery disease




Figure 1. (A) Electrocardiogram at hospital admission showing an accelerated junctional rhythm with ST-segment elevation in the leads I, II, aVL, aVF, V3-V6; (B) electrocardiogram during chest pain (day 2) showing ST-segment elevation and negative T-waves in the leads I, II, aVL, aVF, V3-V6; (C) normal 12-lead electrocardiogram recorded after full recovery




Figure 2. Changes in cardiac markers and C-reactive protein over time after the onset of symptoms to 5 days of admission; numbers in parentheses stand for the value of MB isoenzyme at corresponding time points




Figure 3. (A) Echocardiogram recorded during chest pain (day 2); top panel: apical 2-chamber view showing intracavitary spontaneous echo contrast (yellow arrows); bottom panel: bull's eye display of speckle tracking derived longitudinal peak systolic strain (LPSS) showing significantly decreased segmental strain values (global longitudinal strain of -11.6%); (B) echocardiogram recorded 4 days later; top panel: apical 2-chamber view without SEC; bottom panel: almost complete recovery of segmental LPSS (global longitudinal strain of -18.6%)




Figure 4. Multi-slice computed tomography coronary angiography showing normal coronary arteries; RCA – right coronary artery; LAD – left anterior descending coronary artery; LCx/RI – left circumflex/ramus intermedius coronary artery.




Figure 5. Cardiac magnetic resonance imaging after contrast administration, four-chamber (A) and short-axis (B) view; no myocardial late gadolinium enhancement is present; LV – left ventricle; RV – right ventricle

SUPPLEMENTARY MATERIAL AVAILABLE ONLINE (please click on the text for respective videos)

Video 1. Two-dimensional echo loops recorded at hospital admission (quad view). Note inferposterolateral wall hypokinesia with left ventricular ejection fraction of 50%.

Video 2. Two-dimensional echo loops recorded at hospital day 2, during the episode of chest pain (quad view). There is a global hypokinesia with intracavitary spontaneous echo contrast (“smoke” within the left ventricular cavity, best seen in the apical 2-chamber view – upper right panel).

Video 3. Two-dimensional echo loops recorded at hospital day 6 (quad view). There is a complete recovery of left ventricular systolic function and the disappearance of spontaneous echo contrast within the left ventricle.