



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original article / Оригинални рад

Vladimir Manojlović^{1,2,†}, Đorđe Milošević^{1,2}, Nebojša Budakov^{1,2}, Dragan Nikolić^{1,2}

Incomplete Circle of Willis and cerebrovascular reactivity in asymptomatic patients before and after carotid endarterectomy

Некомплетност Вилисовог прстена и цереброваскуларна реактивност код асимптоматских пацијената пре и после каротидне ендартеректомије

¹University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia;

²Clinical center of Vojvodina, Department for vascular and endovascular surgery, Novi Sad, Serbia

Received: February 3, 2020

Revised: August 31, 2020

Accepted: September 2, 2020

Online First: September 10, 2020

DOI: <https://doi.org/10.2298/SARH200203068M>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

†Correspondence to:

Vladimir MANOJLOVIĆ

University of Novi Sad, Faculty of Medicine,

Hajduk Veljkova 3 21000 Novi Sad, Serbia

Email: vladimir.manojlovic@mf.uns.ac.rs

Incomplete Circle of Willis and cerebrovascular reactivity in asymptomatic patients before and after carotid endarterectomy

Некомплетност Вилисовог прстена и цереброваскуларна реактивност код асимптоматских пацијената пре и после каротидне ендартеректомије

SUMMARY

Introduction/Objective Circle of Willis (CoW) provides the most significant collateral flow in the presence of significant stenosis or occlusion of internal carotid artery. In terms of collateral flow „incomplete“ type and „complete“ type of CoW can be recognized. Patients with carotid artery disease with incomplete CoW have lower cerebrovascular reactivity and higher risk for stroke. Cerebrovascular reactivity refers to the residual capacity of dilatation of cerebral blood vessels in the condition of insufficient blood flow. In this study we analyzed changes in cerebrovascular reactivity after carotid endarterectomy in asymptomatic patients with respect to complete and incomplete CoW morphology.

Methods In this study in 97 patients with asymptomatic carotid artery disease we measured cerebrovascular reactivity before and after carotid endarterectomy by using method of „apnea test“ and “ Breath Hold Index“ (BHI). Patients were divided into two following groups: patients with „complete“ CoW and „incomplete“ CoW based on non-contrast magnetic resonance angio performed previously to the operation. Descriptive statistics, univariate analysis, and ANOVA for comparison of BHI values between groups were used.

Results Results showed significant increase in cerebrovascular reactivity at the side of stenosis in both groups of patients with complete CoW (BHI value increased from 0,897 to 1,090; $F(1,65) = 30,788$, $p < 0,0005$, $parc. \eta^2 = 0,321$) and incomplete CoW (BHI value increased from 0,690 to 1,010; $F(1,27) = 62,318$, $p < 0,0005$, $parc. \eta^2 = 0,698$) and the more significant increase in the group of incomplete CoW compared to the group with complete CoW ($F(1,92) = 4,557$, $p = 0,035$, $parc. \eta^2 = 0,047$)

Conclusion In most asymptomatic patients, cerebrovascular reactivity restores to normal following carotid endarterectomy. Parameters of cerebrovascular reactivity are lower in patients with incomplete CoW and the increase after carotid endarterectomy is more significant in such patients.

Keywords: Breath holding index, extracranial carotid disease, internal cerebral artery, asymptomatic carotid patients risk stratification.

САЖЕТАК

Увод/Циљ Вилисов прстен представља најзначајнији колатерални пут којим се обезбедује проток у хемисферама мозга код екстракранијалне каротидне стенозе или оклузије. У погледу присуства колатералног протока разликујемо комплетни и некомплетни Вилисов прстен. Пацијенти са екстракранијалном каротидном болести који имају некомплетан Вилисов прстен имају слабију цереброваскуларну реактивност и повишен ризик можданог удара. Цереброваскуларна реактивност описује резидуални капацитет за дилатацију церебралних крвних судова у условима мождане хипоперфузије. Циљ ове студије био је да се утврди промене у цереброваскуларној реактивности након каротидне ендартеректомије код асимптоматских пацијената са комплетним и некомплетним Вилисовим прстеном.

Метод Студија је укључила 97 пацијената са асимптоматском каротидном екстракранијалном болести код којих смо одређивали цереброваскуларну реактивност методом „Апнеа теста“ пре и после каротидне ендартеректомије. Пацијенти су на основу налаза безконтрастне магнетне резонантне ангиографије били подељени у две групе: Пацијенти са комплетним Вилисовим прстеном и пацијенти са некомплетним Вилисовим прстеном. Статистичка анализа подразумевала је дескриптивну статистику, униваријатну анализу и ANOVA за поређење параметара цереброваскуларне реактивности пре и после каротидне ендартеректомије.

Резултати Резултати су показали значајно повећање параметара цереброваскуларне реактивности на страни стенозе у обе групе пацијената са комплетним Вилисовим прстеном (порао BHI са 0,897 на 1,090; $F(1,65) = 30,788$, $p < 0,0005$, $parc. \eta^2 = 0,321$) и некомплетним Вилисовим прстеном (порао BHI са 0,690 на 1,010; $F(1,27) = 62,318$, $p < 0,0005$, $parc. \eta^2 = 0,698$) при чему је пораст у групи са некомплетним Вилисовим прстеном био значајнији за $p = 0,035$ ($F(1,92) = 4,557$, $parc. \eta^2 = 0,047$).

Закључак Код већине асимптоматских пацијената са каротидном болести, каротидна ендартеректомија доводи до нормализације параметара цереброваскуларне реактивности. Цереброваскуларна реактивност је нижа код пацијената који имају некомплетан Вилисов прстен, а пораст цереброваскуларне реактивности након операције израженији је код ових пацијената.

Кључне речи: *Breath Hold Index*, Унутрашња каротидна артерија, екстракранијална каротидна болест, стратификација ризика код асимптоматских каротидних болесника

INTRODUCTION

Circle of Willis (CoW) provides the most significant collateral flow in the presence of significant stenosis or occlusion of internal carotid artery (ICA). Anterior collateral segment of CoW (ACA1, AcomA) is a connection between opposite carotid arteries and posterior collateral segment (ACP1, AcomP) provides collateral from posterior cerebral circulation [1].

Morphology of CoW can be evaluated by non-contrast enhanced magnetic resonance angiography (nCEMRA) and it depicts the functional status of collateral flow [2]. Although there is a number of CoW morphology types, in terms of collateral flow „incomplete“ type and „complete“ type of CoW can be recognized. Contrary to the „complete“ CoW that depicts normal CoW morphology, „incomplete“ CoW refers to the hypoplasia or occlusion of anterior and posterior collateral segment and consequent absents of collateral flow provided by CoW.

In the presence of significant ICA stenosis, incomplete CoW can be associated with impaired cerebral blood flow, reduction of cerebral autoregulation decreased circulatory reserve and low cerebrovascular reactivity leading to increased risk of stroke [3].

Cerebrovascular reactivity describes the capacity of adaptation of cerebral blood flow as a reaction to different stimuli. If insufficient cerebral blood flow is present, blood vessels are maximally dilated, and the residual capacity to increase blood flow is limited.

In this study we analyzed changes in cerebrovascular reactivity after carotid endarterectomy in asymptomatic patients with respect to complete and incomplete CoW morphology.

METHODS

Research included 97 patients, out of 171, who were operated from asymptomatic extracranial carotid stenosis in the period January 2017.-June 2019. Prior to the operation, all

patients underwent Duplex Ultrasound Examination (DUS) of carotid arteries and at least one of the following imaging: Magnetic Resonance Imaging and Angiography (MRI and MRA) (Siemens 1.5T) or Computed Tomography Imaging and Angiography (CT and CTA) (64 or 128 lines) of head and neck. MRI and MRA with three-dimensional time of flight (3D TOF) sequence for examination of extracranial portion of carotid and vertebral arteries and CoW and intracranial arteries was mandatory for patients who were included into the study.

Based on aforementioned diagnostic procedures, inclusion and exclusion criteria for this study were defined: patients with unilateral carotid disease (contralateral carotid stenosis was less than 50%) were included, with no significant lesions on intracranial portion of carotid arteries, vertebral and basilar arteries and cerebral arteries and no evidence of „silent brain infarctions“ larger than 1cm. Patients who were presented with insufficient data, poor insonation window for measurement of cerebrovascular reactivity or low compliance with the procedure, patients whose written consent was not p were also excluded from the study.

We collected preoperative data on patients' general characteristics, risk factors and comorbidity: age and gender, presence of hypertension, diabetes, smoking, hyperlipoproteinemia, history of ischemic heart disease or heart failure, left ventricle hypertrophy, significant heart valve diseases, atrial fibrillation, chronic kidney disease, chronic obstructive pulmonary disease and peripheral artery disease (PAD). Assessment of clinical cardiologist has been provided.

Morphology of CoW was determined based on 3D TOF sequence of MRA. By morphology, patients were classified into two groups:

1. Group of patients with „complete CoW“ in which all arteries of CoW were shown on 3D TOF sequence MRA (67 patients – 69%)

2. Group of patients with incomplete CoW: underdeveloped or occluded anterior (ACA1, AcomA) and posterior (ACP1, AcomP) collateral segment of CoW (30 patients - 31%)

Degree of ICA stenosis was estimated by DUS based on ECST criteria and two groups of patients were recognized: group of patients with 75%-84% ICA stenosis and group of patients with 85%-99% ICA stenosis.

All patients underwent operation under general anesthesia; one of the following techniques was used: Carotid Endarterectomy (CEA) with patch plasty and intraluminal shunt protection or Eversion endarterectomy (EEA). Standardized perioperative protocol included administration of clopidogrel 75mg at least five days prior to the operation with no discontinuation for the operation day, administration of statins starting at least 30 days before the operation. In the postoperative period, strict blood pressure control was indicated, with the aim to maintain systolic blood pressure below 160mmHg.

For estimation of cerebrovascular reactivity we used „Apnea test“ method, previously described by M. Silvestrini et al. [4]. In Apnea test patients are asked to hold breath for 30 seconds, and consequent increase in blood CO₂ is used as a stimulus for dilatation of cerebral blood vessels. The increase in blood flow during apnea test is registered with Transcranial Doppler ultrasound (TCD) (Multidop X4, DWL Elektronische Systeme GmbH with 2 MHz probe and Sonara Viasis version 04 with 2MHz probe) on proximal portion of Middle Cerebral Artery (MCA). As a result of the test the Breath Hold Index (BHI) is calculated as ratio between stimulated (CBF_{30 sec}) and basal cerebral blood flow (CBF_{basal}):

$$BHI = ((CBF_{30 \text{ sec}} - CBF_{\text{basal}}) / CBF_{\text{basal}}) / (\text{sec} / 100).$$

„Cut-off“ point for normal finding was set on 0.69. In this research apnea test was done to all patients one day before and one month after surgery.

We compared BHI values before and after surgery in groups of patients with complete and incomplete CoW for both sides: ipsilateral and contralateral to stenosis.

Statistical analysis included descriptive statistics: mean value, frequency (count) and relative frequency (percentage) for categorical data; comparative statistics included univariate analysis of variables with odd's ratio calculation; differences between BHI values before and after surgery in groups of patients with complete and incomplete CoW for both sides: ipsilateral and contralateral to stenosis has been analyzed by ANOVA . SPSS Statistics ver. 25 IBM Inc. Chicago, IL, USA was used.

Subjects' written consents have been obtained. All studies have been approved by the ethics committee of the Clinical centre of Vojvodina and the Medical faculty of the University in Novi Sad, and conforms to the legal standards. In most of the patients, MR examinations were done as a part of the project "Registration of New Ischemic Lesions with Magnetic Resonance Imaging Before and After Carotid Endarterectomy and Carotid Stenting", financed by the Provincial Secretariat for Higher Education and Scientific Research of Vojvodina. Subject's written consent has been approved by ethics Committee of Clinical center of Vojvodina and Medical Faculty University in Novi Sad and conforms to the legal standards.

RESULTS

A total group of 97 asymptomatic patients with ICA stenosis, 75 males, and 24 females, aged 54 to 79 years old, median value 66.33, underwent carotid endarterectomy due to extracranial carotid disease. Both techniques of carotid endarterectomy with patch angioplasty and intraluminal protection (44%) and eversion endarterectomy (54%) were used depending on preferences of surgeon performing the operation, all under general anesthesia. We did not register any major perioperative adverse event (stroke or death) and we registered

one postoperative case of acute coronary syndrome that was successfully treated by percutaneous coronary angioplasty. Patients' general characteristics and comorbidities with respect to complete and incomplete CoW morphology are shown on **Table 1**.

In both groups of patients with complete CoW and incomplete CoW degree of ICA stenosis (patients were classified into groups 75–84% and 85–99% stenosis) was equally distributed as it is shown on **Table 2**.

Preoperative values of BHI at the side of stenosis were 0.897 for the group of patients with complete CoW and 0.617 in the group of patients with incomplete CoW. In 34% of patients with complete CoW and 70% of patients with incomplete CoW preoperative BHI value were lower than previously defined „cut-off“ value of 0.69. After the operation BHI values were 1.09 for group of patients with complete CoW and 1.01 in group of patients with incomplete CoW. Only 5.8% of patients with complete CoW and 6.7 % with incomplete CoW had postoperative BHI value less than 0.69.

Observing BHI values lower than 0.69 as pathological, we registered significant reduction of number of pathological findings of BHI level after the operation in the group of patients with incomplete CoW (63.3%) compared to the group of patients with complete CoW (28,2%) and the difference was statistically significant ($p=0.0016$, OR= 4.36 CI 1.75-10.78) **Figures. 1a and 1b**.

By using ANOVA we tested primary effects of two following variables: morphology of CoW (complete and incomplete CoW) and operation status („before and after surgery“) on BHI values. It was determined that there was a significant difference in BHI values before the surgery between groups of patients with complete and incomplete CoW, $F(1,94) = 16.208$, $p < 0.001$, $\text{parc. } \eta^2 = 0.150$ and significant difference in BHI values between same groups after the surgery, $F(1,94) = 4.134$, $p < 0.05$, $\text{parc. } \eta^2 = 0.043$. We found significant influence of carotid endarterectomy on the BHI values in the group of patients with incomplete CoW, $F($

1.27) = 62.318, $p < 0.0005$, $\text{parc. } \eta^2 = 0.698$ and in the group of patients with complete CoW, $F(1.65) = 30.788$, $p < 0.0005$, $\text{parc. } \eta^2 = 0.321$. **Figure 2a**

By using ANOVA we tested interaction effect of following variables: morphology of CoW (complete and incomplete CoW) and operation status (before and after surgery) on BHI values. There was a significant interaction between the completeness of the CoW and operation status and their impact on the level of BHI, $F(1.92) = 4.557$, $p = 0.035$, $\text{parc. } \eta^2 = 0.047$. More significant increase in BHI value was registered in group of patients with incomplete CoW after the operation compared to patients with complete CoW. The difference proved to be statistically significant for $p = 0.035$. Therefore carotid endarterectomy more affected patients with incomplete CoW in terms of improvement of cerebrovascular reactivity. **Figure 2a**

For the side opposite to stenosis ANOVA showed significant difference in BHI value before and after operation in both groups ($F = 7.357$, $p = 0.008$, $\text{parc. } \eta^2 = 0.072$), but no difference between the groups ($F = 0.831$, $p = 0.34$, $\text{parc. } \eta^2 = 0.009$). **Figure 2b**

DISCUSSION

In asymptomatic significant ICA stenosis revascularization is indicated only in low risk patients who feature increased risk of stroke [5]. In this respect, investigation of cerebrovascular reactivity in asymptomatic patients has been recognized in up-to-date guidelines in preoperative assessment and risk stratification of patients with carotid artery stenosis [6]. Low cerebrovascular reactivity means that cerebral arteries are already dilated to their's maximum due to low cerebral perfusion and there is a limited reserve of adaptation of cerebral flow. In asymptomatic patients with significant carotid artery stenosis reduced cerebrovascular reactivity increases risk of stroke 13-25% per year [7-11]. Decreased circulatory reserve and lack of collateralization may increase the risk of stroke by the

mechanism of impaired hemodynamics and due to the fact that arterio-arterial embolization from diseased carotid artery occurs more often in the zone of reduced circulatory reserve [12, 13].

Association of incomplete CoW finding on nEMRA and low cerebrovascular reserve has been documented as well [14]. nCEMRA that was used in our research, represents functional morphology of CoW as it displays only blood flow within the vessels [2]. Although numerous types of CoW morphology have been described, simplification to „complete“ and „incomplete“ CoW has been accepted for easier use in clinical practice [15]. It is evidenced that 25-30% asymptomatic patients and 45-60% of symptomatic patients with carotid artery disease have incomplete CoW [16]. In symptomatic patients with significant ICA stenosis increased annual risk of stroke up to 13-17% is evidenced if the incomplete CoW is present [17, 18]. For asymptomatic patients with ICA stenosis there is a lack of data from controlled prospective studies [19]. Retrospective post-hoc analyzes of a SMART group showed increased but not statistically significant risk of stroke in patients with „incomplete“ CoW [14].

Our study showed that in patients with incomplete CoW, circulatory reserve at the side of ICA stenosis was significantly lower (median BHI = 0.62) compared to the patients whose MRA findings showed the complete CoW (BHI median = 0.88). As well BHI at the side of ICA stenosis was lower compared to the side opposite to the stenosis (BHI median = 1, 09). BHI in the group of patients with incomplete CoW tended to be lower than the proposed cut-off value for normal findings which is 0.69 [4].

Operative treatment resulted in the significant increase in BHI at the side of the stenosis both in groups of patients with complete and incomplete CoW. We registered both significant improvement circulatory reserve and normalization of the findings in the majority of patients in which BHI was below the threshold of 0.69. Such effect indicates that the

revascularization of stenosed ICA removes the cause of impaired circulatory reserve and reduced vasomotor reactivity. More beneficial effect of surgical treatment we found to be in asymptomatic patients with incomplete CoW with more significant increase of BHI. For the opposed side we found trend of greater postoperative increase in BHI value in the group with complete CoW, which can be explained by the phenomenon of "steal" from the healthy side over active collaterals that was present before the operation.

The literature emphasizes the importance of the effect of carotid endarterectomy on patients with extremely low parameters of cerebral vasoreactivity [20, 21]. Silvestrini and Soine founded beneficial effect of surgery only in symptomatic but not in asymptomatic patients [22]. In aforementioned research the asymptomatic patients were not stratified according to CoW morphology. A significant improvement of cerebrovascular reactivity after carotid endarterectomy in asymptomatic patients can be registered for both sides of brain [23,24]. Surgical treatment of asymptomatic and symptomatic patients is followed by normalization of cerebrovascular reactivity and collateral flow in the CoW [25].

Improvement of cognitive function after carotid endarterectomy along with the improvement of cerebrovascular reactivity is emphasized [26]. Previously mentioned SMART study group was one of the rare studies that followed operated and non-operated asymptomatic patients with complete and incomplete CoW, still it was the retrospective study [14].

Apnea test and its modifications are easy available and can be done in most vascular labs, it is also proved to be comparable to other methods of measurement of cerebrovascular reactivity [27]. Still there is a problem of its reliability especially in patients who are of low compliance with the procedure, which is recognized as a limitation of this study. Association of incomplete CoW and low cerebrovascular reserve is evident, as well as effect of ICA revascularizations on cerebrovascular reactivity, but should presence of incomplete CoW can be observed as risk feature in asymptomatic ICA stenosis is still to be debated.

CONCLUSIONS

In most asymptomatic patients cerebrovascular reactivity restores to normal following carotid endarterectomy. Parameters of cerebrovascular reactivity are lower in patients with incomplete CoW and the increase after carotid endarterectomy is more significant in such patients. This suggests that carotid endarterectomy is more beneficial in asymptomatic patients with incomplete CoW in terms of cerebrovascular reactivity, but does it indicate clinical benefit in such patients (i.e. reduction of the risk of stroke) is yet to be approved by future prospective studies.

ACKNOWLEDGEMENTS

Acknowledgements to Prof dr Petar Slankamenac and Doc dr Aleksandra Lučić Prokin for providing equipment and for expert support.

Conflict of interest: None declared.

REFERENCES

1. Fang H, Song B, Cheng B, Wong KS, Xu YM, Ho SS, et al. Compensatory patterns of collateral flow in stroke patients with unilateral and bilateral carotid stenosis. *BMC Neurol.* 2016;16:39. PMID: 26987874; DOI: 10.1186/s12883-016-0560-0
2. Li J, Wang J, Wei X, Hao YW, Wang F, Li Y. Examination of Structural Variations of the Circle of Willis by 3D Time-of-Flight Magnetic Resonance. *Front. Neurosci.* Feb 2020; DOI:10.3389/fnins.2020.00071
3. Malhotra K, Liebeskind DS. Collaterals in ischemic stroke. *Brain Hemorrhages* 2020;1: 6-12 DOI: 10.1016/j.hest.2019.12.003
4. Silvestrini M, Troisi E, Matteis M, Cupini LM, Caltagirone C. Transcranial Doppler Assessment of Cerebrovascular Reactivity in Symptomatic and Asymptomatic Severe Carotid Stenosis. *Stroke.* 1996;27:1970-73
5. Nicolaides A, Sabetai M, Kakkos SK, Dhanjil S, Tegos T, Stevens JM, et al. ACSRS Study Group. Severity of asymptomatic carotid stenosis and risk of ipsilateral hemispheric ischaemic events: results from the ACSRS study. *Eur J Vasc Endovasc Surg.* 2005;30(3):275-84. PMID: 16130207; DOI: 10.1016/j.ejvs.2005.04.031
6. Naylor AR, Ricco JB, de Borst GJ, Debus S, de Haro J, Halliday A et al. Editor's Choice – Management of Atherosclerotic Carotid and Vertebral Artery Disease: 2017 Clinical Practice Guidelines of the European Society for Vascular Surgery (ESVS). *EJVS.* 2018;55(1): 3–81. PMID: 28851594; DOI: 10.1016/j.ejvs.2017.06.021
7. Lin T, Lai Z, Zuo Z et al. ASL perfusion features and type of circle of Willis as imaging markers for cerebral hyperperfusion after carotid revascularization: a preliminary study. *Eur Radiol* 2019;29(5):2651-2658.
8. Reinhard M, Schwarzer G, Briel M, Altamura C, Palazzo P, King A, et al. Cerebrovascular reactivity predicts stroke in high-grade carotid artery disease. *Neurology.* 2014;83(16):1424–31. PMID: 25217057; DOI: 10.1212/WNL.0000000000000888
9. King A, Serena J, Bornstein NM, Markus HS, ACES Investigators. Does impaired cerebrovascular reactivity predict stroke risk in asymptomatic carotid stenosis? A prospective substudy of the asymptomatic carotid emboli study. *Stroke.* 2011;42:1550–1555. PMID: 21527764 DOI: 10.1161/STROKEAHA.110.609057
10. Ju K, Zhong L, Ni X, Cao H, Cheng G, Ding L. Cerebral vasomotor reactivity predicts the development of acute stroke in patients with internal carotid artery stenosis. *Neurologia i Neurochirurgia Polska* 2018; 52(3):374-8. DOI:10.1016/j.pjnns.2017.12.015
11. Gupta A, Chazen JL, Hartman M, Delgado D, Anumula N, Shao H, et al. Cerebrovascular reserve and stroke risk in patients with carotid stenosis or occlusion: a systematic review and meta-analysis. *Stroke.* 2012;43(11):2884-91. PMID: 23091119; DOI: 10.1161/STROKEAHA.112.663716
12. Zhou C, Yuan C, Li R, Wang W, Li C, Zhao X, and on behalf of the CARE-II Study Collaborators. Association Between Incomplete Circle of Willis and Carotid Vulnerable Atherosclerotic Plaques-A Chinese Atherosclerosis Risk Evaluation Study. *ATVB.* 2018;38:2744–2749. DOI:10.1161/ATVBAHA.118.311797
13. Fang H, Song B, Cheng B et al. Compensatory patterns of collateral flow in stroke patients with unilateral and bilateral carotid stenosis. *BMC Neurol.* 2016;16:39.
14. Hong D, Hen Lee Y, Seo HS, Je BK, Suh S, Seol HY et al. Correlation of Cerebrovascular Reserve as Measured by Acetazolamide-Challenged Perfusion CT with Collateral Circulation in Unilateral High Grade Carotid Stenosis. *J Korean Soc Radiol.* 2014;70(1):1-12. DOI:10.3348/jksr.2014.70.1.1.
15. van Seeters T, Hendrikse J, Biessels GJ, Birgitta K, Velthuis BK, Mali W, Kappelle LJ, van der Graaf Y and SMART Study Group. Completeness of the circle of Willis and risk of ischemic stroke in patients without cerebrovascular disease. *Neuroradiology.* 2015;57(12):1247–51. PMID: 26358136 PMID: DOI: 10.1007/s00234-015-1589-2
16. Manojlović V, Popović V, Nikolić D, Milošević Đ, Pasternak J, Budakov N. Completeness of Circle of Willis in symptomatic and asymptomatic extracranial carotid disease. *Med Pregl* 2016; 69 (11-12):335-416. DOI: 0.2298/MPNS1612351M

17. Pascalau R, Paudrean VA, Bartos D et al. The Geometry of the Circle of Willis Anatomical Variants as a Potential Cerebrovascular Risk Factor. *Turk Neurosurg.* 2019;29(2):151-158.
18. Hoksbergen AW, Legemate DA, Csiba L, Csati G, Siro P, Fulesdi B. Absent collateral function of the circle of Willis as risk factor for ischemic stroke. *Cerebrovasc.Dis.* 2003;16:191-198. PMID: 12865604; DOI: 10.1159/000071115
19. King A, Serena J, Bornstein NM, Markus HM. On behalf of the ACES Investigators. Does impaired cerebrovascular reactivity predict stroke risk in asymptomatic carotid stenosis: a prospective substudy of the asymptomatic carotid emboli study. *Stroke* 2011;42:1550e5. PMID: 25217057; DOI: 10.1212/WNL.0000000000000888
20. Banga PV, Varga V, Csobay-Novák C, Kolossváry M, Szántó E, Oderich GS, Entz L, Sótónyi P. Incomplete circle of Willis is associated with a higher incidence of neurologic events during carotid eversion endarterectomy without shunting. *J Vasc Surg.* 2018 Dec;68(6):1764-1771. DOI: 10.1016/j.jvs.2018.03.429.
21. Lučić Prokin A, Slankamenac P, Kovačević P, Ružička Kaloci S, Živanović Ž. Cerebral Vasomotor Reactivity and Apnea Test in Symptomatic and Asymptomatic High-Grade Carotid Stenosis. *Srp Arh Celok Lek.* 2015;143(9-10):520-524. DOI: 10.2298/SARH1510520L
22. Soenne L, Helenius J, Tatlisumak T, Saimanen E, Salonen O, Lindsberg PJ, et al. Cerebral hemodynamics in asymptomatic and symptomatic patients with high-grade carotid stenosis undergoing carotid endarterectomy. *Stroke.* 2003;34(7):1655-61. PMID: 12805504; DOI: 10.1161/01.STR.0000075605.36068.D9
23. Barzo P, Voros E, Bodosi M. Use of transcranial Doppler sonography and acetazolamide test to demonstrate changes in cerebrovascular reserve capacity following carotid endarterectomy. *Eur J Vasc Endovasc Surg.* 1996;11(1):83-9. PMID: 8564493; DOI: 10.1016/s1078-5884(96)80140-7
24. Saher El-Sayed H, Ehab Shaker B, Montaser MH, Ahmed AEB, Sandra MA, Islam IZ. Cerebrovascular reactivity recovery following carotid angioplasty and stenting. *The Egyptian Journal of Neurology, Psychiatry and Neurosurgery.* 2019;55:32.
25. Telman G, Kouperberg E, Nitecki S, Karram T, Schwarz HA, Sprecher E, et al. Cerebral hemodynamics in symptomatic and asymptomatic patients with severe unilateral carotid stenosis before and after carotid endarterectomy. *EJVES.* 2006;32(4):375-8. PMID: 16395290; DOI: 10.1038/sj.jcbfm.9600260
26. Lattanzi S, Carbonari L, Pagliariccio G, Bartolini M, Cagnetti C, Viticchi G, et al. Neurocognitive functioning and cerebrovascular reactivity after carotid endarterectomy. *Neurology* 2018; 90(4) DOI:10.1212/WNL.0000000000004862
27. Alwatban M, Truemper E, Al-Rethaia A, Murman D, Bashford G. The Breath-Hold Acceleration Index: A New Method to Evaluate Cerebrovascular Reactivity using Transcranial Doppler. 2018 Jul;28(4):429-435. doi: 10.1111/jon.12508.

Table 1. Distribution of patients' general characteristics, risk factors, and comorbidity in the groups with complete and incomplete Circle of Willis (CoW)


Variables	Complete CoW	Incomplete CoW	total	OR	p value
Circle of Willis (No)	70	27	97	-	-
Age (median value)	67.9	65,9	66.33	-	0.465
Males (%)	78%	76%	76%	1.105	0.8734
Hypertension (%)	83%	87%	86%	0.792	0.811
Diabetes mellitus (%)	38%	36%	36.5%	1.131	0.819
Smoking (%)	50%	53%	51%	1.077	0.870
Hyperlipoproteinemia(%)	58%	73%	62%	0.882	0.759
Ishaemic heart disease or heart failure (%)	27%	26%	27%	0.939	0.903
Left ventricle hypertrophy (%)	22%	23%	22%	1.047	0.932
Significant heart valve disease (%)	0%	1%	1%	7.981	0.207
Atrial fibrillation or other arythmia (%)	11%	15%	12%	1.291	0.697
Chronic kidney disease (%)	4.2%	3.7%	4.1%	0.827	0.871
Chronic obstructive pulmonary disease (%)	8.5%	11%	9%	1.333	0.700
Peripheral artery disease (%)	13%	7%	11%	0.629	0.562

Table 2. Degree of Internal Carotid Artery (ICA) stenosis by ECST criteria estimated by duplex ultrasound in the groups of patients with complete and incomplete Circle of Willis (CoW, patients were classified into groups 75–84% and 85–99% stenosis)

CoW morphology	Degree of stenosis		Total	p value
	75–84%	85–99%		
Incomplete CoW	21	9	30	0,78
Complete CoW	45	22	68	
Total	66	31	97	

Paper accepted

a)



b)





Figure 1. Breath Hold Index values before and after operation in asymptomatic patients with a) incomplete Circle of Willis (CoW) b) and complete CoW

a)



b)




Figure 2. Estimated marginal means of Breath Hold Index before and after operation in asymptomatic patients with complete and incomplete Circle of Willis (CoW) at the a) operated side; b) contralateral side.