



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Jovana Milošević^{1†}, Marija Milić¹, Momčilo Mirković¹, Nenad Milošević^{1,2},
Tatjana Novaković^{1,2}, Zdravko Vitošević¹, Slađana Đurić¹, Mirjana Stojanović Tasić^{1,2},
Ljiljana Kulić¹

**Association between non-communicable diseases and satisfaction with
healthcare and self-rated health: experiences from
post-conflict communities**

Повезаност хроничних незаразних болести са задовољством здравственом
заштитом и самопроценом здравственог стања: искуства из
постконфликтне средине

¹University of Priština – Kosovska Mitrovica, Faculty of Medicine, Kosovska Mitrovica, Serbia;

²Clinical hospital Center of Priština, Gračanica, Serbia

Received: October 26, 2019

Revised: May 21, 2020

Accepted: September 22, 2020

Online First: October 1, 2020

DOI: <https://doi.org/10.2298/SARH191026086M>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

†Correspondence to:

Jovana MILOŠEVIĆ

Faculty of Medicine, Institute of Preventive Medicine, University of Priština – Kosovska Mitrovica,

Anri Dinana bb, 38220 Kosovska Mitrovica, Serbia

E-mail: jovana_mil@yahoo.com

Association between non-communicable diseases and satisfaction with healthcare and self-rated health: experiences from post-conflict communities

Повезаност хроничних незаразних болести са задовољством здравственом заштитом и самопроценом здравственог стања: искуства из постконфликтне средине

SUMMARY

Introduction/Objective Estimating the prevalence of non-communicable diseases (NCD), multimorbidities, and their association with self-rated health as well as satisfaction with healthcare.

Methods This cross-sectional study was conducted among ethnic Serb communities at Kosovo and Metohija during 2015–2016. Data of sociodemographic and lifestyle characteristics, self-rated health status and satisfaction with healthcare was obtained through a survey which included 1 067 adults, 535 of whom reported presence of non-communicable disease. Multinomial regression was performed to analysis factors associated with self-rated health and self-rated satisfaction with the healthcare.

Results Presence of one NCD was reported by 50.1% respondents, whereas 23.1% of the respondents reported multimorbidity. While self-reported NCD presence was negatively associated with self-rated health ($p: 0.001–0.016$), no association between NCDs and satisfaction with healthcare was observed ($p: 0.178–0.974$). Being single ($p: 0.011–0.017$), lower educational level ($p: 0.031–0.047$), regular breakfast ($p = 0.032$), frequent vegetable intake ($p: 0.009–0.029$), no alcohol use ($p = 0.010$), shorter waiting time ($p: 0.001–0.004$) and sufficient finance for dental care ($p = 0.021$) were factors statistically significantly correlated with greater satisfaction with the healthcare.

Conclusion Presence of non-communicable diseases was negatively associated with self-rated health status, while shorter waiting time and adequate finances were associated with higher level of satisfaction with the healthcare. The results of our study could be of the importance for policy makers in creating the more effective healthcare service in unstable political and security situations.

Keywords: chronic diseases, health status, adult population

САЖЕТАК

Увод/Циљ Испитивање учесталости хроничних незаразних болести (ХНБ) и мултиморбидитета као и њихове повезаности са самопроценом здравственог стања и задовољством здравственом заштитом.

Метод Студија пресека спроведена је у српским срединама на територији Косова и Метохије током 2015/2016 године. Укупно је анкетирано 1067 одраслих становника коришћењем упитника о социодемографским карактеристикама, животним навикама, самопроцени здравственог стања и задовољства здравственом заштитом, од којих је 535 пријавило присуство неке од ХНБ. Мултиномијална регресија спроведена је за анализу фактора повезаних са задовољством здравственом заштитом и самопроценом здравственог стања.

Резултати: Присуство ХНБ је пријављено од стране 50.1% испитаника, док је присуство мултиморбидитета пријавило 23.1% испитаника. Присуство ХНБ је показало негативну повезаност са самопроценом здравственог стања ($p: 0.001–0.016$), док повезаност ХНБ са задовољством здравственом заштитом није уочена ($p: 0.178–0.974$). Бити без емотивног партнера ($p: 0.011–0.017$), ниже образовање ($p: 0.031–0.047$), редован доручак ($p = 0.032$), редовно конзумирање поврћа ($p: 0.009–0.029$), некоришћење алкохола ($p = 0.010$), краће време чекања ($p: 0.001–0.004$) и поседовање довољно финансија за остваривање стоматолошке здравствене заштите ($p = 0.021$) су показали повезаност са већим задовољством здравственом заштитом.

Закључак Присуство ХНБ је показало негативну повезаност са самопроценом здравственог стања, док су краће време чекања и финансије били повезани са већим задовољством здравственом заштитом. Резултати нашег истраживања могу бити од значаја креаторима здравствене политике приликом доношења одлука у правцу стварања што ефикасније здравствене службе.

Кључне речи: хроничне болести, здравствено стање, одрасло становништво

INTRODUCTION

In the last two decades, global burden of disease expressed as disability-adjusted life year (DALY) has increased mainly due to non-communicable diseases (NCDs) [1, 2].

Population ageing and unhealthy lifestyle are considered to be the main factors contributing to the NCD development [3]. In the primary healthcare setting, especially in high-income

countries, presence of two or more chronic diseases (multimorbidity) is increasingly accepted as the norm rather than exception [4]. Presence of NCD and multimorbidity is related with shorter life expectancy, compromised mental health, frequent hospitalization and overall deterioration in life quality, which places a significant burden on the healthcare system [5].

Reliable data on the NCD and multimorbidity frequency are necessary for optimal healthcare provision and resource management [6]. Owing to their nature, NCDs are often considered a significant predictor of self-rated health and satisfaction with health care [7, 8]. On the other hand, self-rated health status and satisfaction with health care are important indicators of the healthcare system quality and can guide healthcare policy [9].

Studies have shown that residing in post-conflict areas increases the NCD risk, leads to worsening of existing NCDs and consequently to an increase in disability and death [10, 11, 12]. Post-conflict countries usually go through a protracted transition process, which typically includes rapid urbanization, due to which social systems tend to deteriorate [10, 13]. In such a situation, long-term stress exposure leads to mental health problems, unhealthy habits impair physical health and all of that represent ideal conditions for increase in NCD prevalence in the future [10, 13]. Thus, the aims of our study were to assess the prevalence of NCDs and multimorbidities among the population residing in Autonomous Province of Kosovo and Metohija (KM) and to determine their association with self-rated health and satisfaction with healthcare.

METHODS

Study Setting

This cross-sectional study focused on ethnic Serb communities residing on the KM. Data collection was conducted from October 2015 to January 2016. Approval for this study was obtained from the Ethics Committee of the Faculty of Medicine University of Pristina

temporarily settled in Kosovska Mitrovica (approval no. 67-11, issued on May 27, 2014).

Following the ethnic conflict in 1999, the Serbian KM province was segregated into a smaller predominantly Serb territory north of the Ibar River (composed of four municipalities) and a much larger Albanian territory to the south. Ethnic Serb population, however, also resides in six isolated municipalities in the predominantly Albanian southern part of KM, which are (with certain restrictions) still subject to Republic of Serbia (RS) governance [14]. Owing to these political conditions, many social systems and infrastructure services are compromised, including healthcare provision [15]. Given the limited presence of RS institutions on the KM, local population has not been included in population surveys conducted in the RS in the last 20 years [16, 17].

Study Participants

Respondents of interest for this study were adult (aged 18 and above) residents of four KM municipalities, two of which (Kosovska Mitrovica and Zubin Potok) are located north of the Ibar River, while the remaining two (Grasanica and Strpce) are in the southern part of KM.

Households were randomly selected for inclusion in the survey through stratification and two-phase sampling. Four of the ten municipalities with predominantly ethnic Serb population were designated as primary strata, whereby local communities served as sampling units in the first phase of the two-phase sampling process. From each stratum, 50% of local communities were selected, by choosing all odd-numbered local communities from the randomly generated list. In the second phase, households served as sampling units, whereby the units of analysis were identified as study participants from the households.

Out of a total of 400 households, 365 agreed to participate in the study with response household rate of 91.25%. At the end 1067 adults were interviewed.

Data Collection Instrument

Data required to meet the study objectives were collected via a purposefully-developed questionnaire designed in accordance with the European Health Interview Survey (EHIS wave 2), Methodological Manual [18].

Socio-demographic variables included sex, age, residential status (North/South KM), educational level (≤ 8 years/8–12/ > 12 years), employment status (unemployed/employed) and marital status (married/single). Lifestyle variables included daily habits, such as regular breakfast consumption (daily/occasionally), fruit and vegetable intake (rarely or never/weekly/daily), substance abuse (yes/no), smoking (yes/no), alcohol consumption (yes/no), excessive drinking, i.e., consumption of more than six alcoholic beverages on a single occasion (never or rarely/monthly/weekly/daily). Physical activity (active/inactive) was defined as continuous moderate physical activity of at least 10-minute per day that results in increased respiration or elevated heart rate [18, 19].

Presence of 17 predefined NCDs in the preceding 12 months was self-reported by the participants. Based on the number of reported NCDs, the sample was segregated into three groups: no NCDs/one NCD/multimorbidity.

The Likert scale was used to estimate the self-rated health (good/very good/moderate/poor/very poor) and self-rated satisfaction (very satisfied/satisfied/neutral/unsatisfied/very unsatisfied). Further, respondents' self-rated health status were stratified into three categories for the purpose of the study analyses: good or very good/moderate/poor or very poor. Similarly, self-rated satisfaction with the healthcare was grouped under: very satisfied or satisfied/neutral/unsatisfied or very unsatisfied.

Unmet health care needs are defined as the difference between the health services that are considered necessary to deal appropriately with health problems and services that are actually received [20]. Unmet healthcare needs were classified in accordance with the factors

that contribute to not receiving adequate health services (long waiting time, distance to the healthcare facility or transport issues, and inadequate finances for healthcare, dental care, mental care and medical prescription). Unmet health needs were assessed through closed-ended questions in which respondents have chosen between the offered alternatives (I did not need it/No/Yes).

Statistical Analyses

Statistical analyses were conducted using SPSS statistical package for MS Windows ver. 17 [21], where $p < 0.05$ was considered as statistically significant. Descriptive sample statistics included measures of central tendency (mean), variance (standard deviation) and relative values (percentages). Differences between studied parameters were assessed via the Kruskal-Wallis test (KW χ^2).

Multinomial regression analysis was performed when analyzing factors associated with self-rated health and self-rated satisfaction with the healthcare, which were treated as dependent variables. When examining self-rated health status and healthcare system quality, “poor or very poor” and “unsatisfied or very unsatisfied,” respectively, served as referent categories.

RESULTS

The study sample comprised of 1067 respondents aged 42.6 ± 16.0 years and 51.3% of them were female. In Table 1 are summarized participants' socio-demographic characteristics.

Hypertension (24.6%) was the most prevalent NCD in the study sample. The prevalences of other NCDs were: hyperlipidemia (10.3%), allergy (9.9%), deformity of the lower spine (8.1%), renal problems (5.9%), deformity of the upper spine (5.6%), diabetes

(5.2%), depression (5%), angina pectoris (4.1%), arthrosis (3.7%), urinary incontinence (3.1%), chronic bronchitis (2.4%), asthma (1.9%), myocardial infarction (1.4%), cancer (1.3%), stroke (1%) and cirrhosis (0.5%). Moreover, relative to men, a statistically significantly higher percentage of women reported suffering from chronic bronchitis ($p=0.024$), arthrosis ($p=0.001$), skeletal deformities ($p=0.001$) and depression ($p=0.002$) (Data not shown).

One-half of our participants (50.1%) reported that they have one NCD, whereas multimorbidity was declared by 23.1% of the participants (Table 1). Greater multimorbidity frequency was observed among older individuals, women, residents of south municipalities, less educated and unemployed respondents, and those in a relationship. Individuals with multimorbidity more often declared that they did not practice regular physical activity, and were more frequently to be smokers, but did not consume drugs or alcohol (Table 1).

Table 2 described self-rated health status and satisfaction with healthcare. Majority (71.2%) of the respondents rated their health as good or very good. As expected, those with multimorbidity were more frequent to perceive their health as moderate, poor, or very poor. On the other hand, those that did not report any NCDs were least satisfied with the healthcare ($p=0.029$).

Unmet healthcare needs due to long waiting time, distance to health facility or inadequate finances for healthcare, dental care, mental care and medical prescription were more frequently reported by participants with multimorbidities ($p=0.001$) (Table 2).

Table 3 shows factors associated with self-reported health status. Based on the multinomial regression analysis findings ($\chi^2=536.093$; $p=0.001$), absence or smaller number of NCDs, younger age, being employed, and alcohol consumption emerged as the factors most closely associated with positive self-rating health.

Multinomial regression analysis was also performed to assess the association between

presence of NCDs and satisfaction with healthcare. Both models developed for this purpose were statistically significant in explaining this relationship (“General socio-demographic” $\chi^2=108.080$, $p= 0.001$; “Healthcare unavailability” $\chi^2=92.383$; $p= 0.001$). Multinomial regression results failed to reveal any connection between NCD presence and satisfaction with the healthcare.

Being single, lower educational level, healthier lifestyle (regular breakfast, frequent vegetable intake and no alcohol use), shorter waiting time, and sufficient financial means for meeting the dental care needs were factors associated with higher level of satisfaction with the healthcare. Table 4 describes factors associated with healthcare satisfaction among respondents.

DISCUSSION

This study found that more than half of the survey respondents in post-conflict area suffered from at least one NCD. While presence of NCDs was negatively associated with health self-ratings among our participants, it was unrelated with participants’ satisfaction with healthcare.

Non-communicable diseases prevalence in the study sample was congruent with that reported for other Serbian territories [17]. The survey respondents were most frequent to report hypertension, which coincided with the findings of previous studies conducted in other countries [22, 23]. Comprehensive literature review revealed that armed conflicts tend to result in greater cardiovascular disease (CVD) incidence, as well as more frequent adoption of risky behaviors [10]. For example, Spiegel at al. reported that CVDs are more frequent cause of mortality in areas affected by interethnic conflict, compared to contagious diseases and conflict-related violence [24].

In a five-year study focusing on the 12 most common NCDs was found that

multimorbidity prevalence varies by the countries, with estimated range between 45% and 70% [25]. The lower multimorbidity prevalence found in this study relative to the results reported by other authors could be attributed to the nature of self-reported data, whereas in most previous studies prevalence was calculated based on data from official medical records [25]. Likewise, disparities in the multimorbidity prevalence for different countries are ascribed to incongruence in the utilized data sources, differences in target population characteristics and NCD types included in the analysis [22]. In the present study, respondents with multimorbidity more frequently reported unemployment, lower educational level and sedentary lifestyle. On the other hand, they more frequently adopted other healthy habits. It is possible that severely compromised health of persons living with multimorbidities precludes them from performing everyday physical activities, but following medical advice regarding healthy lifestyle and diet.

A higher percentage of participants in our study rated their health as very good and good compared to that of the previous study conducted throughout Serbia [17]. Our results coincide with those reported for European Union, where almost 70% of the surveyed population rated their health status as good [26]. In our sample, absence or a smaller number of NCDs, younger age, being employed and alcohol consumption were correlated with more positive self-rated health, which is in line with the findings reported in literature [27]. It is interesting to note that the survey respondents who consumed alcohol tended to rate their health more favorably. Given the anxiolytic effects of smoking and alcohol consumption, these habits are more prevalent among individuals living in post-conflict zones, as they serve as a coping mechanism [10, 28]. The studies have found that individuals who consume alcohol regularly in moderate quantities perceive their health status more favorably compared to those that abstain from alcohol or consume it occasionally [29].

In contrast to the health self-rated, presence of self-reported NCDs was not associated

with satisfaction with healthcare. Available evidence indicates that satisfaction with healthcare is determined by the educational level, income, age, and residence type [30]. Most of the studies examining patients' satisfaction with healthcare in the United States and European countries indicate that shorter waiting time, better-qualified medical personnel and lower healthcare cost are associated with more favorable healthcare system ratings [31, 32]. Our findings coincide with these observations. Available evidence further indicates that positive interactions with healthcare professionals correlate with greater satisfaction with the healthcare irrespective of the number and complexity of present diseases [33, 34]. In general, most Serbs residing within KM struggle to access required medical treatments and attribute these issues to inadequate healthcare system organization [16]. In sum, patient satisfaction is derived from a combination of prior experiences with medical services and perceived efficacy, safety and utility of available medical transportation [35].

The significance of the present study stems from the large sample size, thus ensuring its representativeness of ethnic Serb population residing on the KM. Consequently, the results reported in our study provide valuable insight into the NCD prevalence and perceived health status among individuals living in post-conflict zones, and their satisfaction with the healthcare, which is inevitably compromised due to previous conflict. Namely, previous studies from other countries affected by the conflict have found a higher burden of NCDs and mental illness among the population and exacerbations of existing NCDs due to serious psychological distress, less access to health care, interruptions in medication supply, lack of finances [11, 12, 23]. However, there are some limitations. Cross-sectional design adopted in the present study precluded us from reaching any conclusions regarding the causality of the observed relationships. Moreover, given that analyses were based on self-reported data that to a degree relied on accurate recall, informer bias could have affected the obtained results. Finally, our lifestyle measures were broad. For example, the measure of alcohol intake and

smoking has not been accurately quantified.

CONCLUSION

Based on our findings could be concluded that presence of NCDs is negatively associated with the self-rated health, but not with satisfaction with healthcare. Interestingly, being single, lower educational level, regular breakfast, frequent vegetable intake, no alcohol use, sufficient finance for dental care and shorter waiting time that mostly depends of organization of health service have positive influence on patients' satisfaction with healthcare. As patients' satisfaction with healthcare is one of the determinants of healthcare system quality, the findings reported in this work could be utilized when formulating healthcare policy, in particular when allocating inevitably limited resources, to ensure that they are most optimally utilized to provide as effective and efficient healthcare as possible.

ACKNOWLEDGEMENT

The paper is part of Jovana M. Milošević's doctoral thesis and presents some of the results that will be published in the thesis.

Conflict of interest: None declared.

REFERENCES

1. GBD 2016 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390(10100):1260-1344. DOI:10.1016/S0140-6736(17)32130-X
2. World Health Organization. Global status report on non-communicable diseases, 2014. Geneva: 2014. Available from: <https://www.who.int/nmh/publications/ncd-status-report-2014/en/>. [last assessed September 29, 2019]
3. Fabbri E., Zoli M., Gonzalez-Freire M., Salive M. E., Studenski S. A. & Ferrucci L. Aging and Multimorbidity: New Tasks, Priorities, and Frontiers for Integrated Gerontological and Clinical Research. *J Am Med Dir Assoc*. 2015; 16(8):640-647. DOI:10.1016/j.jamda.2015.03.013
4. Academy of Medical Sciences. Multimorbidity: a priority for global health research. 2018. <https://acmedsci.ac.uk/policy/policy-projects/multimorbidity> [last assessed January 22, 2020]
5. Atun R. Transitioning health system for morbidity. *Lancet*. 2015;386(9995):721-2. DOI: 10.1016/S0140-6736(14)62254-6.
6. Wallace E, Salisbury C, Guthrie B, Lewis C, Fahey T, Smith SM. Managing patients with multimorbidity in primary care. *Bmj*. 2015;350:h176. DOI: 10.1136/bmj.h176
7. Moffat K, Mercer SW. Challenges of managing people with multimorbidity in today's healthcare systems. *BMC Fam Pract* . 2015; 16:129. DOI: 10.1186/s12875-015-0344-4.
8. Gómez Ramírez OJ, Carrillo GM, Cárdenas DC. Survey on Satisfaction with Health care of People with Chronic Disease. *Enfermeria Global*. 2016;15(4):321-30. DOI: 10.6018/eglobal.15.4.226851
9. Mavaddat N, Valderas JM, van der Linde R, Khaw KT, Kinmonth AL. Association of self-rated health with multimorbidity, chronic disease and psychosocial factors in a large middle-aged and older cohort from general practice: a cross-sectional study. *BMC Fam Pract*. 2014; 15:185. DOI: 10.1186/s12875-014-0185-6
10. Jawad M, Vamos EP, Najim M, Roberts B, Millett C. Impact of armed conflict on cardiovascular disease risk: a systematic review. *Heart*. 2019;105(18):1388-94. DOI: 10.1136/heartjnl.
11. Greene-Cramer B, Summers A, Lopes-Cardozo B, Husain F, Couture A et al. Noncommunicable disease burden among conflict-affected adults in Ukraine: A cross-sectional study of prevalence, risk factors, and effect of conflict on severity of disease and access to care. *PLoS One*. 2020; 15(4):e0231899.
12. Slama S, Kim H-J, Roglic G, Boulle P, Hering H, Varghese C et al. Care of non-communicable diseases in emergencies. *The Lancet*. 2017; 389(10066), 326–330. DOI: [https://doi.org/10.1016/S0140-6736\(16\)31404-0](https://doi.org/10.1016/S0140-6736(16)31404-0)
13. Roberts B, Patel P, McKee M. Noncommunicable diseases and post-conflict countries. *Bull World Health Organ*. 2012;90(1):2, 2A. DOI: 10.2471/BLT.11.098863
14. Office for Kosovo and Metohija. Government of the Republic of Serbia. Available from: <http://www.kim.gov.rs/eng/index.php>. [last assessed September 08, 2019].
15. European Centre for Minority Issues Kosovo. Community profile: Serb community. Available from: <http://www.ecmikosovo.org/uploads/Serbcommunity1.pdf/>. [last assessed September 08, 2019].
16. Министарство здравља Републике Србије. Истраживање здравља Републике Србије: 2006. година: Основни резултати. Београд, 2007. Available from: <http://www.batut.org.rs/download/publikacije/Finalni%20izvestaj%202006.pdf> [last assessed September 19, 2019].
17. Институт за јавно здравље “Др Милан Јовановић Батут“. Резултати истраживања здравља становништва Србије 2013. година. Београд, 2014. Available from: <http://www.batut.org.rs/download/publikacije/IstrazivanjeZdravljaStanovnistvaRS2013.pdf/>. [last assessed September 19, 2019].
18. European Health Interview Survey (EHIS wave 2) – Methodological manual-2013 edition, Eurostat, 2013. Available from: <https://ec.europa.eu/eurostat/web/products-manuals-and-guidelines/-/KS-RA-13-018>. [last assessed September 09, 2019].
19. Physical activity. 2018. Available on: <https://www.who.int/news-room/fact-sheets/detail/physical-activity>.

20. Carr W, Wolfe S. Unmet Needs as Sociomedical Indicators. *International Journal of Health Services*. 1976;6 (3):417±430. DOI: 10.2190/MCG0-UH8D-0AG8-VFNU
21. SPSS Inc. Released 2008. *SPSS Statistics for Windows, Version 17.0*. Chicago: SPSS Inc.
22. Noncommunicable diseases country profiles 2018. Geneva: World Health Organization; 2018. Available from: <https://www.who.int/nmh/publications/ncd-profiles-2018/en/>. [last assessed September 09, 2019].
23. Mirković M, Simić S, Marinković J, Đurić Đ. Health state of the citizens of northern Kosovska Mitrovica. [Article in Serbian]. *Srp Arh Celok Lek*. 2010;138:746-51. DOI: 10.2298/SARH1012746M
24. Spiegel PB, Salama P. War and mortality in Kosovo, 1998-99: an epidemiological testimony. *Lancet* 2000; 355:2204–9. DOI:10.1016/S0140-6736(00)02404-1.
25. Garin N, Koyanagi A, Chatterji S, Tyrovolas S, Olaya B, Leonardi M, et al. Global Multimorbidity Patterns: a cross-sectional, population-based, multi-country study. *J Gerontol A Biol Sci Med Sci*. 2016;71(2):205-14. DOI: 10.1093/gerona/qlv128.
26. EUROSTAT. *Statistic Explained. Self-perceived health statistics*. 2018. https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Self-perceived_health_statistics#Self-perceived_health [last assessed September 22, 2020]
27. Gallagher JE, Wilkie AA, Corder A, Hudgens EE, Ghio AJ, Birch RJ, et al. Factors associated with self-reported health: implications for screening level community-based health and environmental studies. *BMC Public Health*. 2016;16:640. DOI: 10.1186/s12889-016-3321-5.
28. Hendler RA, Ramchandani VA, Gilman J, Hommer DW. Stimulant and sedative effects of alcohol. *Curr Top Behav Neurosci*. 2013;13:489-509. DOI: 10.1007/7854_2011_135.
29. Kaplan MS1, Huguet N, Feeny D, McFarland BH, Caetano R, Bernier J, Giesbrecht N, Oliver L, Ross N. Alcohol use patterns and trajectories of health-related quality of life in middle-aged and older adults: a 14-year population-based study. *J Stud Alcohol Drugs*. 2012;73(4):581-90. DOI:10.15288/jsad.2012.73.581
30. Woldeyohanes TR, Woldehaimanot TE, Kerie MW, Mengistie MA, Yesuf EA. Perceived patient satisfaction with in-patient services at Jimma University Specialized Hospital, Southwest Ethiopia. *BMC Res Notes*. 2015;8:285. DOI: 10.1186/s13104-015-1179-8
31. Tehrani AB, Feldman SR, Camacho FT, Balkrishnan R. Patient Satisfaction with Outpatient Medical Care in the United States. *Health Outcomes Res Med*. 2011;2(4):e197–e202. DOI:10.14744/nci.2014.30074.
32. Stepurko T, Paylova M, Groot W. Overall satisfaction of health care users with the quality of and access to health care services: a cross-sectional study in six Central and Eastern European countries. *BMC Health Serv Res*. 2016;16(a):342. DOI: 10.1186/s12913-016-1585-1.
33. Epping-Jordan JE, Pruitt SD, Bengoa R, Wagner EH. Improving the quality of health care for chronic conditions. *Qual Saf Health Care*. 2004;13(4):299-305. DOI: 10.1136/qhc.13.4.299
34. Fufa BD, Negao EB. Satisfaction of Outpatient Service Consumers and Associated Factors Towards the Health Service Given at Jimma Medical Center, South West Ethiopia. *Patient Relat Outcome Meas*. 2019 ;10:347-54. DOI: 10.2147/PROM.S220404.
35. Doyle C, Lennox L, Bell D. A systematic review of evidence on the links between patient experience and clinical safety and effectiveness. *BMJ Open*. 2013;3(1). DOI:10.1136/bmjopen-2012-001

Table 1. Socio-demographics and lifestyle characteristics

Variables	Number of non-communicable diseases				p-value
	None (532)	One (288)	Multimorbid (247)	Total (1067)	
	n (%)	n (%)	n (%)	n (%)	
Gender					
Male	266 (51.2)	153 (29.4)	101 (19.4)	520 (48.7)	0.013
Female	266 (48.6)	135 (24.7)	146 (26.7)	547 (51.3)	
Age	34.8 (12.3)	46.8 (15.8)	52.7 (15.4)	42.6 (16)	0.001
Place of residence					
North of Kosovo	235 (48.5)	149 (30.7)	101 (20.8)	485 (45.5)	0.031
South part of Kosovo	296 (51)	139 (23.9)	146 (25.1)	581 (54.5)	
Relationship status					
Single	209 (19.6)	70 (6.6)	59 (5.5)	338 (31.7)	0.001
In relationship	323 (30.3)	218 (20.4)	188 (17.6)	729 (68.3)	
Education					
≤ 8 years	7 (9.1)	20 (26.0)	50 (64.9)	77 (7.2)	0.001
8-12 years	310 (51)	178 (29.3)	120 (19.7)	608 (57)	
>12 years	215 (56.3)	90 (23.6)	77 (20.1)	382 (35.8)	
Employment					
Unemployed	226 (44.8)	138 (27.4)	140 (27.8)	504 (47.2)	0.001
Employed	306 (54.4)	150 (26.6)	107 (19)	563 (52.8)	
Having breakfast					
Irregular	93 (48.9)	50 (26.3)	47 (24.8)	190 (17.8)	0.848
Regular	439 (50.1)	238 (27.1)	200 (22.8)	877 (82.2)	
Fruits intake					
Rare or never	86 (45.5)	53 (28)	50 (26.5)	189 (17.7)	0.562
Weekly	274 (51)	151 (28.1)	112 (20.9)	537 (50.3)	
Daily	172 (50.4)	84 (24.6)	85 (25)	341 (32)	
Vegetable intake					
Rare or never	62 (44.6)	36 (25.9)	41 (29.5)	139 (13)	0.757
Weekly	293 (50.9)	161 (28)	122 (21.1)	576 (54)	
Daily	177 (50.3)	91 (25.9)	84 (23.9)	352 (33)	
Physical activity					
No	247 (41.3)	166 (27.8)	185 (30.9)	598 (56)	0.001
Yes	285 (60.8)	122 (26.0)	62 (13.2)	469 (44)	
Drug use					
No	527 (49.9)	286 (27.1)	243 (23)	1056 (99)	0.549
Yes	5 (45.5)	2 (18.2)	4 (36.3)	11 (1)	
Smoking					
No	301 (53.7)	143 (25.5)	117 (20.8)	561 (52.6)	0.029
Yes	231 (45.6)	145 (28.7)	130 (25.7)	506 (47.4)	
Alcohol use					
No	243 (42.6)	158 (27.7)	170 (29.7)	571 (53.5)	0.001
Yes	289 (58.3)	130 (26.2)	77 (15.5)	496 (46.5)	
Alcohol use (binge drinking)					
Never or rare	318 (45.6)	188 (27)	191 (27.4)	697 (65.3)	0.001
Monthly	162 (57.6)	80 (28.5)	39 (13.9)	281 (26.3)	
Weekly	49 (62)	17 (21.5)	13 (16.5)	79 (7.4)	
Daily	3 (30)	3 (30)	4 (40)	10 (1)	

Statistics: KW χ^2 , mean and SD; Bold values are statistically significant

Table 2. Self-rated health status and satisfaction with healthcare compared to the number of non-communicable diseases

Variables	Number of non-communicable diseases				p-value
	None (532)	One (288)	Multimorbid (247)	Total (1067)	
	n (%)	n (%)	n (%)	n (%)	
Self-rated health status					
Poor, very poor	3 (5.5)	10 (18.2)	42 (76.3)	55 (5.2)	0.001
Moderate	36 (14.3)	93 (36.9)	123 (48.8)	252 (23.6)	
Good, very good	493 (64.9)	185 (24.3)	82 (10.8)	760 (71.2)	
Satisfaction with healthcare					
Unsatisfied, very unsatisfied	55 (46.6)	31 (26.3)	32 (27.1)	118 (11.1)	0.029
Neutral	152 (45.8)	90 (27.1)	90 (27.1)	332 (31.1)	
Satisfied, very satisfied	325 (52.7)	167 (27.1)	125 (20.2)	617 (57.8)	
Long waiting time					
I did not need it	336 (58.7)	144 (25.2)	92 (16.1)	572 (53.6)	0.001
No	172 (42)	116 (28.4)	121 (29.6)	409 (38.3)	
Yes	24 (27.9)	28 (32.5)	34 (39.6)	86 (8.1)	
Availability due to transportation					
I did not need it	343 (57.3)	156 (26.1)	99 (16.6)	598 (56)	0.001
No	178 (42)	123 (29.1)	135 (31.9)	436 (40.9)	
Yes	11 (33.3)	9 (27.3)	13 (39.4)	33 (3.1)	
Lack of finance for healthcare					
I did not need it	338 (58.6)	139 (24.1)	100 (17.3)	577 (54.1)	0.001
No	179 (42.3)	129 (30.5)	115 (27.2)	423 (39.6)	
Yes	15 (22.4)	20 (29.9)	32 (47.7)	67 (6.3)	
Lack of finance for dental care					
I did not need it	324 (57.3)	139 (24.6)	102 (18.1)	565 (53)	0.001
No	188 (42.3)	134 (30.2)	122 (27.5)	444 (41.6)	
Yes	20 (34.5)	15 (25.9)	23 (39.6)	58 (5.4)	
Lack of finance for medication prescription					
I did not need it	334 (58.1)	141 (24.5)	100 (17.4)	575 (53.9)	0.001
No	190 (41.9)	134 (29.6)	129 (28.5)	453 (42.5)	
Yes	8 (20.5)	13 (33.3)	18 (46.2)	39 (3.6)	
Lack of finance for mental care					
I did not need it	50 (56.4)	150 (24.2)	121 (19.4)	621 (58.2)	0.001
No	182 (41.7)	136 (31.1)	119 (27.2)	437 (41.0)	
Yes	0 (0)	2 (22.2)	7 (77.8)	9 (0.8)	

Statistics: KW χ^2 , mean and SD; Bold values are statistically significant

Table 3. Multinomial regression model describing factors associated with self-reported health status

Variables	Self-rated health status (Moderate vs. Poor, very poor)		Self-rated health status (Good, very good vs. Poor, very poor)	
	<i>p-value</i>	OR (95% CI of OR)	<i>p-value</i>	OR (95% CI of OR)
Number of chronic diseases				
None	0.195	2.35 (0.64–8.56)	0.001	23.33 (6.53–83.35)
One	0.016	2.68 (1.20–5.98)	0.001	7.38 (3.19–17.06)
Multimorbidity	I		I	
Gender				
Male	0.184	0.60 (0.29–1.26)	0.991	0.99 (0.46–2.13)
Female	I		I	
Age	0.272	0.99 (0.96–1.01)	0.001	0.93 (0.90–0.95)
Place of residence				
North of Kosovo	0.851	1.07 (0.54–2.11)	0.242	0.65 (0.32–1.33)
South part of Kosovo	I		I	
Relationship status				
Single	0.293	0.67 (0.32–1.41)	0.325	0.67 (0.30–1.49)
In relationship	I		I	
Education				
≤ 8 years	0.339	0.59 (0.20–1.73)	0.082	0.34 (0.10–1.15)
8–12 years	0.892	1.06 (0.44–2.58)	0.462	0.72 (0.29–1.75)
> 12 years	I		I	
Employment				
Unemployed	0.128	0.53 (0.24–1.20)	0.032	0.40 (0.18–0.93)
Employed	I		I	
Having breakfast				
Irregular	0.101	2.43 (0.84–7.02)	0.341	1.71 (0.57–5.13)
Regular	I		I	
Fruits intake				
Rare or never	0.631	1.31 (0.44–3.89)	0.324	0.56 (0.18–1.78)
Weekly	0.733	1.16 (0.50–2.69)	0.908	0.39 (0.39–2.28)
Daily	I		I	
Vegetable intake				
Rare or never	0.607	0.73 (0.23–2.39)	0.729	0.80 (0.23–2.79)
Weekly	0.945	1.03 (0.44–2.40)	0.901	0.95 (0.39–2.27)
Daily	I		I	
Physical activity				
No	0.495	1.31 (0.60–2.86)	0.909	0.96 (0.43–2.11)
Yes	I		I	
Drug use				
No	0.474	2.55 (0.19–32.93)	0.301	4.27 (0.27–66.83)
Yes	I		I	
Smoking				
No	0.903	1.04 (0.52–2.09)	0.362	1.40 (0.68–2.89)
Yes	I		I	
Alcohol use				
No	0.004	0.23 (0.09–0.62)	0.002	0.20 (0.06–0.55)
Yes	I		I	

Bold values are statistically significant; **I** - reference category

Table 4. Multinomial regression model describing factors associated with healthcare satisfaction among respondents

Variables	Satisfaction with healthcare (Neutral vs. Unsatisfied, very unsatisfied)		Satisfaction with healthcare (Satisfied, very satisfied vs. Unsatisfied, very unsatisfied)	
	<i>p-value</i>	OR (95% CI of OR)	<i>p-value</i>	OR (95% CI of OR)
Model 1. Socio-demographic and lifestyle				
Number of chronic diseases	0.974		0.178	
None	0.778	0.99 (0.55–1.79)	0.209	1.48 (0.84–2.60)
One	1	1.09 (0.59–2)	1	1.45 (0.81–2.58)
Multimorbidity				
Gender	0.315	0.78 (0.48–1.27)	0.592	1.13 (0.72–1.79)
Male	1		1	
Female				
Age	0.806	0.99 (0.98–1.02)	0.580	1.01 (0.99–1.02)
Place of residence	0.498	1.17 (0.75–1.83)	0.206	0.76 (0.49–1.16)
North of Kosovo	1		1	
South part of Kosovo				
Relationship status	0.017	2.02 (1.14–3.59)	0.011	2.06 (1.18–3.57)
Single	1		1	
In relationship				
Education	0.164	1.97 (0.76–5.11)	0.496	0.72 (0.28–1.85)
≤ 8 years	0.031	1.69 (1.05–2.73)	0.047	1.58 (1.01–2.47)
8–12 years	1		1	
>12 years				
Employment	0.563	0.87 (0.55–1.39)	0.406	0.83 (0.53–1.29)
Unemployed	1		1	
Employed				
Having breakfast	0.384	0.79 (0.47–1.34)	0.032	0.57 (0.34–0.95)
Irregular	1		1	
Regular				
Fruits intake	0.423	1.35 (0.65–2.84)	0.212	0.64 (0.32–1.29)
Rare or never	0.225	1.426 (0.80–2.53)	0.225	0.72 (0.42–1.23)
Weekly	1		1	
Daily				
Vegetable intake	0.009	0.32 (0.14–0.75)	0.633	0.82 (0.37–1.83)
Rare or never	0.013	0.48 (0.27–0.85)	0.029	0.54 (0.31–0.94)
Weekly	1		1	
Daily				
Physical activity	0.549	1.16 (0.72–1.86)	0.543	0.87 (0.56–1.36)
No	1		1	
Yes				
Drug use	0.374	2.53 (0.33–19.50)	0.931	1.08 (0.19–5.87)
No	1		1	
Yes				
Smoking	0.472	0.85 (0.54–1.34)	0.520	1.15 (0.75–1.77)
No	1		1	
Yes				
Alcohol use	0.352	1.26 (0.77–2.07)	0.010	1.85 (1.16–2.95)
No	1		1	
Yes				
Model 2. Unavailability of healthcare				

Number of chronic diseases				
None	0.368	0.76 (0.45–1.35)	0.720	1.10 (0.65–1.88)
One	0.909	0.97 (0.53–1.78)	0.542	1.20 (0.67–2.16)
Multimorbidity	I		I	
Long waiting time				
I did not need it	0.001	7.27 (2.15–24.59)	0.004	4.95 (1.65–14.88)
No	0.001	5.39 (2.59–11.19)	0.001	7.37 (3.68–14.74)
Yes	I		I	
Availability due to transportation				
I did not need it	0.368	0.52 (0.13–2.16)	0.185	2.69 (0.62–11.64)
No	0.660	0.79 (0.27–2.29)	0.320	1.82 (0.56–5.94)
Yes	I		I	
Lack of finance for healthcare				
I did not need it	0.555	0.68 (0.19–2.45)	0.712	0.79 (0.22–2.79)
No	0.775	0.88 (0.35–2.17)	0.940	1.04 (0.42–2.56)
Yes	I		I	
Lack of finance for dental care				
I did not need it	0.960	1.03 (0.31–3.47)	0.021	4.37 (1.25–15.27)
No	0.607	0.79 (0.31–1.97)	0.168	1.99 (0.75–5.27)
Yes	I		I	
Lack of finance for medication prescription				
I did not need it	0.184	0.35 (0.08–1.64)	0.150	0.33 (0.07–1.50)
No	0.131	0.34 (0.09–1.38)	0.297	0.47 (0.12–1.94)
Yes	I		I	
Lack of finance for mental care				
I did not need it	0.315	3.14 (0.34–29.36)	0.784	1.37 (0.15–12.87)
No	0.548	1.94 (0.22–16.88)	0.912	1.13 (0.13–9.99)
Yes	I		I	

Bold values are statistically significant; **I** - reference category