



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Case Report / Приказ болесника

Mladen Bila^{1,*}, Tanja Kalezić^{1,2}, Igor Kovačević^{1,2}, Goran Damjanović¹, Dijana Risimić^{1,2}

**Uncommon severe blunt ocular injury associated with large retinal dialysis
caused by a fishing sinker**

Неуобичајено тешка контузиона повреда ока удружена са великом
дијализом ретине настала металним тегом за пецање

¹Clinical Center of Serbia, Clinic for Eye Diseases, Belgrade, Serbia;

²University of Belgrade Faculty of Medicine, Belgrade, Serbia

Received: June 19, 2020

Revised: October 9, 2020

Accepted: October 11, 2020

Online First: October 14, 2020

DOI: <https://doi.org/10.2298/SARH200619094B>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

***Correspondence to:**

Mladen BILA

Clinic for Eye Diseases, Clinical Center of Serbia, Pasterova 2, 11000 Belgrade, Serbia

E-mail: mladen.bila@gmail.com

Uncommon severe blunt ocular injury associated with large retinal dialysis caused by a fishing sinker

Неуобичајено тешка контузиона повреда ока удружена са великом дијализом ретине настала металним тегом за пецање

SUMMARY

Introduction Severe ocular injuries caused by fishing equipment are relatively rare. The visual prognosis for fishing-related injuries depends on the involved ocular structures, the presence of complications and surgical techniques applied.

Case outline A 40-year-old man reported the sudden severe sharp pain and loss of vision in his left eye while he was pulling the fishing rod during recreational fishing. At admission, his best corrected visual acuity (BCVA) was 1/60 at the left eye. Clinical examination revealed a laceration in the temporal quadrant of bulbar conjunctiva of the 0.2-0.3mm size and a dark tumefaction under the conjunctiva in the same region. It was identified during primary surgical exploration as a fishing sinker of the 1cm size lodged under the lateral rectus muscle. The bulbar wall was intact and the fishing sinker was safely removed. Phacoemulsification with implantation of artificial foldable intraocular lens and 23G pars plana vitrectomy were performed. During vitrectomy, subretinal hemorrhage in the macular region and large retinal dialysis in the temporal segment were revealed. Vitrectomy was finished with silicone oil tamponade. One week later, the patient's BCVA was 2/60 at the left eye. At 8 months after surgery spontaneous resorption of subretinal hemorrhage in the macular region and the attached retina were observed. The patient's BCVA was 3/60 due to destruction of photoreceptors and retinal pigment epithelium and formation of epiretinal membrane.

Conclusion To our knowledge this is the first case report of severe blunt ocular injury associated with large retinal dialysis caused by a fishing sinker.

Keywords: ocular trauma; fishing sinker; subretinal hemorrhage; retinal dialysis; vitrectomy

САЖЕТАК

Увод Тешке повреде ока настале опремом за пецање су релативно ретке. Функционална прогноза по вид, код повреда насталих опремом за пецање зависи од захваћених окуларних структура, присуства компликација и хируршких техника које се примењују код различитих типова повреда.

Приказ болесника Мушкарац, узраста 40 година, током рекреативног пецања је, повлачећи штап, одједном осетио веома оштар бол и губитак вида на левом оку. Најбоље коригована видна оштрина на левом, повређеном оку, на пријему је износила 1/60. Клиничким прегледом, уочена је лацерација булбарне конјунктиве величине 0,2–0,3 мм као и тамна тумефакција испод конјунктиве у истом, темпоралном сегменту. Тамна тумефакција испод конјунктиве је током примарне хируршке експлорације идентификована као метални тег за пецање величине 1 цм, који се налазио испод спољашњег правог мишића. Булбарни зид је био интактан а метални тег пажљиво уклоњен. Затим је урађена факоемулзификација природног сочива са уградњом савитљивог, вештачког, интраокуларног сочива као и 23Г *pars plana* витректомија. Током витректомије откривено је присуство субретиналне хеморагије у макули као и велика дијализа ретине у темпоралном квадранту. Витректомија је завршена тампонадом силиконским уљем. На првој постоперативној контроли, недељу дана касније, најбоље коригована видна оштрина на левом оку је износила 2/60. Након осам месеци од операције контролни налаз је показао спонтану ресорцију субретиналне хеморагије у макули са налегнутим ретином. Најбоље коригована видна оштрина је износила 3/60 као последица деструкције фоторецептора и ретиналног пигментног епитела у макули и формирања епиретиналне мембране.

Закључак Према нашим сазнањима, ово је први приказ тешке контузионе повреде ока удружене са великом дијализом ретине настале металним тегом за пецање.

Кључне речи: траума ока; метални тег за пецање; субретинална хеморагија; дијализа ретине; витректомија

INTRODUCTION

Ocular injury is a main cause of visual morbidity and blindness in the adult age worldwide [1]. Numerous causes of eye trauma exist and sport-related injuries have been described in the literature. Fishing is a popular activity and usually no particular safety measures are undertaken. However, ocular traumas ranging from simple to severe can occur during recreational fishing. Severe eye injuries caused by fishhooks and other parts of fishing equipment are relatively rare. Cases of ocular injuries caused by fishhook are mostly reported and the severity of ocular injuries depends on the involved ocular structures. Various structures of the eye including the lid, conjunctiva, cornea, sclera, anterior chamber, lens and the posterior segment structures may be affected in fishhook ocular injury. Possible complications of these injuries may involve anterior and posterior segment with partial or complete loss of vision and even loss of the eye in certain circumstances. Treatment of these injuries depends on the location of the injury, the involved ocular structures, and the type of fishhook or other parts of fishing equipment [2]. In a suspicion of metal foreign body eye injury detailed examination of anterior and posterior segment structures is required including X-ray or CT of orbits and ultrasound examination [3].

We present uncommon, very severe blunt eye injury associated with large retinal dialysis caused by a fishing sinker. We describe the unusual mechanism, severity, treatment and clinical outcome of injury in a patient which occurred during recreational fishing.

CASE REPORT

A 40-year-old man was admitted at the Clinic for eye diseases after suffering trauma to his left eye during recreational fishing. On initial presentation the patient reported the sudden severe sharp pain in his left eye and loss of vision with a sensation of dark curtain coming down across his left eye that occurred while he was pulling the fishing rod.

The patient's best corrected visual acuity was 1.0 on the right eye and 1/60 on the left eye. Intraocular pressure on both eyes was 16mmHg. Slit lamp examination of the anterior and posterior segment of the right eye was normal. Slit lamp examination of the left eye anterior segment showed suffusion and hyperemia of the bulbar conjunctiva with the presence of small laceration in temporal quadrant of the 0.2-0.3mm size. The dark

tumefaction under the bulbar conjunctiva in the region of small laceration in the zone of lateral rectus muscle was observed. Cornea was intact, anterior chamber was deeper and clear with traumatic mydriasis and slightly irregular pupil with clear lens in physiological position. Eye fundus examination demonstrated haemophthalmus. Ultrasound examination of the left eye revealed the presence of mobile vitreous opacities of low to medium reflection, the findings consistent to haemophthalmus, no echo signs of intraocular foreign body and the zone of high reflection of the detached retina (Figure 1).

The presence of dark tumefaction in the zone of lateral rectus muscle under the bulbar conjunctiva raised the suspicion of globe rupture with the prolapsed uveal tissue. Primary surgical exploration was done in the local anesthesia. Intact bulbar wall and unexpected presence of a metal foreign body, a fishing sinker, under the lateral rectus muscle of the 1cm size were revealed. The fishing sinker was extracted (Figures 2A and 2B). Phacoemulsification of natural lens with implantation of artificial foldable intraocular lens (Figure 3) and 23G pars plana vitrectomy with silicone oil tamponade were performed. Following three ports sclerotomies at pars plana, at 3.5mm from limbus and the placement of the infusion line, „core“ vitrectomy and removal of haemophthalmus were performed enabling the visualization of fundus. Several pre-retinal haemorrhages, subretinal haemorrhage in the macular region and large retinal dialysis in the temporal segment were observed (Figures 4A and 4B). In this segment retina appeared wrinkled, immobile with irregular posterior border with attached vitreous and small quantity of subretinal fluid. Triamcinolon-acetonid (Kenalog) was applied in vitreous for better visualization, especially in the zone of posterior border of dialysis and at the vitreous base. Posterior border of the dialysis was relaxed from vitreal traction and the retina became mobile. Vitreous base was cleared along the complete circumferention. Perfluorocarbon (Decalin) was installed and the retina was stabilized and attached in the zone of damage. Laser photocoagulation was applied in several rows along complete circumferention of peripheral retina and in the zone of posterior border of dialysis. Perfluorocarbon-air exchange was done and surgical treatment was finished with air-silicone oil exchange (Figures 5A and 5B). During hospitalization, patient received antibiotics, corticosteroids and midriatic drug locally as well as systemic antibiotics and corticosteroids. In the early postoperative period the mild increase of intraocular pressure of 26mmHg was noticed and treated locally with antiglaucomatous therapy. He was discharged from the hospital and at the control visit one week later the patient's best corrected visual acuity was 2/60 at the left eye with intraocular pressure of

15mmHg. At 8 months after surgery the patient's best corrected visual acuity was 3/60. At the same control visit, photo fundus showed the spontaneous resorption of subretinal haemorrhage in the macular region, attached retina under the silicone oil with scar tissue at the posterior border of dialysis (Figures 6A and 6B). Macular optical coherence tomography findings showed thinning, photoreceptors and retinal pigment epithelium destruction and epiretinal membrane (Figure 7).

All procedures performed were in accordance with the ethical standards of the institutional and national research committee and with the 1964 Helsinki declaration and its later amendment. Written consent to publish all shown material was obtained from the patient.

DISCUSSION

The majority of reported cases of fishing-related ocular injuries were caused by fishhook. These reports describe the types and severity of injuries, and the surgical techniques applied for fishhook removal [4, 5]. To our knowledge this is the first case report of a patient with severe blunt ocular injury caused by a fishing sinker during recreational fishing. The eye injury was caused by a metal fishing sinker of 1cm size which penetrated bulbar conjunctiva leaving the laceration of 0.2-0.3mm size. The fishing sinker was revealed during the primary surgical exploration of eye globe and it was embedded under the lateral rectus muscle and removed. During vitrectomy large retinal dialysis was observed in the temporal quadrant in the projection of contusion scleral wall injury caused by the fishing sinker. Large retinal dialysis, subretinal haemorrhage in macular region and haemophthalmus determined the severity of eye injury in this patient.

In retrospective analysis using United States Eye Injury Registry, Alfaro et al. [6] reported that 19.54% of sport-related ocular traumas occurred during fishing. Corneal laceration, globe rupture and hyphaema are the most frequent eye injuries caused by fishhook, fishing lure or weights. In a study by Hoskin et al. [7] fishing-related eye injuries were found in 7% of sport-related injuries in a Western Australian pediatric population. Purtskhvanidze et al. [2] reported nine patients who experienced eye injury caused by fishhook. Five patients had only eyelid injury (55%) and four patients suffered open globe injuries (45%). Among those, one patient had eye trauma limited to cornea (25%), one patient

had scleral injury (25%) and two patients had injuries of lens, iris and posterior segment (50%). Surgical treatment described in this report included primary pars plana vitrectomy performed for endophthalmitis and for retinal detachment with the giant retinal tear and choroid haemorrhage. Choovuthayakorn et al. [4] described primary wound management, the “back-out” technique of removing fishhook from upper eyelid which was embedded during the fishing. Fishhook penetrated cornea, iris and was lodged in the ciliary body behind the clouded natural lens. The patient underwent lensectomy with pars plana vitrectomy and silicone oil tamponade for the treatment of the local retinal detachment. Uncommon mechanism of eye injury and the surgical treatment is reported by Iannetti and Tortorella [5] in patients who suffered fishhook injury that penetrated through sclera at 2mm from corneal limbus and trabeculum and was lodged in the anterior chamber. Nakatsuka et al. [8] described “cut-out” technique for removal of large fishhook that caused full thickness corneal penetration at the nasal limbus and was lodged in the anterior chamber angle of the eye.

We describe severe blunt eye trauma associated with haemophthalmus, subretinal haemorrhage in the macular region and large retinal dialysis in the temporal quadrant. Retinal tears in the region of the ora serrata and peripheral retina are typically caused by blunt trauma [9]. A retinal dialysis is a tear in the retina whose anterior edge is at ora serrata and whose posterior edge is attached to the vitreous base [10]. Retinal dialysis has been described as the most common complication after ocular contusion injury [11]. The incidence of rhegmatogenous retinal detachment caused by dialysis is 8-17% and the retinal dialyses are most often seen in younger men following trauma [12]. In contrast to eye trauma related giant retinal tears which are caused by vitreal traction, in retinal dialysis vitreous base is firmly attached to the posterior border, but the avulsion of vitreous base may occur and represents the pathognomonic sign of blunt eye injury [11]. Blunt injury causes retinal dialysis by the compression of the eye in anteroposterior plane which results in the expanding in equatorial plane and causes pressure in the vitreous base [11]. Retinal dialyses are most often located in inferotemporal quadrant. However, multiple dialyses, small dialyses and dialyses in several quadrants may exist [13].

In conclusion, to our knowledge this is the first case report of severe blunt ocular injury associated with large retinal dialysis caused by a fishing sinker. Accidents due to fishing equipment are rare and may result in serious ocular injuries and significant visual loss. The outcome of fishing-related ocular injury depends on affected eye structures, mechanism of

injury, the presence of complications and surgical techniques applied. We report treatments that were applied in a patient with severe ocular trauma caused by a fishing sinker that involved the posterior segment including subretinal macular haemorrhage and retinal dialysis, which enabled visual and anatomical satisfactory outcomes. However, to reduce the occurrence of fishing-related eye injuries, preventive measures should be undertaken such as the use of protective eyewear during this activity.

Conflict of interest: None declared.

Paper accepted

REFERENCES

1. Négrel AD, Thylefors B. The global impact of eye injuries. *Ophthalmic Epidemiol.* 1998;5(3):143-169. doi:10.1076/oep.5.3.143.8364
2. Purtskhvanidze K, Saeger M, Treumer F, Nölle B, Roider J. Open globe and penetrating eyelid injuries from fish hooks. *BMC Ophthalmol.* 2019;19(1):26. Published 2019 Jan 21. doi:10.1186/s12886-019-1040-2
3. Gaković A, Kovačević I, Biševac J, Radović B, Čubrilo K, Stefanović I. [Big intraocular foreign body: case report]. *Srp Arh Celok Lek.* 2013 Jul-Aug;141(7-8):516-8. Serbian. PMID: 24073560. doi: 10.2298/SARH1308516G
4. Choovuthayakorn J, Chavengsaksongkram P, Watanachai N, Chaidaroon W. Penetrating Eyelid and Ocular Fishhook-Related Injury. *Case Rep Ophthalmol.* 2019;10(1):41-46. Published 2019 Jan 24. doi:10.1159/000496382
5. Iannetti L, Tortorella P. Penetrating fish-hook ocular injury: management of an unusual intraocular foreign body. *Case Rep Med.* 2014;2014:901285. doi:10.1155/2014/901285
6. Alfaro DV 3rd, Jablon EP, Rodriguez Fontal M, Villalba SJ, Morris RE, Grossman M, Roig-Melo E. Fishing-related ocular trauma. *Am J Ophthalmol.* 2005;139(3):488-492. doi:10.1016/j.ajo.2004.10.011
7. Hoskin AK, Yardley AM, Hanman K, Lam G, Mackey DA. Sports-related eye and adnexal injuries in the Western Australian paediatric population. *Acta Ophthalmol.* 2016;94(6):e407-e410. doi:10.1111/aos.12911
8. Nakatsuka AS, Khanamiri HN, Merkley KH. Fishhook Injury of the Anterior Chamber Angle of the Eye. *Hawaii J Med Public Health.* 2019;78(6):200-201.
9. Ayalon A, Okrent L, Rubowitz A. Posterior pole retinal tears following blunt ocular trauma. *Am J Ophthalmol Case Rep.* 2020;18:100642. Published 2020 Feb 29. doi:10.1016/j.ajoc.2020.100642
10. Vote BJ, Casswell AG. Retinal dialysis: are we missing diagnostic opportunities?. *Eye (Lond).* 2004;18(7):709-713. doi:10.1038/sj.eye.6701310.
11. Stiff HA, Abbassi S, Cunningham CM, Peairs JJ, Folk JC, Sohn EH. Retinal Dialysis. *EyeRounds.org.* posted June 13, 2018; Available from <https://EyeRounds.org/cases/272-retinal-dialysis.htm>
12. Chang JS, Marra K, Flynn HW Jr, Berrocal AM, Arroyo JG. Scleral Buckling in the Treatment of Retinal Detachment Due to Retinal Dialysis. *Ophthalmic Surg Lasers Imaging Retina.* 2016;47(4):336-340. doi:10.3928/23258160-20160324-06
13. Qiang Kwong T, Shunmugam M, Williamson TH. Characteristics of rhegmatogenous retinal detachments secondary to retinal dialyses. *Can J Ophthalmol.* 2014;49(2):196-199. doi:10.1016/j.jcjo.2013.12.013

P. ar




Figure 1. Ultrasound of the left eye shows the presence of mobile vitreous opacities of low to medium reflection (haemophthalmus), no echo signs of intraocular foreign body and the zone of high reflection of the detached retina




Figure 2. Intact bulbar wall and extracted fishing sinker during primary surgical exploration (A); dimensions of the fishing sinker (B)

Paper accepted




Figure 3. Implanted, artificial foldable intraocular lens and 3-port PPV 23G system with infusion line

Paper accepted




Figure 4. Several pre-retinal hemorrhages and subretinal hemorrhage in the macular region (A); large retinal dialysis in temporal quadrant (B)

Paper accepted




Figure 5. Perfluorocarbon–air exchange and laser spots in several rows along posterior border of dialysis (A); surgical treatment was finished with air–silicone oil exchange (B)

Paper accepted




Figure 6. Photo fundus shows the spontaneous resorption of subretinal hemorrhage in the macular region (A) with attached retina under the silicone oil and the scar tissue at the posterior border of dialysis (B)




Figure 7. Optical coherence tomography of the macula shows thinning, photoreceptors and retinal pigment epithelium destruction and epiretinal membrane

Paper accepted