



СРПСКИ АРХИВ
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО
SERBIAN ARCHIVES
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: office@srpskiarhiv.rs, Web address: www.srpskiarhiv.rs

Paper Accepted*

ISSN Online 2406-0895

Original Article / Оригинални рад

Nevena Kalezić^{1,2}, Milica Karadžić-Kočica², Nemanja Dimić³, Mladen Kočica^{2,*},
Anka Tošković², Milan Jovanović², Ivan Dimitrijević^{1,2}

Alcohol abuse as a risk factor for developing thyroid cancer

Злоупотреба алкохола као фактор ризика за развој рака штитне жлезде

¹University of Belgrade, Faculty of Medicine, Belgrade, Serbia;

²Clinical Center of Serbia, Belgrade, Serbia;

³Dr. Dragiša Mišović Clinical Hospital Center, Belgrade, Serbia

Received: October 21, 2020

Accepted: November 23, 2020

Online First: November 26, 2020

DOI: <https://doi.org/10.2298/SARH201123113K>

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

***Correspondence to:**

Mladen J. KOČICA

Clinical Centre of Serbia, Clinic for Cardiac Surgery, 8 Kosta Todorović St., Belgrade 11000, Serbia

E-mail: kocica@sbb.rs

Alcohol abuse as a risk factor for developing thyroid cancer

Злоупотреба алкохола као фактор ризика за развој рака штитне жлезде

SUMMARY

Introduction/Objective Alcohol abuse influence on developing thyroid cancer is controversial. While some studies consider it a protective factor, others deny any impact on thyroid cancer.

The objective of the paper was to establish a possible link between alcohol abuse and certain types of thyroid cancers.

Methods The retrospective study included 502 patients with thyroid cancer and control group of 600 patients with benign forms of thyroid diseases (e.g. nodular, multinodular and toxic nodular goiter). Thyroid cancer patients were divided into 4 groups: I – papillary, II – medullary, III – anaplastic, and IV – follicular carcinoma, and grouped by sex, age (< 30 yrs.; > 30 yrs.) and alcohol abuse, as defined by WHO.

Results Thyroid cancer patients were predominantly male of younger age. This distribution difference was statistically significant in groups I and II ($p < 0.001$). Of total 10 (0.9%) patients with chronic alcohol abuse, 8 (1.6%) had thyroid cancer, while 2 (0.3%) belonged to the control group ($p < 0.001$). In thyroid cancer patients, chronic alcohol abuse was absent in group III and IV. Distribution in group I and II was 6 (1.6%) and 2 (2%) respectively ($p < 0.001$).

Conclusion Alcohol abuse deserves to be considered as a risk factor for papillary and medullary forms of thyroid cancer, while it does not stay the same for anaplastic and follicular thyroid cancers.

Keywords: thyroid cancer; papillary cancer; medullary cancer; anaplastic cancer; follicular cancer; alcohol abuse

САЖЕТАК

Увод/Циљ Утицај злоупотребе алкохола на развој рака штитне жлезде је контроверзан. Док неке студије то сматрају заштитним фактором, друге негирају било какав утицај на рак штитне жлезде.

Циљ рада је анализа могуће везе између хроничне злоупотребе алкохола и одређених врста рака штитне жлезде.

Метод Ретроспективна студија обухватила је 502 пацијента оболелих од рака штитне жлезде и контролну групу од 600 пацијената са доброћудним облицима болести ове жлезде (нпр. нодуларна, мултинодуларна и токсична нодуларна струма). Оболели од рака штитне жлезде подељени су у четири групе: I – папиларни, II – медуларни, III – анапластични и IV – фоликуларни карцином и груписани по полу, узрасту (< 30 год.; > 30 год.) и злоупотреби алкохола, у складу са дефиницијом СЗО.

Резултати Оболели од рака штитне жлезде били су претежно мушкарци млађег узраста. Ова разлика у расподели је статистички значајна и групама I и II ($p < 0.001$). Од укупно 10 (0.9%) пацијената са хроничном злоупотребом алкохола, 8 (1.6%) је имало рак штитне жлезде, док је 2 (0.3%) припадало контролној групи ($p < 0.001$). Код оболелих од рака штитне жлезде, хронична злоупотреба алкохола није забележена у групама III и IV. Дистрибуција у групама I и II је била 6 (1.6%), односно 2 (2%) ($p < 0.001$).

Закључак Злоупотреба алкохола заслужује да буде размотрена као фактор ризика за папиларне и медуларне облике рака штитне жлезде, док то исто не важи за анапластични и фоликуларни рак штитне жлезде.

Кључне речи: рак штитњаче; папиларни рак; медуларни рак; анапластични рак; фоликуларни рак; злоупотреба алкохола

INTRODUCTION

Apart from the social, mental and behavioral disturbances, chronic alcohol abuse (CAA) causes and/or affects many serious somatic diseases, including cancer [1, 2]. Among the surgical patients, different drinking patterns may also affect specific features in the management of anesthesia, patient behavior and different complications in the perioperative period [2, 3, 4].

Alcohol abuse was addressed as a possible cause or contributing factor for thyroid cancers and other non-cancerous thyroid diseases by many observational studies. The results of these studies are different, sometimes inconclusive or even conflicting [5, 6]. Yet, the abundant evidence of the increasing incidence of thyroid cancers, attributed mainly to increased detection of papillary thyroid cancer, deserves a careful analysis of all possible risk factors, including CAA [7, 8]

We designed a retrospective, cross-sectional, study to determine a possible influence of CAA on thyroid cancer incidence.

The objective of this study was to determine if CAA was a risk factor for thyroid cancer in general, as well as for different types of thyroid cancers (i.e. I-papillary, II-medullary, III-anaplastic and IV-follicular carcinoma).

METHODS

Total of 1102 consecutive patients who underwent thyroid surgery at the Center for Endocrine Surgery, Clinical Center of Serbia during the three consecutive years were analyzed. The study group included 502 patients with different forms of thyroid cancer and the control group included 600 patients with benign or degenerative diseases of the thyroid gland. Thyroid cancer patients were divided, according to histopathological findings, into four groups: I - papillary carcinoma (380 patients, 75.7%), II - medullary carcinoma (102 patients, 20.3%), III - anaplastic carcinoma (10 patients, 2%) and IV - follicular carcinoma (10 patients, 2%). The control group consisted of patients with thyroid nodule (233 patients, 38.8%), multinodular goiter (337 patients, 56.2%), and toxic adenoma (30 patients, 5%) (Table 1). All patients with autoimmune thyroid diseases were excluded from the study.

Patients' records were used to collect demographic (age, gender) and clinical data (present and past diseases and surgeries) as well as socio-epidemiological questionnaire (exposures, habits, abuses) with a particular accent on CAA (type, dose, pattern), as defined by the World Health Organization (WHO) [9]. For this study, CAA was defined at least as moderate alcohol intake and/or alcohol dependence. Accordingly, moderate alcohol intake

implies daily consumption of 1–2 (women) or 3–4 (men) standard drinks. Standard drink implies 0.03L of distilled beverage or 0.2L of wine or 0.3L of beer. Alcohol dependence is present if in the past year, at least 3/7 criteria were present: craving, the irresistible need for alcohol; increased tolerance; loss of control; abstinence syndrome; use of the same or of related substances to relieve the withdrawal syndrome; progressive neglect of alternative pleasures (socializing, hobbies, sports, etc.) and specific drinking pattern.

All data were collected into an electronic database (IBM SPSS® Statistics, V26) and presented in tables. Pearson's chi-squared test was used to compare the difference between categorical variables and the p-value was set < 0.05 for the statistically significant difference.

This study was approved by the Ethical Committee of the Faculty of Medicine, University of Belgrade (No.1575/7).

RESULTS

Papillary carcinoma was the most common form in thyroid cancer group (75.7%) while multinodular (56.2%) and nodular goiter (38.8%) made 95% of the control group pathologies (Table 1).

The mean age of patients was similar in a group with thyroid cancer (50.34 years) and control group (50.88 years). Patients under the age of 30 were significantly more represented in cancer than in the control group (13.3% vs. 6.8%, $p < 0.001$). The same is true for the male gender (19.7% vs. 11.7%, $p < 0.001$) distribution (Table 2).

The I-group (i.e. papillary carcinoma), compared with the control group, had significantly more patients under the age of 30 (14.5% vs. 6.8%, $p = 0.000$) and patients of male gender (18.7% vs. 11.7%, $p = 0.002$). The II-group (i.e. medullary carcinoma) had no age difference but did show a significant male predominance, compared to control (26.5% vs. 11.7%, $p = 0.000$). All patients from the III-group (i.e. anaplastic carcinoma) were over the age of 30, but this fact provided no statistically significant difference with the control group (100% vs. 93.2%, $p = 0.392$). The IV-group (i.e. follicular carcinoma) had no significant

difference in age ($p = 0.695$) and gender ($p = 0.251$) distribution, compared with the control group, despite old age and female predominance (Table 3).

There was an overall significant difference in CAA distribution between cancer and control group (1.6%, 8/502 patients vs. 0.3%, 2/600 patients, $p < 0.001$). The presence of CAA was recorded only in I- and II-group of thyroid cancer patients, with incidences significantly higher than the control group (1.6% and 2.0% vs. 0.3%, $p = 0.034$ and $p = 0.044$). There were no records of CAA in III- and IV-group. The incidence in these groups was significantly lower than the control group ($p = 0.001$) (Table 4).

DISCUSSION

Almost 10% of men in the Republic of Serbia (ROS) had alcohol use disorders, compared to 2.1% of women [10]. Alcohol consumption is an attributable risk for 5.1% of all-cause deaths in our country [11]. There is epidemiological evidence that alcohol causes cancer at seven sites in the body (oropharynx, larynx, esophagus, liver, colon, rectum and breast), although without exact and complete knowledge of underlying biological mechanisms. (1) Pandemic increase in thyroid cancer incidence over the two past decades resulted in significant efforts towards early detection and therapy, but also deeper analyses of possible toxic, environmental and socio-economic causes [12].

Many studies, so far, have addressed alcohol abuse as a possible risk factor for thyroid cancer [6, 13, 14]. A recent and, so far, the most comprehensive meta-analysis of 33 observational studies which involved a total of 7725 thyroid cancer patients and 3113679 participants without it, suggested that alcohol intake may decrease the risk of thyroid cancer. In a subgroup meta-analyses by geographic region, alcohol intake was associated with a decreased risk of thyroid cancer in the American, but not in the European or Asian regions [6]. Previous studies of risk factors for thyroid cancer, published in ROS, also haven't found any correlation with CAA [15, 16].

Yet, our study has shown that younger (under the age of 30 years), male patients, with history of CAA were at higher risk for overall and particularly for papillary (I-group) and

medullary (II-group) forms of thyroid carcinoma, compared to control group of non-cancerous thyroid patients.

Results of a study from South Korea, which has the highest incidence of thyroid cancers in the world, based on data collected from 12276 individuals, among others, reveals CAA (OR, 1.89; 95%CI 1.08-3.32) as a significant risk factor for thyroid cancer [14]. Data from the Thyroid Cancer Longitudinal Study (T-CALOS) on 2258 thyroid cancer patients and 22580 healthy individuals showed that acute high-dose and chronic lifetime exposure (>31 years) to alcohol are linked to an increased risk of development of thyroid cancer [13]. In addition to these findings, another study from the same country, comparing health behaviours of 942 thyroid cancer survivors with 9420 matched non-cancer controls, found that clustering of smoking, drinking, and physical inactivity is more often present in male thyroid cancer survivors [12].

Inconsistent reports from different studies of CAA and thyroid cancer are commonly based on a small number of patients with cancer involved (i.e. less than 500), restriction to certain patient sub-population (e.g. postmenopausal females) and failure to evaluate the effect modifiers (e.g. cigarette smoking, obesity, physical inactivity, etc.) [6, 12-14].

Rare studies have precisely defined thresholds of alcohol intake (i.e. amount, duration) in terms of thyroid cancer risk. Honnamurthy et al. revealed a significant influence of alcohol consumption duration, but not alcohol dependence on thyroid function tests [17]. Whang et al. report a reduction in thyroid cancer risk with decreased alcohol consumption (25 g or less) per event (i.e. mild to moderate consumption) and a drinking duration of less than 10 years, compared to never-drinkers. On the contrary, acute heavy alcohol consumption (151 g or more per event) consumption of alcohol for 31 or more years, was associated with increased risk for thyroid cancer in both men and women [13]. Our study has set the threshold of alcohol intake to moderate and higher levels, which may explain similar results in terms of a positive correlation between CAA and thyroid cancer.

The precise mechanisms by which alcohol possibly induce thyroid oncogenicity remains unclear. Whether the alcohol at certain blood levels and duration of exposure has a direct toxic effect on thyroid cells is not yet firmly established, but abnormal functioning of the hypothalamic-pituitary-thyroid axis has been observed in chronic alcoholics.

Experimental data have shown that chronic ethanol exposure in rats elevated thyroid-releasing hormone messenger RNA in hypothalamic neurons. Whether this effect, in long term, may produce hyperproliferation and/or cancerogenesis, remains unclear. (18-20)

This study has several limitations. Being a retrospective cross-sectional study, its results necessitate further validation in a wider scope, prospective study, including a larger number of patients and variables to allow more powerful statistics.

CONCLUSION

The results of our study suggest that the CAA is positively correlated with the appearance of papillary and medullar forms of thyroid carcinoma, whereas in anaplastic and follicular forms this correlation was absent. Further prospective investigations are needed to confirm these findings.

Conflict of interest: None declared.

REFERENCES

1. Connor J. Alcohol consumption as a cause of cancer. *Addiction*. 2017;112(2):222-8. PMID: 27442501, DOI: 10.1111/add.13477
2. Katherine E. Marschall, Roberta L. Hines. *Psychiatric Disease, Substance Abuse, and Drug Overdose In: Roberta L. Hines, Katherine E. Marschall, editors. Stoelting's Anesthesia and Co-Existing Disease E-Book: Elsevier Health Sciences; 2017. p. 611-34. ISBN: 0323508626, 9780323508629*
3. Kalezić N, Dimitrijević I, Leposavić L, Kocica M, Bumbasirević V, Vucetić C, et al. Postoperative cognitive deficits. *Srp Arh Celok Lek*. 2006;134(7-8):331-8. PMID: 17009615, DOI: 10.2298/sarh0608331k
4. Flórez G, Espandian A, Villa R, Sáiz PA. Clinical implications of cognitive impairment and alcohol dependence. *Adicciones*. 2019;31(1):3-7. *Adicciones*. 2019;31(1):3-7. PMID: 30699230, DOI: 10.20882/adicciones.1284
5. Rachdaoui N, Sarkar DK. Pathophysiology of the Effects of Alcohol Abuse on the Endocrine System. *Alcohol Res*. 2017;38(2):255-76. PMID: 28988577, PMCID: PMC5513689
6. Wang X, Cheng W, Li J, Zhu J. A meta-analysis of alcohol consumption and thyroid cancer risk. *Oncotarget*. 2016;7(34):55912-23. PMID: 27385005, PMCID: PMC5342461, DOI: 10.18632/oncotarget.10352
7. Wiltshire JJ, Drake TM, Uttley L, Balasubramanian SP. Systematic Review of Trends in the Incidence Rates of Thyroid Cancer. *Thyroid*. 2016;26(11):1541-52. PMID: 27571228, DOI: 10.1089/thy.2016.0100
8. Kitahara CM, Sosa JA. The changing incidence of thyroid cancer. *Nat Rev Endocrinol*. 2016;12(11):646-53. PMID: 27418023 DOI: 10.1038/nrendo.2016.110
9. World Health Organization. *Global Status Report on Alcohol and Health 2018: World Health Organization; 2019. ISBN: 9241565632, 9789241565639*
10. Statista. Prevalence of alcoholism in Serbia in 2016, by gender and type 2016 [cited 2020]. Available from: <https://www.statista.com/statistics/983902/serbia-alcoholism-prevalence-by-gender-and-type/>.
11. WHO. Alcohol-attributable fractions, all-cause deaths. 2016 [cited 2020]. Available from: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/alcohol-attributable-fractions-all-cause-deaths\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/alcohol-attributable-fractions-all-cause-deaths(-)).
12. Yoon J, Park B. Factors Associated with Health Behaviors in Thyroid Cancer Survivors. *J Cancer Prev*. 2020;25(3):173-80. PMID: 33033711, PMCID: PMC7523035, DOI: 10.15430/JCP.2020.25.3.173
13. Hwang Y, Lee KE, Weiderpass E, Park YJ, Chai YJ, Kwon H, et al. Acute High-Dose and Chronic Lifetime Exposure to Alcohol Consumption and Differentiated Thyroid Cancer: T-CALOS Korea. *PLoS One*. 2016;11(3):e0151562. PMID: 26985827, PMCID: PMC4795733, DOI: 10.1371/journal.pone.0151562
14. Choi SW, Ryu SY, Han MA, Park J. The association between the socioeconomic status and thyroid cancer prevalence; based on the Korean National Health and Nutrition Examination Survey 2010-2011. *J Korean Med Sci*. 2013;28(12):1734-40. PMID: 24339702, PMCID: PMC3857368, DOI: 10.3346/jkms.2013.28.12.1734
15. Zivaljevic V, Vlajinac H, Marinkovic J, Sipetic S, Paunovic I, Diklic A, et al. Case-Control Study of Anaplastic Thyroid Cancer: Papillary Thyroid Cancer Patients as Controls. *The Endocrinologist*. 2010;20(6):308-11. DOI: 10.1097/TEN.0b013e3181fd02f2
16. Sokić SI, Adanja BJ, Vlajinac HD, Janković RR, Marinković JP, Zivaljević VR. Risk factors for thyroid cancer. *Neoplasma*. 1994;41(6):371-4. PMID: 7870223
17. Honnamurthy JB, Shivashankara AR, Avinash SS, John Mathai P, Malathi M. Effect of Interaction Between Duration of Alcohol Consumption and Alcohol Dependence on Thyroid Function Test: Cross-Sectional Observational Study. *Indian J Clin Biochem*. 2018;33(1):61-8. PMID: 29371771, PMCID: PMC5766464, DOI: 10.1007/s12291-017-0645-6
18. Balhara YP, Deb KS. Impact of alcohol use on thyroid function. *Indian J Endocrinol Metab*. 2013;17(4):580-7. PMID: 23961472, PMCID: PMC3743356, DOI: 10.4103/2230-8210.113724
19. Zoeller RT, Fletcher DL, Simonyl A, Rudeen PK. Chronic ethanol treatment reduces the responsiveness of the hypothalamic-pituitary-thyroid axis to central stimulation. *Alcohol Clin Exp Res*. 1996;20(5):954-60. PMID: 8865974, DOI: 10.1111/j.1530-0277.1996.tb05277.x
20. Hegedüs L. Decreased thyroid gland volume in alcoholic cirrhosis of the liver. *J Clin Endocrinol Metab*. 1984;58(5):930-3. PMID: 6707193, DOI: 10.1210/jcem-58-5-930.

Table 1. Distribution of patients by diseases

Disease	n (%)
Thyroid cancer group	
I – Papillary cancer	380 (75.7)
II – Medullary cancer	102 (20.3)
III – Anaplastic cancer	10 (2)
IV – Follicular cancer	10 (2)
Total	502 (100)
Control group	
Nodular goiter	233 (38.8)
Multinodular goiter	337 (56.2)
Toxic nodular goiter	30 (5)
Total	600 (100)

Paper accepted

Table 2. Distribution of patients by age and sex

Characteristics	Thyroid cancer (n = 502) n (%)	Control group (n = 600) n (%)	p
Age			
≤30 years	67 (13.3)	41 (6.8)	< 0.001
> 30 years	435 (86.7)	559 (93.2)	n.s.
Sex			
Male	99 (19.7)	70 (11.7)	< 0.001
Female	403 (80.3)	530 (88.3)	n.s.

n.s. – non-significant

Paper accepted

Table 3. Distribution of patients by age and sex according to the type of thyroid cancer

Parameter (p-value)	Control n = 600 n (%)	I – Papillary n = 380 n (%)	II – Medullary n = 102 n (%)	III – Anaplastic n = 10 n (%)	IV – Follicular n = 10 n (%)
Age					
≤30	41 (6.8)	55 (14.5)	11(10.8)	0 (0)	1 (10)
> 30	559 (93.2)	325 (85.5)	91 (89.2)	10 (100)	9 (90)
p		(0.000)	(0.159)	(0.392)	(0.695)
Sex					
Male	70 (11.7)	71 (18.7)	27 (26.5)	1 (10)	0 (0)
Female	530 (88.3)	309 (81.3)	75 (73.5)	9 (90)	10 (100)
p		(0.002)	(0.000)	(0.871)	(0.251)

Table 4. Alcohol abuse among thyroid cancer patients

Alcohol abuse	All cancers n = 502 n (%)	I – Papillary n = 380 n (%)	II – Medullary n = 102 n (%)	III – Anaplastic n = 10 n (%)	IV – Follicular n = 10 n (%)
Yes	8 (1.6)	6 (1.6)	2 (2)	0 (0)	0 (0)
No	494 (98.4)	374 (98.4)	100 (98)	10 (100)	10 (100)
p*	< 0.001	0.034	0.044	0.001	0.001

*Statistical significance was measured against the control group incidence of alcohol abuse;

bold – significantly higher incidence;

bold-italic – significantly lower incidence