



**СРПСКИ АРХИВ**  
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО  
**SERBIAN ARCHIVES**  
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: [office@srpskiarhiv.rs](mailto:office@srpskiarhiv.rs), Web address: [www.srpskiarhiv.rs](http://www.srpskiarhiv.rs)

**Paper Accepted\***

**ISSN Online 2406-0895**

**Original Article / Оригинални рад**

Milanka Tatić<sup>1,2,†</sup>, Dragan Lađinović<sup>2</sup>, Sanja Skeledžija-Mišković<sup>2</sup>, Ranko Zdravković<sup>3</sup>, Dragana Radovanović<sup>1,2</sup>, Ferenc Vicko<sup>1,2</sup>, Goran Malenković<sup>1,2</sup>, Jelena Vučković Karan<sup>3</sup>, Arsen Uvelin<sup>1,4</sup>

**Efficiency and safety of intrathecal morphine for analgesia  
after hysterectomy**

Ефикасност и сигурност примене интратекалног морфина за аналгезију  
након хистеректомије

<sup>1</sup>University of Novi Sad, Faculty of Medicine, Novi Sad, Serbia;

<sup>2</sup>Institute of Oncology of Vojvodina, Sremska Kamenica, Serbia;

<sup>3</sup>Institute for Cardiovascular Diseases of Vojvodina, Sremska Kamenica, Serbia;

<sup>4</sup>Clinical Center of Vojvodina, Emergency Center, Novi Sad, Serbia

**Received: March 12, 2019**

**Revised: October 8, 2019**

**Accepted: October 25, 2019**

**Online First: November 1, 2019**

**DOI: <https://doi.org/10.2298/SARH190312116T>**

\* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal.

The date the article was made available online first will be carried over.

**†Correspondence to:**

Milanka TATIĆ

Oncology Institute of Vojvodina,

Hajduk Veljkova 10, Sremska Kamenica, Serbia

Email: [milanka.tatic@mf.uns.ac.rs](mailto:milanka.tatic@mf.uns.ac.rs)

## Efficiency and safety of intrathecal morphine for analgesia after hysterectomy

### Ефикасност и сигурност примене интратекалног морфина за аналгезију након хистеректомије

#### SUMMARY

**Introduction/Objective** This prospective, randomized study was aimed at determining the efficiency and safety of the administration of intrathecal morphine chloride combined with the local anesthetic levobupivacaine given to female patients subjected to hysterectomy to ensure postoperative analgesia.

**Methods** The study sample consisted of 50 patients who were to undergo hysterectomy with adnexectomy and they were divided into two groups of 25 each. The group A patients were given the combination of 0.3 mg of morphine chloride with 1.7 ml of 0.5% levobupivacaine immediately before the surgery, whereas the group B patients were administered 5 mg of morphine chloride iv before the end of surgery and after the surgery at certain time intervals. The postoperative pain was assessed at the first, sixth, 12th, and 24th hours by Numeric Rating Scale (NRS). Side effects, such as nausea, vomiting, itching and respiratory depressions were followed as well.

**Results** The postoperative pain was less expressed at any time interval both at rest and on exertion in group A ( $p < 0.001$ ) and therefore the need for additional analgesia was less in group A ( $p < 0.001$ ). The subjective feeling of satisfaction with postoperative analgesia was statistically significant in group A ( $p < 0.001$ ).

**Conclusion** Intrathecal administration of morphine chloride combined with levobupivacaine ensures better postoperative analgesia after hysterectomy than intravenous morphine chloride, their side effects being equally frequent.

**Keywords:** Hysterectomy; intrathecal morphine; postoperative analgesia

#### САЖЕТАК

**Увод/циљ** Циљ ове проспективне, рандомизоване студије је да детерминише ефикасност и сигурност примене интратекалног морфин хлорида у комбинацији са локалним анестетиком левобупивакаином, за обезбеђивање постоперативне аналгезије код пацијенткиња које се подвргавају хистеректомији.

**Метод** Педесет пацијенткиња којима је индикована хистеректомија са аднексектомијом и лимфаденектомијом је подељено у две групе од по 25 испитаница. Испитаницама из Групе А је непосредно пре операције интратекално апликована комбинација 0,3 mg морфин хлорида и 1,7 ml 0,5% левобупивакаина, док су оне из Групе Б интравенски добијале 5 mg морфин хлорида пре завршетка операције и постоперативно у одређеним временским интервалима. Постоперативни бол је оцењен првог, шестог 12. и 24. сата, уз помоћ *Numeric Rating Scale (NRS)*. Праћена су и нежељена дејства у виду мучнине, повраћања, свраба и респираторне депресије.

**Резултати** Постоперативни бол је у свим интервалима праћења, како у миру тако и приликом напора, био мање изражен у Групи А ( $p < 0,001$ ), због тога је и потреба за додатном аналгезијом била мања у групи А ( $p < 0,001$ ). Субјективни осећај задовољства постоперативном аналгезијом је статистички значајан, у корист Групе А ( $p = 0,001$ ).

**Закључак** Интратекално давање морфин хлорида у комбинацији са левобупивакаином обезбеђује бољу постоперативну аналгезију након хистеректомије, у односу на ив морфин хлорид, уз подједнаку учесталост нежељених дејстава.

**Кључне речи:** Хистеректомија; интратекални морфин; постоперативна аналгезија

## INTRODUCTION

Acute pain is the body's mechanism to signal tissue injury and danger [1]. The intensity of postoperative pain after hysterectomy ranges from moderate to severe. The efficient pain control leads to the earlier mobilization of patients and their faster recovery, thus leading to shorter hospitalization and lower treatment expenses [2–5]. Controlling acute

pain after surgery is important not only in the immediate postoperative phase but also to prevent chronic postsurgical pain, which can develop in as many as 10% of patients [6].

The aim is to achieve good analgesia with minimum side effects. Currently, it cannot be said that there is an ideal mode of analgesia. Intravenous opioids and other analgesics are most frequently used drugs to alleviate the postoperative pain during the first 24 hours. The use of opioids for patients who have surgery presents a particularly challenging problem requiring clinicians to balance managing acute pain in the postoperative period and minimising the risks of persistent opioid use after surgery [7].

Another good technique is epidural analgesia. A systemic review has shown that the continuous epidural analgesia is superior to the patient controlled analgesia (PCA) in the patients having had a major intra-abdominal surgery [8]. However, this procedure is a bit more invasive because a catheter must be inserted, which has to remain in the epidural space for at least 24 to 48 hours.

Intrathecal administration of local anesthetics with small doses of opioids is an attractive analgesic technique because the drugs are given directly into the liquor, close to the structures of the central nervous system through which these drugs act [9]. This procedure is simple, easy to be performed, with a relatively low risk of failure. It is expected to decrease the postoperative pain as well as the necessity for additional analgesia. It has been shown to be effective in the alleviation of postoperative pain after prostatectomy [10], transurethral resection of the prostate [11] and hepatectomy [12]. The most frequently used opioid is morphine. Analgesic effect of morphine administered intrathecally may persist for up to 48 hours [13]. A disadvantage of this technique is the occurrence of side effects such as nausea, vomiting, itching, urine retention, and respiratory depression, being the most severe complication [14, 15, 16, 17].

This prospective, randomized study was aimed at determining the efficiency and safety of administration of intrathecal morphine combined with a local anesthetic given to ensure analgesia in the patients to undergo hysterectomy.

## METHODS

This prospective, randomized study included 50 patients from the elective surgical program performed at the Department of Surgical Oncology, Institute of Oncology of Vojvodina in Sremska Kamenica. The patients underwent hysterectomy with bilateral adnexectomy and lymphadenectomy. The study, which had been approved by the Ethics Committee of the Institute of Oncology of Vojvodina, was conducted in the period from September 1, 2017 to February 15, 2018. The patients were divided into two groups according to the applied technique of postoperative analgesia:

Group A (25 subjects) – the patients having been given morphine chloride intrathecally in the combination with levobupivacaine before the surgery;

Group B (25 subjects) – the patients given morphine chloride intravenously before the end of surgery (when suturing began) and after the surgery.

The inclusion criteria were:

- Age  $\geq$  18 years
- ASA classification I or II
- Diagnosed neoplasm of the uterine body or cervix

The exclusion criteria were:

- Allergic reaction to morphine, local anesthetic and Metamizole
- Age  $<$  18 years

The patients meeting criteria were given both verbal and written details about the study method and aims as well as about the technique of premedication and anesthesia for the

planned surgical intervention. The ones who had given their written consent to participate were included in the study sample.

The group A patients were given 0.3 mg of morphine chloride intrathecally with 1.7 ml of 0.5% levobupivacaine (L3–L4 space, 26G needle) 15 minutes before anesthesia, and the group B patients were intravenously given 5 mg of morphine chloride before the end of surgery when suturing began. Anesthesia in both groups of patients was general and balanced. All patients were premedicated with midazolam (0.05 mg/kg iv), and propofol (1.5 mg/kg) was used to introduce anesthesia along with opioid analgesic fentanyl (1.5 µg/kg). Rocuronium (1 mg/kg) was administered for muscle relaxation for intubation as well as during the surgery at a dose adequate to maintain the relaxation. Anesthesia was maintained with sevoflurane (1 MAC). The lung ventilation was ensured with the gas mixture O<sub>2</sub>:N<sub>2</sub>O 40:60 so that EtCO<sub>2</sub> < 38 mmHg. Electrocardiogram, blood pressure, heart frequency, SpO<sub>2</sub>, EtCO<sub>2</sub> and body temperature were monitored during the surgery. Boluses of fentanyl (50 µg) and of rocuronium (10 mg) were given when necessary. Anesthetics were administered in such a way so to provide satisfactory anesthesia, blood pressure values and heart frequency in the values ± 30% as compared with the values prior to anesthesia. Drugs for reversal of neuromuscular blockade (neostigmine 0.02 mg/kg and atropine 0.01 mg/kg) were given at the end of surgery. In addition, metoclopramide 10 mg intravenous was administered at the end of surgery, immediately before the patients woke up, to prevent nausea and vomiting. The group B patients were given metoclopramide at 8 hours in first 24 hours after the surgery.

The group A patients were given intravenous 2.5 g (maximum 5g/24h) of Metamizol when necessary for postoperative analgesia in case of pain exceeding 4 according to the numeric rating scale (NRS), whereas morphine chloride (5 mg iv/6-8 hours) and metamizol 2.5 g (maximum 5 g/24 h) were administered alternatively after the surgery to the group B patients and 2–5 mg of intravenous morphine chloride was given when pain exceeded 4

according to NRS. The dose of additional morphine chloride in group B was dependent on the age and body weight of the patients, according to the protocol of our clinic.

Numeric values of postoperative pain intensity assessed according to NRS at rest, while coughing and active moving were recorded at intervals of 1h, 6h, 12h, and 24h. These values ranged from 0 to 10 (0 – no pain, 10 – the most severe possible pain). In addition, the patients were asked to keep records of their subjective feeling of satisfaction (SFS) with analgesia, assessing it as: 1 – poor, 2 – medium, 3 – good, and 4 – excellent. Complications such as nausea, vomiting, itching, and respiratory depression were also recorded. All patients had urinary catheter so we could not follow urinary retention.

The data were analyzed and processed by IBM SPSS statistics 10.0 software and given in tables and figures created in Word and Excel Microsoft Office 2003 packs. The results were presented using standard statistical methods: frequency (f), arithmetic means ( $\bar{x}$ ), standard deviation (SD), value intervals (maximum and minimum), and percentages (%). The patients' characteristics were compared with Student's t test and  $\chi^2$  test,  $p < 0.05$  being the statistically significant value.

## RESULTS

The study sample consisted of 50 female patients divided into two groups of 25 each. There were no statistically significant differences in age, body weight, ASA (American Society of Anesthesiologists) classification and the length of surgery between the group A and B (Table 1).

Table 2 and Figure 1 show the mean values of pain at rest assessed according to NRS during the period of 24 hours after the surgery. A statistically significant difference was recorded.

The group A patients, who had received morphine chloride intrathecally, had the values

lower than the numeric values found in the group B patients, and the difference was even more expressed after the 1<sup>st</sup> and 6<sup>th</sup> hour than after the 12<sup>th</sup> and 24<sup>th</sup> hour.

Table 3 and Figure 2 show the mean values of pain on exertion assessed according to NRS during the period of 24 hours after the surgery, with a statistically significant difference to the benefit of the group A patients.

The results of cross comparison of the need for additional analgesia an hour after the end of surgery are shown in Table 4. As it can be seen, only two patients, who had received morphine chloride intrathecally, needed additional analgesia, whereas the majority the group B patients needed additional analgesia. This difference was highly statistically significant ( $\chi^2 = 12.834$ ;  $p < 0.001$ ).

Tables 5, 6, 7 and 8 show the development of nausea as a side effect of analgesic. No statistically significant difference was found between the groups in various time intervals during the first 24 hours after the surgery.

There was no statistically significant difference in vomiting as a side effect of analgesic during the period of 12 hours after the surgery, as shown in Tables 9, 10, and 11. None of the patients had this problem 24 hours after surgery.

Possible side effects of opioid analgesia are itching and respiratory depression. However, none of our patients developed any of them.

Table 12 shows the subjective feeling of satisfaction with analgesia. Our patients assessed analgesia as: 1 – poor, 2 – medium, 3 – good, and 4 – excellent.

## DISCUSSION

There were not statistically significant differences between the two groups regarding age, body weight, ASA classification and the length of surgery. We opted for the dose of 0.3 mg of morphine chloride based on a meta-analysis and a review [18, 19].

The results of pain assessment show that pain, both at rest and on exertion, was statistically significantly less expressed in the patients having received morphine chloride intrathecally (Group A). The difference was even more expressed immediately after the surgery. Only two patients from group A asked for additional analgesia after the first postoperative hour and they received 2.5 gr of intravenous Metamizole. The patients from this group required additional analgesia only 27 hours after the moment of intrathecal administration of morphine chloride. However, as many as 15 patients from group B asked for additional analgesia after the first hour and they received 2-5 mg of morphine chloride iv. The average consumption of morphine chloride after 24 hours was 21.7 mg in group B, which confirms the statement made El Sherif et al who claim that intrathecal morphine has about 100 times more potency than intravenous morphine [20]. Jakobson L et al believe that intrathecally administered morphine yields effective long-lasting analgesia (over 20 hours) and that the dose from 0.3 mg to 1 mg should ensure good analgesia without the major complication – respiratory depression [21]. A team of Danish researchers has found in their study that it takes 27 hours on average from the administration of 0.1 mg and 0.2 mg of intrathecal morphine given for Cesarean section until the first postoperative analgesic [22]. They recommend 0.1 mg as a dose of choice. Slappendel R et al have compared various doses of intrathecal morphine in the patients having had hip replacement and their conclusion is that the dose of 0.1 mg ensures good analgesia with minimum side effects [23]. An Egyptian team of researchers performed a study on safety and analgesic efficiency of intrathecal morphine (0.2, 0.5, and 1 mg) in the patients undergoing a major abdominal surgery for cancer. They concluded that 1 mg of morphine resulted in superior analgesia 48 hours after surgery in comparison with 0.2 and 0.5 mg, without a significant difference in the frequency of side effects [24]. Urban MK and his colleagues have found that the dose of 20 µg/kg of intrathecal morphine considerably decreases the need for additional postoperative



analgesia after lumbar fusion surgery [25]. It can be concluded that the dose of morphine, which is necessary for good postoperative analgesia, depends on the type of surgical procedure.

The following side effects were monitored: nausea, vomiting, itching, and respiratory depression. Since all patients had the urinary catheter, urinary retention, as a side effect, was not followed.

Nausea was less expressed in the patients who had received morphine chloride intrathecally; however, the difference was not statistically significant. Vomiting was equally frequent in both groups. It should be emphasized that the group A patients were given metoclopramide at the end of surgery as well as in the postoperative period if necessary because of nausea and vomiting, whereas the group B patients were given metoclopramide at the end of surgery and 8 hours after the surgery. Similar results have been reported in other studies as well [26, 27].

None of our patients reported itching nor was respiratory depression observed as a side effect of morphine. Itching is often mentioned in literature as a frequent side effect of intrathecal morphine, particularly in obstetrics, where gestation hormones may lead to certain changes on opioid receptors [28]. Some researchers from Holland have shown that intrathecal administration of morphine for laparoscopic colon resection is followed with itching more often [29].

Based on their study Jacobson L et al have concluded that nausea, vomiting, and itching develop irrespectively of the dose given, whereas respiratory depression occurs after a dose exceeding 1 mg [21]. In 2009, Gehling M and Tryba M performed a meta-analysis of randomized controlled studies, which had dealt with the application of intrathecal morphine combined with spinal anesthesia [18]. Their analysis included the following surgical procedures: Cesarean section, orthopedic surgery, gynecologic surgery, arthroscopy,

transurethral prostatectomy, and hemorrhoidectomy. Morphine was administered in the doses ranging from 0.025 mg to 2.5 mg. It has been found that the doses  $\geq 0.3$  mg increase the incidence of itching. The incidence of respiratory depression in the patients given a dose  $< 0.3$  mg was 1%, whereas it was 9% in case of a dose  $\geq 0.3$  mg. The authors believe that intrathecal morphine requires prophylaxis and treatment of side effects. In addition, the respiratory function must be continuously followed for 24 hours. The same must be done in the patients who received systemic opioids. There is no evidence to corroborate the extended monitoring of patients receiving low doses of morphine.

The subjective feeling of satisfaction with analgesia may have to be the most important factor when opting for postoperative analgesia. As many as 96% of the group A patients assessed analgesia as excellent, whereas only 20% of those from group B were of that opinion, which is in discrepancy with a similar study which showed almost equal satisfaction of patients with analgesia [29].

At the end, we should also mention a study performed by an American team of researchers on the pregnant women who underwent Cesarean section delivery [30]. The study sample was divided according to the wish of the pregnant women when asked whether they wanted 0.1 mg or 0.2 mg of morphine after having been adequately informed about the advantages and shortcomings of such analgesia. The need for additional analgesia was found to be bigger in the patients who had received 0.2 mg of morphine. Therefore, it can be concluded that postoperative analgesia is a multifactorial problem, where the subjective feeling of the patients themselves and their fear of surgery, postoperative pain, and side effects are of uppermost importance.

Potential limitations to our study are a low number of subjects and a subjective assessment of pain intensity according to NRS. It is difficult to observe a rare side effect in a small study sample. A similar study including a higher number of subjects would contribute

to the strength of conclusions. Urinary retention could not be followed as a side effect in this study because of routine placement of urinary catheter.

## CONCLUSION

According to our study, it can be concluded that the combination of 0.3 mg of morphine chloride and 1.7 ml of 0.5% levobupivacaine provides very good analgesia after hysterectomy with adnexectomy and lymphadenectomy lasting for 24 hours after the surgery. Side effects do not differ from those associated with intravenous administration of opioid. Nausea and vomiting can be treated with antiemetics. Although respiratory depression was not observed in our study sample, intrathecal morphine must be administered with great caution and respiratory function must be monitored during the first 24 hours. Finally, our study has shown that the patients who received morphine chloride intrathecally were more satisfied than those who were given morphine intravenously.

**Conflict of interest:** None declared.

**REFERENCES**

1. Loke MF, Wei H, Yeo J, Sng BL, Sia AT, Tan EC. Deep sequencing analysis to identify novel and rare variants in pain-related genes in patients with acute postoperative pain and high morphine use. *J Pain Res.* 2019; 12:2755-2770. doi: 10.2147/JPR.S213869.
2. Apfelbaum JL, Chen C, Mehta SS, Gan TJ. Postoperative pain experience: results from a national survey suggest postoperative pain continues to be undermanaged. *Anesth Analg.* 2003; 97(2): 534–540. PMID: 12873949
3. Azari L, Santoso JT, Osborne SE. Optimal pain management in total abdominal hysterectomy. *Obstet Gynecol Surv.* 2013; 68(3):215–227. doi.org/10.1097/OGX.0b013e31827f5119
4. Koning MV, Teunissen AJW, van der Harst E, Ruijgrok EJ, Stolker RJ. Intrathecal Morphine for Laparoscopic Segmental Colonic Resection as Part of an Enhanced Recovery Protocol: A Randomized Controlled Trial. *RegAnesth Pain Med.* 2018; 43(2):166-173. doi: 10.1097/AAP.0000000000000703.
5. Farzi F, NaderiNabi B, Mirmansouri A, Fakoor F, Atrkar Roshan Z, Biazar G et al. Postoperative Pain After Abdominal Hysterectomy: A Randomized, Double-Blind, Controlled Trial Comparing the Effects of Tramadol and Gabapentin as Premedication. *Anesth Pain Med.* 2016; 6(1):e32360. doi: 10.5812/aapm.32360.
6. The Lancet. Best practice in managing postoperative pain. *Lancet.* 2019; 393(10180):1478. doi: 10.1016/S0140-6736(19)30813-X.
7. Glare P, Aubrey KR, Myles PS. Transition from acute to chronic pain after surgery. *Lancet.* 2019; 393(10180):1537-1546. doi: 10.1016/S0140-6736(19)30352-6.
8. Werawatganon T, Charuluxananan S. Patient controlled intravenous opioid analgesia versus continuous epidural analgesia for pain after intra-abdominal surgery. *Cochrane Database Syst Rev.* 2005; 1:CD004088. doi. org/10.1002/14651858.CD004088.pub2
9. Ferraz S, Caria T, Da Silva AV, Candeias MJ, Cenicante T. Persistent Hypothermia and Excessive Sweating Following Intrathecal Morphine Administration in a Teenage Boy: A Case Report. *Anesth Pain Med.* 2018; 8(1):e66724. doi: 10.5812/aapm.66724.
10. Andrieu G, Roth B, Ousmane L, Castaner M, Petillot P, Vallet B et al. The efficacy of intrathecal morphine with or without clonidine for postoperative analgesia after radical prostatectomy. *Anesth Analg.* 2009; 108: 1954–1957. doi: 10.1213/ane.0b013e3181a30182.
11. Sakai T, Use T, Shimamoto H, Tanabe T, Terao Y, Sumikawa K. Minidose (0.05 mg) intrathecal morphine provides effective analgesia after transurethral resection of the prostate. *Can J Anaesth.* 2003; 50: 1027–1030. doi: 10.1007/s00540-009-0817-5.
12. Ko JS, Choi SJ, Gwak MS, Kim GS, Ahn HJ, Kim JA et al. Intrathecal morphine combined with intravenous patient-controlled analgesia is an effective and safe method for immediate postoperative pain control in live liver donors. *Liver Transpl.* 2009; 15: 381–389. doi: 10.1002/lt.21625.
13. Gehling MH, Luesebrink T, Kulka PJ, Tryba M. The effective duration of analgesia after intrathecal morphine in patients without additional opioid analgesia: a randomized double-blind multicentre study on orthopaedic patients. *Eur J Anaesthesiol.* 2009; 26: 683–688. doi: 10.1097/EJA.0b013e328329b045.
14. Rathmell JP, Lair TR, Nauman B. The role of intrathecal drugs in the treatment of acute pain. *Anesth Analg.* 2005; 101:S30-43. doi: 10.1213/01.ANE.0000177101.99398.22.
15. Lui MW, Li TKT, Lui F, Ong CYT. A randomised, controlled trial of rectus sheath bupivacaine and intrathecal bupivacaine, without or with intrathecal morphine, vs. intrathecal bupivacaine and morphine after caesarean section. *Anaesthesia.* 2017; 72(10):1225-1229. doi: 10.1111/anae.13998.

16. Woods JM, Lim AG. Prevalence and management of intrathecal morphine-induced pruritus in New Zealand Māori healthcare recipients. *Br J Pain*. 2018; 12(1):20-25. doi: 10.1177/2049463717719773.
17. Ladha KS, Kato R, Tsen LC, Bateman BT, Okutomi T. A prospective study of post-cesarean delivery hypoxia after spinal anesthesia with intrathecal morphine 150µg. *Int J Obstet Anesth*. 2017; 32:48-53. doi: 10.1016/j.ijoa.2017.06.003.
18. Gehling M, Tryba M. Risks and side-effects of intrathecal morphine combined with spinal anaesthesia: a meta-analysis. *Anaesthesia*. 2009; 64(6):643-51. PMID: 19462494.
19. MugabureBujedo B. A clinical approach to neuraxial morphine for the treatment of postoperative pain. *Pain Res Treat*. 2012; 2012:11. doi:10.1155/2012/612145.
20. El Sherif FA, Othman AH, Abd El-Rahman AM, Taha O. Effect of adding intrathecal morphine to a multimodal analgesic regimen for postoperative pain management after laparoscopic bariatric surgery: a prospective, double-blind, randomized controlled trial. *Br J Pain*. 2016; 10(4):209-216. doi:10.1177/2049463716668904.
21. Jacobson L, Chabal C, Brody MC. A dose-response study of intrathecal morphine: efficacy, duration, optimal dose, and side effects. *Anesth Analg*. 1988; 67(11):1082-8. PMID: 3189898.
22. Dahl JB, Jeppesen IS, Jørgensen H, Wetterslev J, Møiniche S. Intraoperative and postoperative analgesic efficacy and adverse effects of intrathecal opioids in patients undergoing cesarean section with spinal anesthesia: a qualitative and quantitative systematic review of randomized controlled trials. *Anesthesiology*. 1999; 91(6):1919-27.
23. Slappendel R, Weber EW, Dirksen R, Gielen MJ, van Limbeek J. Optimization of the dose of intrathecal morphine in total hip surgery: a dose-finding study. *Anesth Analg*. 1999; 88(4):822-6. doi: 10.1213/0000539-199904000-00026.
24. Fares KM, Mohamed SA, Abdel-Ghaffar HS. High dose intrathecal morphine for major abdominal cancer surgery: a prospective double-blind, dose-finding clinical study. *Pain Physician*. 2014; 17(3):255-64.
25. Urban MK, Jules Elysee K, Urquhart B, Cammisa FP, BoachieAdjei O. Reduction in postoperative pain after spinal fusion with instrumentation using intrathecal morphine. *Spine*. 2002; vol. 27:535-7. PMID: 11880840.
26. Jun JH, Kim GS, Lee JJ, Ko JS, Kim SJ, Jeon PH. Comparison of intrathecal morphine and surgical-site infusion of ropivacaine as adjuncts to intravenous patient-controlled analgesia in living-donor kidney transplant recipients. *Singapore Med J*. 2017; 58(11):666-673. doi: 10.11622/smedj.2017077.
27. Kim HC, Bae JY, Kim TK, Jeon Y, Min JJ, Goo EK et al. Efficacy of intrathecal morphine for postoperative pain management following open nephrectomy. *J Int Med Res*. 2016;44(1):42-53. doi: 10.1177/0300060515595650.
28. Hindle A. Intrathecal opioids in the management of acute postoperative pain. *Continuing Education in Anaesthesia Critical Care & Pain*. 2008; Volume 8, Issue 3:81–85.
29. Koning MV, Teunissen AJW, van der Harst E, Ruijgrok EJ, Stolker RJ. Intrathecal Morphine for Laparoscopic Segmental Colonic Resection as Part of an Enhanced Recovery Protocol: A Randomized Controlled Trial. *RegAnesth Pain Med*. 2018; 43(2):166-173. doi: 10.1097/AAP.0000000000000703.
30. Carvalho B, Mirza F, Flood P. Patient choice compared with no choice of intrathecal morphine dose for caesarean analgesia: a randomized clinical trial. *Br J Anaesth*. 2017; 118(5):762-771. doi: 10.1093/bja/aex039.

**Table 1.** Patient characteristics

Parameters	Group A	Group B	p
Age (years $\pm$ SD)	56.6 $\pm$ 11.1	57.7 $\pm$ 9.8	0.707
Weight (kg $\pm$ SD)	73.6 $\pm$ 13.2	75.5 $\pm$ 13.8	0.624
ASA I n (%)	2 (8)	3 (12)	1.000
ASA II n (%)	23 (92)	22 (88)	
Duration of the operation (min $\pm$ SD)	174.2 $\pm$ 38.8	150 $\pm$ 49.7	0.061

Paper accepted

**Table 2.** Mean NRS (at rest) one, six, 12 and 24 hours after surgery

Parameters	Group A				Group B				t-test	p
	Mean	Min	Max	Standard deviation	Mean	Min	Max	Standard deviation		
<b>NRS 1h</b>	1.04	0	5	1.485	4.32	0	9	1.97	-6.640	< 0.001
<b>NRS 6h</b>	0.08	0	1	0.277	2.44	0	5	1.356	-8.523	< 0.001
<b>NRS 12h</b>	0.04	0	1	0.2	1.44	0	3	1	-6.842	< 0.001
<b>NRS 24h</b>	0	0	0	0	1.2	0	4	1	-6.000	< 0.001

NRS – Numeric Rating Scale

**Table 3.** Mean NRS on exertion, one, six, 12 and 24 hours after surgery

Parameters	Group A				Group B				t-test	p
	Mean	Min	Max	Standard deviation	Mean	Min	Max	Standard deviation		
<b>NRS 1h</b>	1.48	0	6	2.044	6	2	10	1.848	-8.201	< 0.001
<b>NRS 6h</b>	0.20	0	3	0.707	3.88	2	6	1.301	-12.424	< 0.001
<b>NRS 12h</b>	0.08	0	2	0.400	2.76	1	5	0.879	-13.87	< 0.001
<b>NRS 24h</b>	0.00	0	0	0.000	2.24	1	5	0.969	-11.552	< 0.001

NRS – Numeric Rating Scale



**Table 4.** Need for additional analgesia one hour after surgery

<b>Additional analgesia after 1h</b>	<b>Total</b>	<b>Group A</b>	<b>Group B</b>	<b><math>\chi^2</math></b>	<b>p</b>
<b>No n (%)</b>	33 (66)	23 (92)	10 (40)	12.834	< 0.001
<b>Yes n (%)</b>	17 (34)	2 (8)	15 (60)		
<b>Total n (%)</b>	50 (100)	25 (100)	25 (100)		

Paper accepted

**Table 5.** Nausea one hour after surgery

Parameter		Group		$\chi^2$	p
		Group A	Group B		
Nausea	No	n	23	1.495	0.417
		%	92		
	Yes	n	2		
		%	8		
Total	n	25			
	%	100			

Paper accepted

**Table 6.** Nausea six hours after surgery

Parameters			Group		$\chi^2$	p
			Group A	Group B		
Nausea 6h	No	n	23	21	0.758	0.667
		%	92	84		
	Yes	n	2	4		
		%	8	16		
Total		n	25	25		
		%	100	100		

Paper accepted

**Table 7.** Nausea 12 hours after surgery

Parameters			Group		$\chi^2$	p
			Group A	Group B		
Nausea 12h	No	n	23	20	1.495	0.417
		%	92	80		
	Yes	n	2	5		
		%	8	20		
Total		n	25	25		
		%	100	100		

Paper accepted

**Table 8.** Nausea 24 hours after surgery

Parameters			Group		$\chi^2$	p
			Group A	Group B		
Nausea 24h	No	n	25	22	3.191	0.235
		%	100	88		
	Yes	n	0	3		
		%	0	12		
Total		n	25	25		
		%	100	100		

Paper accepted

**Table 9.** Vomiting one hour after surgery

Parameters			Group		$\chi^2$	p
			Group A	Group B		
Vomiting 1h	No	n	24	24	0.000	1.000
		%	96	96		
	Yes	n	1	1		
		%	4	4		
Total	n	25	25			
	%	100	100			

Paper accepted

**Table 10.** Vomiting six hours after surgery

Parameters			Group		$\chi^2$	p
			Group A	Group B		
Vomiting 6h	No	n	24	24	0.000	1.000
		%	96	96		
	Yes	n	1	1		
		%	4	4		
Total		n	25	25		
		%	100	100		

Paper accepted

**Table 11.** Vomiting 12 hours after surgery

Parameters			Group		$\chi^2$	p
			Group A	Group B		
Vomiting 12h	No	n	23	23	0.000	1.000
		%	92	92		
	Yes	n	2	2		
		%	8	8		
Total		n	25	25		
		%	100	100		


Paper accepted



**Table 12.** The subjective feeling of satisfaction (SFS) with analgesia


Parameters			Group		$\chi^2$	p
			Group A	Group B		
SFS	2 (medium)	n	0	4	29.684	0.001
		%	0	16		
	3 (good)	n	1	16		
		%	4	64		
	4 (excellent)	n	24	5		
		%	96	20		
Total		n	25	25		
		%	100	100		

**Figure 1.** Mean NRS at rest, one, six, 12 and 24 hours after surgery



NRS – Numeric Rating Scale

**Figure 2.** Mean NRS on exertion, one, six, 12 and 24h after surgery



NRS – Numeric Rating Scale

Paper accepted