



**СРПСКИ АРХИВ**  
ЗА ЦЕЛОКУПНО ЛЕКАРСТВО  
**SERBIAN ARCHIVES**  
OF MEDICINE

Address: 1 Kraljice Natalije Street, Belgrade 11000, Serbia

+381 11 4092 776, Fax: +381 11 3348 653

E-mail: [office@srpskiarhiv.rs](mailto:office@srpskiarhiv.rs), Web address: [www.srpskiarhiv.rs](http://www.srpskiarhiv.rs)

**Paper Accepted\***

**ISSN Online 2406-0895**

**Case Report / Приказ болесника**

Vladimir Milosavljević<sup>1</sup>, Boris Tadić<sup>2,3,†</sup>, Nikola Grubor<sup>2,3</sup>, Đorđe Knežević<sup>2,3</sup>, Slavko Matic<sup>2,3</sup>

**Large choledochal cyst initially interpreted as Mirizzi syndrome –  
case report and literature review**

Циста холедоха иницијално дијагностикована као Мирицијев синдром –  
приказ болесника и преглед литературе

<sup>1</sup>Gracia Medica Polyclinic, Belgrade, Serbia;

<sup>2</sup>Clinical Centre of Serbia, First Surgical Clinic – Clinic for Digestive Surgery, Belgrade, Serbia;

<sup>3</sup>University of Belgrade, Faculty of Medicine, Belgrade, Serbia

**Received: November 12, 2019**

**Revised: December 17, 2019**

**Accepted: December 19, 2019**

**Online First: December 23, 2019**

**DOI: <https://doi.org/10.2298/SARH191112133M>**

\***Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy-edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

†**Correspondence to:**

Boris TADIĆ

Clinic for Digestive Surgery, Clinical Centre of Serbia, Dr Koste Todorovića 6, 11000 Belgrade, Serbia

E-mail: [tadicboris@yahoo.com](mailto:tadicboris@yahoo.com)

## Large choledochal cyst initially interpreted as Mirizzi syndrome – case report and literature review

Циста холедоха иницијално дијагностикована као Мирицијев синдром – приказ болесника и преглед литературе

### SUMMARY

**Introduction** Choledochal cysts are congenital anomalies manifested as focal or diffuse cystic dilatation of the bile ducts. They are mostly diagnosed in childhood. The magnetic resonance and surgical management are the gold standard diagnostic and treatment modality.

**Case outline** We present a male patient who was presumed to have Mirizzi syndrome. This presumption was discarded by additional imaging procedures and by later surgical treatment. It was confirmed that it was a large choledochal cyst filled with stones. Considering the diagnosis and good patient's general condition, we opted for surgical treatment.

**Conclusion** There are several surgical techniques that can be used in the treatment of choledochal cysts, wherein each is intended as a complete resection of cyst with histological confirmation. Operative techniques do not affect the final outcome of the treatment, but the time and extent of surgical resection, as well as any metaplasia of the epithelium, does.

**Keywords:** choledochal cyst; Todani; common hepatic duct; icterus; Mirizzi syndrome

### САЖЕТАК

**Увод** Цисте холедоха су урођене аномалије које се манифестују као фокалне или дифузне дилатације жучних водова. Најчешће се симптоматски испоље и дијагностикују током детињства. Магнетна резонанца је златни стандард за постављање дијагнозе, а хируршко лечење једини сигуран и ефикасан модалитет лечења.

**Приказ болесника** Пацијент стар 53 године примљен је на клинику због болова у трбуху, мучнине и иктеруса. Из медицинске документације се сазнаје да пацијент 10 година у назад зна за калкулозу жучне кесе, те је након иницијалне ултразвучне дијагностике постављена сумња на Мирицијев синдром. Након спроведене допунске имџинг дијагностике утврђено је да се заправо ради о великој цисти холедохуса која је испуњена масом калкулуса. Имајући у виду добро опште стање пацијента, одлучили смо се за хируршко лечење.

**Закључак** Описано је неколико оперативних техника за хируршки третман цисте холедоха. Без обзира на примењену технику циљ је да се одстрани комплетна циста и хистопатолошки прегледају маргине ресекције као и сам препарат. Избор оперативне технике не утиче на исход лечења. Обзиром на познату тенденцију метапластичне промене епитела холедоха код пацијената са цистом холедоха од суштинског је значаја оперативном лечењу приступити на време.

**Кључне речи:** циста холедоха; Тодани; холедохус; иктерус; Мирицијев синдром

### INTRODUCTION

Choledochal cysts are congenital anomalies that manifest as focal or diffuse cystic dilation of the biliary tree. It is a rare clinical entity with an incidence of 1/150.000 births approximately [1]. There is a significant female predominance with a female to male ratio of 4:1 [2]. Five types of cysts have been described, of which a choledochal cyst type I has around 68% of percentage coverage out of all subtypes [1, 2].

The majority of patients are usually diagnosed in childhood, in the first decade of life. Only 20% of cases are diagnosed in adults [3]. Symptomatology is different in children and in adults. In children, a classical triad of symptoms in the form of abdominal pain, a palpable mass in the upper right quadrant of the abdomen and obstructive jaundice frequently occur. In adults, the ailments are related to biliary and pancreatic symptoms, accompanied by

abdominal pain. The ruptures of the cysts are rare and such an occurrence is reported only in neonates [4].

In a differential diagnosis, biliary lithiasis, sclerosing cholangitis, and pancreatic pseudocyst may be taken into account. Biliary atresia is often associated with choledochal cysts and therefore should be excluded at neonatal obstructive jaundice [2, 4].

Choledochal cyst Type I treatment involves complete surgical resection with reconstruction with an isolated small bowel segment by the method of Roux-en-Y. Sometimes, the scope of surgical resection may be much more radical. When the cyst is spreading into the head of the pancreas a cephalic duodenopancreatectomy is preferred surgical option [5].

The aim of this paper is a presentation of the rare disease, the surgical technique, and the literature review.

## CASE REPORT

In our paper, we present a male patient, 53 years of age, who was admitted to our hospital because of pain in the upper abdomen, subicteric and afebrile. The initial abdominal ultrasound was performed. Distended gallbladder, with a thickened wall and with multiple stones in the lumen were seen. Choledochus was with concrements inside the lumen, and in close relationship with the gallbladder, differential diagnosis Mirizzi syndrome. A magnetic resonance imaging (MRI) and magnetic resonance imaging cholangiopancreatography (MRCP) were performed as a part of additional diagnostics, in which a large fusiform ductus choledochus cyst was seen. A large choledochal cyst (CC), 7 cm in longitudinal diameter and 4 cm in transverse diameter, was localized at about 2 cm below the primary biliary confluence, without communication with the lumen of the gallbladder, in close contact with the portal vein and with proper hepatic artery. The cyst was filled with numerous stones of different sizes (Figure 1). The laboratory check showed elevated WBC  $18 \times 10^9/L$ , CRP 65mg/L, Total bilirubin 49mg/dL, and Alkaline phosphatase 197 IU/L.

In medical history, the patient provides information that gallbladder calculus was verified by abdominal ultrasound more than 10 years ago. In addition to the examinations that were performed several times during this period, no additional diagnostic procedures were performed. The patient states that he repeatedly felt difficulties in the form of biliary colic and sometimes spontaneously resolved mild jaundice, which were treated in a conservative way.

The patient was in good general condition, so we opted for surgical treatment. Given the size, the position of the cyst and its close relations with surrounding structures, primarily vascular, we applied the open surgical approach. In terms of general endotracheal anesthesia (GETA), the abdomen was opened with the right subcostal laparotomy. After adhesiolysis and the inspection, the preoperative diagnosis was confirmed with an inflammatory block surrounding hepatoduodenal ligament. Hard adhesions were obscuring normal anatomy, cholecystectomy was performed and the careful dissection of hepatoduodenal ligament, with difficult separation of the vascular structures from the cystic structures. Resection of the common bile duct was made along with the large cyst. The upper and lower resection was at a distance of 1 cm with respect to both ends of the cyst. Resection margins have been sent to *ex tempore* histopathologic examination. In the meantime, we have done extensive lavage of bile ducts, proximal part first and then the distal part from previously resected CC. Using choledochoscope, a proximal part relative to the branch of left and right hepatic duct and then the distal stump of the resected hepatic ductus to the papilla of Vater were inspected. The finding was normal, with no residual stones. *Ex tempore* findings were negative for malignancy. Hepaticojejunostomy was performed using interrupted, monofilament, slowly absorbable suture (4/0) at about 1 cm below the biliary confluence (Figure 2). The abdomen was drained with two abdominal drains and the operating incision was reconstructed. The tissue of the cyst (Figure 3) and of the gall bladder has been sent to histopathological examination.

Histopathological examination revealed only mild inflammatory changes in the wall of the gallbladder, with no dysplasia or other significant epithelial changes. Specimen with a cyst measuring 75 × 48 mm in diameter was extensively reviewed. It was seen that areas of papillated and partly atypical epithelial proliferation are present in the bile duct, but it is coupled with poorly expressed atypia, so there were only focuses of light epithelial dysplasia (grade I–II). In the surroundings, a chronic inflammation with some of the sporadic

multiplicity of the diverticula of the same epithelium was present, as well as the light multiplication of such tubulo-glandular structures. Not one of the sections showed any invasiveness elements. Cystic part of sections showed mostly flattened and only reactively altered, but mostly non-dysplastic epithelial changes.

The postoperative course was uneventful. Abdominal drains were removed on the third postoperative day and the patient was discharged on the seventh postoperative day. A month after, an abdominal ultrasound was performed and the finding was normal, as well as laboratory analysis. Six months after surgery, an MRI and MRCP were performed showing that the anastomosis is passable and that the other findings in the abdomen were normal. The patient's condition is still monitored.

This case report was approved by the institutional ethics committee, and written consent was obtained from the patient for the publication of this case report and any accompanying images.

## DISCUSSION

The choledochal cysts (CC) were first reported by Vater and Ezler [6]. This congenital malformation is characterized by dilatation of the biliary tree without acute obstruction of the flow of bile. The cyst may be present in any part of the biliary tree. According to the volume, it may be complete or a segmented, and according to the shape, it may be saccular or fusiform [7]. In our case, it was a fusiform cyst, which was filled with numerous concretions.

Etiology of the CC, it is still unclear and there is still no clear experts consensus. The most frequently mentioned hypothesis in the current literature suggests that choledochal cysts occur after partial obstruction of the bile duct, which produces an increase in pressure in the proximal part of the bile duct, which leads to the dilatation of this part [7]. CC prevalence is much higher in Asian countries such as in Japan and predominantly occur in females [8].

The first classification of CC was announced by Alonso-Lej et al. [9] in 1959. This initial classification was completed by Todani et al. [10], wherein the choledochal cysts were classified into five types. Type I cysts represent the dilatation of the extrahepatic bile ducts.

Also, they represent the most common type according to the previously mentioned Todani classification, where the incidence is 1:1,000, compared to western countries where the incidence ranges from 1:150,000 up to 1:1,000,000 births. Type I choledochal cysts have the greatest frequency of occurrence (75–85%) compared to other types [1, 10]. Isolated cystic dilatation of the cystic duct was added to the Todani classification in 1991 as type VI [11].

Choledochal cysts are presented with different symptoms, but they can often be asymptomatic. In symptomatic patients, they are commonly presented as abdominal pain, nausea, and vomiting. In these patients, in 60–80% of cases biliary stones, cholangitis, liver abscess, and biliary cirrhosis are present [4, 12].

Ultrasound examination of the abdomen is the first diagnostic procedure, particularly in children. As an additional diagnosis, a computed tomography is used. Both methods cannot always provide sufficient information. The gold standard is magnetic resonance imaging (MRI), as well as magnetic resonance imaging cholangiopancreatography (MRCP), with an efficiency of 96–100% [1, 12]. Endoscopic retrograde cholangiopancreatography and percutaneous transhepatic cholangiography represent very reliable diagnostic procedures, but both are invasive procedures and their application is not routinely performed [12, 13].

In our case, only after completed MRI and MRCP, we were able to remove a dilemma on the possible presence of Mirizzi syndrome. There has been no communication of choledochal cystic dilation with the lumen of the gall bladder, nor any expressed compression in the relations of biliary structures, as originally seen in abdominal ultrasound examination. Since the only reliable and correct treatment modality of these biliary tree anomalies is a complete surgical resection, surgical treatment should not be delayed, especially if you take into account that these anomalies represent the premalignant condition. Malignant alteration depends on the cyst type and that percentage is the largest for Type I cysts (about 70%), then for cysts Type IV (about 20%), followed by cysts Type II (about 5%) and cysts Type III (about 5%) [14].

Within frequent cholangitis and inflammatory processes of the surrounding structures and their consequences, sometimes it is difficult to achieve complete excision of the CC, especially because of the close relationship with the blood vessels, especially the portal vein. CC sometimes spreads into the parenchyma of the pancreas and it is necessary, in order to

apply the adequate and radical surgical approach, to perform a cephalic duodenopancreatectomy with all the risks that this procedure can cause [5, 7].

After the excision of the cyst, the reconstruction can be done in two ways: hepatico-duodenal anastomosis, or hepatico-jejunal anastomosis by the method Roux-en-Y [15]. The success of operational procedures and of the selection of anastomosis is measured by the ease of implementation, as well as by short- and long-term results of the surgical treatment. The data from the current literature suggest that the success of hepaticojejunostomy is about 92% with the complication rate of 7%, compared to hepatico-duodenostomy in which complications rate about 42% [7, 15].


Surgical management can be carried out by using several surgical techniques. These include the classical operational approach, which we also apply, then minimally-invasive or laparoscopic access, and the most modern, robotic-assisted surgical approach [16, 17].

Depending on technical equipment and on training of the surgical team, in institutions where this type of surgery is performed, it is possible to effectively apply several surgical techniques, which in turn aim to the same thing, and that is the complete excision of the choledochal cyst with the appropriate reconstruction of biliary ducts. After reviewing the literature and bibliographic databases (PubMed, Scopus), we came to the conclusion that the short- and long-term results of surgical treatment outcomes of patients operated for choledochal cyst Type I are similar, regardless of the applied surgical technique. It can be concluded that the applied surgical technique does not affect considerably the final result of the treatment, but the period required to diagnose the disease, the extent of surgical resection, and the presence of the bile duct epithelium metaplasia, does.


**Conflict of interest:** None declared

**REFERENCES**

1. Khandelwal C, Anand U, Kumar B, Priyadarshi RN. Diagnosis and management of choledochal cysts. *Indian J Surg.* 2012;74(1):29-34 DOI: 10.1007/s12262-011-0388-1
2. Fan F, Xu DP, Xiong ZX, Li HJ, Xin HB, Zhao H, et al. Clinical significance of intrapancreatic choledochal cyst excision in surgical management of type I choledochal cyst. *J Int Med Res.* 2018;46(3):1221-9 DOI: 10.1177/0300060517728598
3. Mabrut JY, Bozio G, Hubert C, Gigot JF. Management of congenital bile duct cysts. *Dig Surg.* 2010;27(1):12-8 DOI: 10.1159/000268109
4. Badebarin D, Aslanabadi S, Teimouri-Dereshki A, Jamshidi M, Tarverdizadeh T, Shad K, et al. Different clinical presentations of choledochal cyst among infants and older children: A 10-year retrospective study. *Medicine (Baltimore).* 2017;96(17):e6679 DOI: 10.1097/MD.0000000000006679
5. Ronnekleiv-Kelly SM, Soares KC, Ejaz A, Pawlik TM. Management of choledochal cysts. *Curr Opin Gastroenterol.* 2016;32(3):225-31 DOI: 10.1097/MOG.0000000000000256
6. Vater A, Ezler CS. *Dissertio de scirrhes viscerum occasione sectionis vire typanite defunte.* Wittenburgae Pamphlets 1723.4:881
7. Machado NO, Chopra PJ, Al-Zadjali A, Younas S. Choledochal Cyst in Adults: Etiopathogenesis, Presentation, Management, and Outcome-Case Series and Review. *Gastroenterol Res Pract.* 2015;2015:602591 DOI: 10.1155/2015/602591
8. Nambiar L, Alex A, Siskind E, Shen AW, Fan C, Grimaldi G, et al. Type VI Choledochal Cyst-An Unusual Presentation of Jaundice. *Int J Angiol.* 2016;25(4):263-5 DOI: 10.1055/s-0034-1376317
9. Alonso-Lej F, Rever WB, Jr., Pessagno DJ. Congenital choledochal cyst, with a report of 2, and an analysis of 94, cases. *Int Abstr Surg.* 1959;108(1):1-30
10. Todani T, Watanabe Y, Narusue M, Tabuchi K, Okajima K. Congenital bile duct cysts: Classification, operative procedures, and review of thirty-seven cases including cancer arising from choledochal cyst. *Am J Surg.* 1977;134(2):263-9 DOI: 10.1016/0002-9610(77)90359-2
11. Ray S, Bhat BK, Yadav A, Nundy S. Isolated dilatation of the cystic duct-Type VI choledochal cyst: a rare case presentation and review of the literature. *J Surg Case Rep.* 2017;2017(4):rjx067 DOI: 10.1093/jscr/rjx067
12. Katabathina VS, Kapalczynski W, Dasyam AK, Anaya-Baez V, Menias CO. Adult choledochal cysts: current update on classification, pathogenesis, and cross-sectional imaging findings. *Abdom Imaging.* 2015;40(6):1971-81 DOI: 10.1007/s00261-014-0344-1
13. Jovanovic M, Bilanovic D, Colovic R, Grubor N, Ugljesic M. [Anomalous choledocho-pancretic ductal junction in a choledochal cyst--a case report]. *Srp Arh Celok Lek.* 2004;132(5-6):179-81 DOI: 10.2298/sarh0406179j
14. Tarallo N, Curti M, Molinelli V, Leonardi A, Fugazzola C. Diverticulum of common hepatic duct leading to obstructive jaundice, a case report. *BJR Case Rep.* 2019;5(2):20180105 DOI: 10.1259/bjrcr.20180105
15. Jablonska B. Biliary cysts: etiology, diagnosis and management. *World J Gastroenterol.* 2012;18(35):4801-10 DOI: 10.3748/wjg.v18.i35.4801
16. Xu D, Tang K, He S. A modified technique of single-incision laparoscopic hepaticojejunostomy for children with choledochal cysts. *BMC Surg.* 2019;19(1):36 DOI: 10.1186/s12893-019-0499-3
17. Ten Hove A, de Kleine RHJ, Nijkamp MW, Gouw ASH, Koopman T, Klaase JM. Robot-Assisted Laparoscopic Resection of a Todani Type II Choledochal Malformation. *Case Rep Gastroenterol.* 2019;13(2):230-7 DOI: 10.1159/000500080




**Figure 1.** Abdominal MRI – gallbladder (GB) and Type I choledochal cyst (CC) in sagittal (A) and axial (B) plane



**Figure 2.** Intraoperative photo showing hepaticojejunostomy performed using interrupted suture at about 1 cm below the primary biliary confluence

Paper accepted



**Figure 3.** Macroscopic image of the specimen, before and after opening showing multiple stones in its lumen