



Paper Accepted*

ISSN Online 2406-0895

Letter to the Editor / Писмо уреднику

Alcohol use and clinical manifestations of tuberculosis

Злоупотреба алкохола и клиничке манифестације туберкулозе

Dear Editor,

Tuberculosis (TB) remains a major cause of morbidity and mortality from infectious diseases worldwide. In 2011 there were 8.7 million new cases of active tuberculosis worldwide (13% of which involved coinfection with the human immunodeficiency virus-HIV) and 1.4 million deaths [1]. Studies have shown TB to be a stigmatized disease, which makes efforts in controlling. Improved counseling of patients and communication, patient choice of treatment support and reinforcement of supervision activities are associated with improved treatment outcomes [1]. Alcohol Use Disorders (AUDs) among tuberculosis patients are associated with nonadherence and poor treatment outcomes [1, 2]. AUD has been reported as a risk factor for an impaired immune system and increases a person's susceptibility to active TB infection, as well as to the reactivation of latent disease. Many studies have demonstrated that alcoholism has been one of the major reasons for default and mortality

* **Accepted papers** are articles in press that have gone through due peer review process and have been accepted for publication by the Editorial Board of the *Serbian Archives of Medicine*. They have not yet been copy edited and/or formatted in the publication house style, and the text may be changed before the final publication.

Although accepted papers do not yet have all the accompanying bibliographic details available, they can already be cited using the year of online publication and the DOI, as follows: the author's last name and initial of the first name, article title, journal title, online first publication month and year, and the DOI; e.g.: Petrović P, Jovanović J. The title of the article. *Srp Arh Celok Lek*. Online First, February 2017.

When the final article is assigned to volumes/issues of the journal, the Article in Press version will be removed and the final version will appear in the associated published volumes/issues of the journal. The date the article was made available online first will be carried over.

Received: September 1, 2016

Accepted: October 11, 2017

Online First: October 27, 2017

DOI: <https://doi.org/10.2298/SARH160901188S>

under the Directly Observed Tuberculosis Treatment. Persons who abuse alcohol, smoking and illicit drugs are increasingly affected by tuberculosis, with reported tuberculosis rates up to 28 times higher than the age-matched general population [3, 4, 5]. All 90 consecutive patients with active TB hospitalized from January 2015. to September 2015. at the Clinic for Pulmonary Diseases, Clinical Centre of Serbia, were included in the study. Tuberculosis was confirmed by direct microscopy (positive sputum sample for acid-fast bacili) and/or culture sputum, but rarely patohistologically. Basic demographic data, sputum sample for direct microscopy or culture, chest X-ray, alcohol use, smoking history and body mass index were collected. This study was approved by the institutional review board and patients' consents were obtained (Table 1).

Table 1. Baseline and clinical characteristics of study population (n=90).

Parameters	n(%)
Gender (male/female)	62(68.9) / 28(31.1)
Marital status (married/single/widowed/ divorced)	46(51.1) / 23(25.6) / 5(5.6) / 16(17.8)
Place of residence (urban/rural)	68(75.6) / 22(24.4)
Education (without/primary school/ secondary school /high school/college/university)	7(7.8) / 20(22.2) / 14(15.6) / 35(38.9) /5(5.6) / 9(10)
Employment (employed/unemployed/retired/ students)	43(47.8) / 29(32.2) / 16(17.8) / 2(2.2)
Alcohol consumption (never/sometimes/ ofen/excessively)	45(50) / 3(3.3) / 41(45.6) / 25(27.8)

Comparison between the groups was performed with χ^2 test. The basic and clinical variables are listed in Table 1. There was a significant difference in the distribution of the categories of chest X-Ray stages ($\chi^2=62.740$, $df=2$, $p<0.0001$) between the subjects who drank alcohol excessively compared to those who did not drink or drank occasionally. Patients who drank alcohol excessively had more extensive

changes in the lungs (25/32) compared to those who did not drink or drank occasionally. Significantly more patients with excessive alcohol consumption ad irregular nutrition in comparison with those subjects who did not drink alcohol or drank occasionally ($\chi^2=26.894$, $p<0.001$).

Hemoptysis as a symptom of serious illness had been significantly more frequent in TB patients with alcohol use (odds ratio=4.699, $p=0.040$). It was also a higher risk that TB patients who drank alcohol had extensive chest-X Ray changes at the moment of establishing the diagnosis of tuberculosis (odds ratio=63.424, $p<0.0001$). There is a strong association between AUD and TB [2, 5]. There are only a few studies which assessed alcohol use disorders in tuberculosis patients in low and middle income countries: Kazakhstan: 4% alcohol abusers; Russia: 24–62% alcohol abusers/dependents; India: 14.9–32% alcohol abusers/alcoholics; Brazil: 14–24% alcohol abusers and South Africa: 31–62% alcohol misuse [1, 5]. We found high rates of hazardous or harmful drinking habit among tuberculosis patients in Serbia, which is consistent with studies conducted in low and middle income countries. Alcohol exerts immunosuppressive effects but the exact mechanisms are difficult to define because excessive alcohol use is often associated with other comorbidities such as nutritional deficiencies, liver disease, and cigarette smoking.

It is not clear whether excess alcohol use actually leads to increased transmission of Mycobacterium tuberculosis. This study which found that excessive alcohol use was not independently associated with cavitary lesions in a multivariate analysis and tobacco use is, in univariate analysis only, may not be generalized because the researchers seropositive individuals, those patients who could not be located, and those subjects with negative sputum culture for Mycobacterium tuberculosis.

In sum, there is an evidence of a social marginalization and drift pathway on how alcohol could lead to TB. Multidisciplinary approach is essential in the treatment of these patients, and the findings suggest the importance of integrating alcohol treatment into TB care. Studies of interventions that concurrently treat excess alcohol use and tuberculosis are needed to make progress in tuberculosis elimination.

REFERENCES

1. Zumla A, Raviglione M, Hafner R, von Reyn CF. Tuberculosis. N Engl J Med. 2013; 368(8): 745–55.
2. Siemion-Szcześniak I, Kuś J. Impact of social risk factors on treatment outcome in patients with culture positive pulmonary tuberculosis (CPPTB). Pneumonol Alergol Pol. 2012; 80(5): 412–21.
3. Škodrić-Trifunović V, Mihailović-Vučinić V, Stjepanović M, Radišavljević-Pavlović S, Adžić-Vukučević T, Ilić B. Analysis of tuberculosis prevalence among cigarette smokers. Med Pregl. 2013; LXVI(Suppl 1): 84–7.
4. Golo DL, Cirić-Janković S, Santrić-Milićević M, Simić S. Alcohol use among adolescents in Serbia. Srp Arh Celok Lek. 2013; 141(3-4): 207–13.
5. Azevedo MJ, Conwill DE, Lawrence S, Jackson A, Bhuiyan AR, Hall D, et al. Tuberculosis Containment among the Homeless in Metropolitan Jackson, Mississippi. J Miss State Med Assoc. 2015; 56(8): 243–8.

*Mihailo I. STJEPANOVIĆ,
Violeta MIHAILOVIĆ-VUČINIĆ,
Jovana MAŠKOVIĆ,
Nataša ČOLOVIĆ,
Branislav GVOZDENOVIĆ,
Milica STOJKOVIĆ-LALOŠEVIĆ,
Mirjana STOJKOVIĆ*

Clinical Centre of Serbia, Belgrade, Serbia
School of Medicine, University of Belgrade, Belgrade, Serbia